



Saurashtra University

Re – Accredited Grade 'B' by NAAC
(CGPA 2.93)

Gajera, Rave V., 2010, "*Synthesis and Physicochemical Studies of some Molecules of Meducinal Interest*", thesis PhD, Saurashtra University

<http://etheses.saurashtrauniversity.edu/id/eprint/496>

Copyright and moral rights for this thesis are retained by the author

A copy can be downloaded for personal non-commercial research or study, without prior permission or charge.

This thesis cannot be reproduced or quoted extensively from without first obtaining permission in writing from the Author.

The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the Author

When referring to this work, full bibliographic details including the author, title, awarding institution and date of the thesis must be given.

Saurashtra University Theses Service
<http://etheses.saurashtrauniversity.edu>
repository@sauuni.ernet.in

Ph.D. Thesis

“SYNTHESIS AND PHYSICOCHEMICAL STUDIES OF SOME MOLECULES OF MEDICINAL INTEREST”

RAVI V. GAJERA

**DEPARTMENT OF CHEMISTRY
(DST-FIST FUNDED & UGC-SAP SPONSORED)
SAURASHTRA UNIVERSITY
RAJKOT-360 005.
GUJARAT (INDIA)
JANUARY-2010.**

**“SYNTHESIS AND PHYSICOCHEMICAL
STUDIES OF SOME MOLECULES OF
MEDICINAL INTEREST”**

**A THESIS
SUBMITTED TO THE
SAURASHTRA UNIVERSITY
FOR THE DEGREE OF**

Doctor of Philosophy

**IN
THE FACULTY OF SCIENCE (CHEMISTRY)
BY**

RAVI V. GAJERA

**UNDER THE GUIDANCE
OF**

Dr. SHIPRA BALUJA

**Department of Chemistry
(DST-FIST Funded and UGC-SAP Sponsored)
Saurashtra University
Rajkot- 360 005
Gujarat - (INDIA)**

2010



Dedicated To My
Parents &
Beloved Guide

ACKNOWLEDGEMENT

First and foremost, I wish to pay my homage and devote my emotions to "LORD GAYATRI MAA", "The Wonderful Chemist" of this lovely world without whose blessings this task would not have been accomplished. I bow my head in utter humility and complete dedication.

At the outset, I would like to extol Dr. Shipra Baluja for is advise during my doctoral research endeavor during the yester years. As my supervisor, she constantly forced me to remain focused on achieving my goal. Her observation and comments helped me to establish the overall direction of the research and to move forward expeditiously with investigation in depth. I thank her for providing me the opportunity to work with numerous local and global peers.

I owe a great deal to Dr. P. H. Parshania, Professor and Head, Department of Chemistry, who always showed deep concern and was always approachable in time to show the silver lining in every dark cloud. I will never forget Dr. A. K. Shah, Dr. V. H. Shah, Dr. H. S. Joshi, Dr. M. K. Shah, Dr. Y. T. Naliyapara, for their constant inspiration with keen interest and ever vigilant guidance without which this task could not have been achieved.

I wish to express my gratitude to Prof. S. V. Chanda, Department of Biosciences, Saurashtra University, Rajkot, for help in conducting biological activities and reviewing the manuscript of this thesis.

Above all, I bow my head with utter respect to my beloved mother Smt. Diyyaben for her continuous source of inspiration, motivation and devotion to the family, and my father Shri Vithalbhair for the uncompromising principles that guided my life. Through the stress and strain of this study. Also I can never ever forget my dear brother Rakshit. However I assure them to be worthy of whatever they have done for me.

Above all, needless to say "Thanks" to express my deep indebtedness to my seniors Dr. Nilesh Godvani, Dr. Nikunj Kachhadia, Dr. Anchal Kulshrestha, Dr. Asif Solanki, Dr. J. C. Javiya, Dr. P. K. Kasundra, and Dr. K. P. Vaishnani. They ever stood beside me with their helping hands and moral support.

I am thankful to the S.A.I.F., C.I.L., Punjab University, Chandigarh for NMR spectra, Department of Chemistry, Saurashtra University (Dr. Pankaj Kachhadia and Mr. Ram Vijay) for Mass and IR spectral data. I really very much thankful to the Saurashtra University for giving me valuable opportunity of being part of this prestigious institution. I would like to thank all whose direct and indirect support help me completing my thesis in time.

I am equally thankful to all of the non-teaching staff for extending help and cooperation.

I would like to thank my dear lab mates Rakesh, Rahul, Mehul, Nayan, Jagdish, Ashish, Sandip, Suresh, Jignesh, Pankaj, Naimish, Leena, Pooja, Amit, Renish, Bhavesh, Chirag, Axay, Nilay, Anil, Piyush, Govind, Savant, Vipul, Piyush .

I offered my gratitude to my friends Jaydeep, Piyush, Vikas, Ashish, Praful (Panu), Nilesh (Nilo), Bipin, Mepal, Vipul, Anil.

At this juncture I thank my whole family for encourage me and providing every help to fulfill my task especially My fiancée Sneha, who blessed me with their good wishes relieving all type of distress from me and fir bwing always with me continuing to boost my moral. I am gratified for their eternal love, trust, support and becoming my strongest source of motivation and inspiration. I owe a lifelong indebttness.

Ravi V. Gajera

CONTENTS

| | Page. No. |
|--|-----------|
| <u>CHAPTER-1</u> <u>GENERAL INTRODUCTION</u> | 1-5 |
| <u>CHAPTER-2</u> <u>SYNTHESIS AND CHARACTERIZATION</u> | |
| <u>PART-1</u> <u>SYNTHESIS OF PYRIMIDINE DERIVATIVES</u> | |
| SECTION-I SYNTHESIS OF 1,4-DIHYDROPYRIMIDINONES | 6-20 |
| SECTION -II SYNTHESIS OF DIHYDROPYRIMIDINETHIONES | 21-36 |
| SECTION -III SYNTHESIS OF N-METHYLPYRIMIDINES | 37-48 |
| SECTION – IV SYNTHESIS OF TETRAHYDROPYRIMIDINES | 49-63 |
| <u>PART-2</u> <u>SYNTHESIS OF BENZOTHAZOLE DERIVATIVES</u> | |
| SYNTHESIS OF AZOMETHINES | 64-81 |
| <u>CHAPTER-3</u> <u>PHYSICO CHEMICAL PROPERTIES</u> | |
| SECTION-I ACOUSTICAL PROPERTIES | 82-105 |
| SECTION -II DENSITY AND REFRACTIVE INDEX | 106-123 |
| SECTION -III CONDUCTANCE | 124-138 |
| SECTION -IV SOLUBILITY | 139-154 |
| SECTION –V THERMAL PROPERTIES | 155-171 |
| SECTION -VI DISSOCIATION CONSTANTS | 172-201 |
| <u>CHAPTER-4</u> <u>BIOLOGICAL ACTIVITIES</u> | 202-214 |
| A COMPREHENSIVE SUMMARY OF THE WORK | 215-218 |
| LIST OF PAPERS | 219-220 |

Synopsis

SYNOPSIS

"SYNTHESIS AND PHYSICOCHEMICAL STUDIES OF SOME MOLECULES OF MEDICINAL INTEREST"

Ravi V. Gajera

**Department of Chemistry
(FIST-DST funded & UGC-SAP sponsored)
Saurashtra University
Rajkot-360 005 (Gujarat)
India**

**Synopsis of the Thesis submitted to the Saurashtra
University for the degree of Doctor of Philosophy in
Chemistry**

| | | |
|------------------------------|----------|--|
| Faculty | : | Science |
| Subject | : | Chemistry |
| Title | : | Synthesis and physicochemical studies of some molecules of medicinal interest |
| Name of the Candidate | : | Ravi V. Gajera |
| Registration Number | : | 3674 |
| Date of Registration | : | 17/09/2007 |
| Name of the Guide | : | Dr. Shipra Baluja |

Place of Work

**Department of Chemistry
Saurashtra University
Rajkot-360005.
Gujarat (India)**

SYNTHESIS AND PHYSICOCHEMICAL STUDIES OF SOME MOLECULES OF MEDICINAL INTEREST.

Medicinal or pharmaceutical chemistry is a scientific discipline at the intersection of chemistry and pharmacy involved with designing, synthesizing and developing pharmaceutical drugs. It deals with identification, synthesis and development of new chemical entities suitable for therapeutic use. In recent years, many researchers have been work on medicinal chemistry which involves various aromatic compounds including heterocyclic ring systems. Heterocyclic compounds are biologically active and they have a large number of applications in medicine and pharmacy.

Taking in view of the applicability of heterocyclic organic compounds, the present work was undertaken to synthesize some new heterocycles bearing pyrimidines and thiazole nucleus.

The work to be presented in the thesis entitled "Synthesis and Physicochemical studies of some molecules of medicinal interest" is divided into four Chapters.

Chapter-1 General Introduction

Chapter-2 Synthesis and characterization

Part-1 Synthesis of Pyrimidine Derivatives

Section-I Synthesis of Dihydropyrimidinones

Section-II Synthesis of Dihydropyrimidinethiones

Section-III Synthesis of N-methyl Dihydropyrimidinones

Section-IV Synthesis of Tetrahydropyrimidines

Part-2 Synthesis of Thiazole Derivatives

Synthesis of Azomethines

Chapter-3 Physicochemical properties

Section-I Acoustical Properties

| | |
|-------------|------------------------------|
| Section-II | Density and Refractive index |
| Section-III | Conductance |
| Section-IV | Solubility |
| Section-V | Thermal Properties |
| Section-VI | Dissociation Constants |

Chapter-4 Biological activities

CHAPTER 1: General Introduction

This chapter describes literature survey of heterocyclic organic compounds, specially Pyrimidines and Thiazole derivatives. The synthesis, characterization, applications, physicochemical properties and antibacterial activities of pyrimidines and thiazole derivatives have also been discussed.


CHAPTER 2: Synthesis and Characterization

In this chapter, synthesis and characterization of various compounds are reported. It consists of two parts:


Part 1: Synthesis of Pyrimidine Derivatives

Four different types of pyrimidine Derivatives were synthesized which are given in four different sections.


Section 1: Synthesis of Dihydropyrimidinones




Section 2: Synthesis of Dihydropyrimidinethiones



Section 3: Synthesis of N-methyl Dihydropyrimidinones




Section 4: Synthesis of Tetrahydropyrimidines



Part 2: Synthesis of Azomethines

This part comprises synthesis of azomethines from benzthiazole amine.



CHAPTER 3

In this chapter, some physicochemical properties have been studied for some **dihydropyrimidinones derivatives (RVG series)** in N, N-dimethylformamide (DMF) and dimethyl sulphoxide (DMSO) solutions of different concentrations at 298.15 K. This chapter divided into six sections.

Section 1: Acoustical studies

In this section, sound velocity of some synthesized compounds (RVG series) in DMF and DMSO solutions of different concentrations were studied at 298.15 K. Further, density and viscosity also have been measured. Using these experimental data, some acoustical parameters were calculated which

are interpreted in terms of solute-solute and solute-solvent interactions in different solutions.

Section 2: Density and Refractive index

Since refractive index is a fundamental physical property of a substance, it is often used to identify a particular substance, confirm its purity, or measure its concentration. In this section, the density and refractive index of compounds of RVG series were measured in DMF and DMSO solutions at 298.15 K. Further, the density of these compounds was also evaluated theoretically and was compared with experimental values. The molar refraction and refractive index of compounds have also been calculated.

Section 3: Conductance

This section deals with the conductance measurement of solutions of compounds of RVG series in DMF and DMSO solutions over a wide range of concentration at 298.15 K. From these experimental values, specific conductance and equivalent conductance for studied compounds was measured.

Section 4: Solubility

The solubility of some derivatives of RVG series compounds in DMF and DMSO were measured by a gravimetric method at different temperatures (298.15 to 318.15 K) under atmospheric pressure. Further, some thermodynamic parameters such as Gibbs energy, enthalpy and entropy of solutions have been evaluated.

Section 5: Dissociation constants

This section deals with the dissociation constant of compounds of RVG series were studied in DMF-water mixtures at 298.15, 308.15 and 318.15 K.

From the experimental data, some thermodynamic parameters have been evaluated.

Section 6: Thermal Properties

In this section, the thermal properties of some dihydropyrimidinones derivatives were studied by TGA and DSC technique. From this data, various kinetic parameters such as energy of activation, order of degradation, frequency factor and entropy change were evaluated. The stability of various compounds was also determined from these data. Further, the melting point observed by DSC was compared with open capillary method for each derivative.

CHAPTER 4: Biological activity

In the present chapter, antibacterial activity of all the synthesized compounds was studied against some Gram positive and Gram negative bacteria in DMF and DMSO.

Signature of Guide

(Dr. Shipra Baluja)

Associate Professor,
Department of Chemistry,
Saurashtra University,
Rajkot - 360 005.

Signature of Student

(Ravi V. Gajera)

Chapter -1

General Introduction

GENERAL INTRODUCTION

The primary objective of medicinal chemistry is the design and discovery of new compounds that are suitable for use as drugs. This process involves a team of workers from a wide range of disciplines such as chemistry, biology, biochemistry, pharmacology mathematics, medicine and computing, amongst others.

The discovery or design of a new drug requires synthesis of drug, method of administration, development of tests and procedures to establish how it operates in the body and safety assessment. Drug discovery may also require fundamental research into the biological and chemical nature of the diseased state. For this, specialists from various fields including medicinal or organic chemists are needed.

The first rational development of synthetic drugs was carried out by Ehrlich and Hata who produced arsphenamine in 1910 by combining synthesis with reliable biological screening and evaluation procedures¹.

Heterocycles form by far the largest of the classical divisions of organic chemistry and are of immense importance biologically, industrially, and indeed to the functioning of any developed human society. The majority of pharmaceuticals and biologically active agrochemicals are having heterocyclic nucleus.

The heterocyclic compounds occupy a key position in the area of drugs and pharmaceuticals. Almost 80% of the drugs in clinical use are based on heterocyclic constitution because they have specific chemical reactivity.

Majority of drugs being introduced in pharmacopeias in recent years are heterocyclic compounds. A wide variety of drugs such as chlordiazepoxide (tranquillizer)^{2, 3}, imipromine (antidepressant)⁴, guanethidine (antihypertensive)⁵, indapamide (diuretic and antihypertensive)^{6,7}, ketoprofen⁸, fenoprofen and flurbiprofen penicillin⁹, cephalosporin¹⁰, norfloxacin¹¹, streptomycin^{12, 13} etc., contain heterocyclic nucleus.

Further, in industries various additives and modifiers having heterocyclic moiety are used. In addition to these, in cosmetics, reprography, information storage, plastics etc. various heterocyclic compounds have been used. Thus, for many recruits to positions in the pharmaceutical industry, a fast introduction to heterocyclic chemistry is a must.

Various plants contain heterocyclic compounds and these compounds are known to be biologically active¹⁴⁻¹⁶. Some heterocycles are fundamental to life, such as haem derivatives in blood and the chlorophylls essential for photosynthesis^{17, 18}. Further, the paired bases found in RNA and DNA are heterocycles¹⁹, sugars that in combination with phosphates²⁰ (which provide the backbones and determine the topology of these nucleic acids), some dyestuffs of plant origin include indigo blue²¹ and a poison strychnine²², obtained from the plant resin curare etc., contain heterocyclic ring. Pyrrole²³ was detected in the dry distillation of bones and pyridine²⁴ was also isolated from bone oil.

A heterocyclic nucleus is a ring in which at least one atom is not carbon. Compound containing aromatic heterocyclic nuclei have been known since the earliest studies in organic chemistry. In 1818, Brugnatelli had isolated alloxan by oxidizing of uric acid²⁵. Other derivatives of uric acid (purines and pyrimidines) were described by Wohler and Liebig²⁶.

Taking in view of the applicability of heterocyclic compounds, the present work was undertaken to synthesize some new heterocycles bearing pyrimidine and benzothiazole nucleus.

LITERATURE CITED

- (1) Ehrlich, P., Sacachiro, H. "Die Experimentelle Chemotherapie der Spirillosen." *Springer, Berlin.*, **1910**.
- (2) Cooper, J. R., Floyd, E. B., Robert, H. R. "The Complete Story of the Benzodiazepines." Oxford University Press, 7th ed., **2007**.
- (3) Skerritt, J. H., Johnston, G. A. "Enhancement of GABA binding by benzodiazepines and related anxiolytic." *Eur. J. Pharmacol.*, **1983**, 89(3-4), 193-198.
- (4) Mayo, C. P. "Pharmacology of Antidepressants" **2001**, 76.
- (5) Joyce, P. I., Rizzi, D., Calo, G., Rowbotham, D. J., Lambert D. G. "The effect of guanethidine and local anesthetics on the electrically stimulated mouse vas deferens." *Anesth. Analg.*, **2002**, 95(5), 1339-1343.
- (6) Mutschler, E. "Drug action: Basic Principles and Therapeutic aspects." *Med. Pharma Scientific Pub.*, **1995**.
- (7) Guyton and Hall; "Text book of Medical Physiology" Elsevier-Saunders (7th Ed.), **2005**.
- (8) Kantor, T. G. "Ketoprofen: a review of its pharmacologic and clinical properties." *Pharmacotherapy*, **1986**, 6(3), 93-103.
- (9) Fleming, A. "On the antibacterial action of cultures of a penicillium, with special reference to their use in the isolation of B. influenza." *Br. J. Exp. Pathol.*, **1929**, 10(31), 226-236.
- (10) Dash, C. H. "Penicillin allergy and the cephalosporins." *J. Antimicrob. Chemother.*, **1975**, 1(3), 107-118.
- (11) Goldstein, E. "Norfloxacin, a fluoroquinolone antibacterial agent, Classification, mechanism of action and in vitro activity" *Am. J. Med.*, **1987**, 82(6B), 3-17.
- (12) Kingston, W. "Streptomycin and the Balance of Credit for Discovery." *J. History Med. Allied Sci.*, **2004**, 59(3), 441-462.
- (13) Mistean, V. "The history behind the discovery of streptomycin" **2002**.
- (14) Sadykov, A. S. "Conformation states of nitrogen containing heterocyclic compounds and their biological significance." *Russ. Chem. Bull.*, **1974**, 23(5), 1129-1135.
- (15) Orhan, I., Ozcelik, B., Sener, B. "Antiviral and antimicrobial evaluation of some heterocyclic compounds from Turkish plants." *Top Heterocycl. Chem.*, **2007**, 11, 303-323.


- (16) Fuster, M. D., Alyson, E. M., Hiroto, O., Takayuki, S. "Antioxidative Activities of Heterocyclic Compounds Formed in Brewed Coffee." *J. Agri. Food Chem.*, **2000**, 48, 5600-5603.
- (17) Inman, O. L. "Studies on the chlorophylls and photosynthesis of thermal algae from Yellowstone national park, California and Nevada." *J. Gen. Physio.*, **1940**, 1, 661-666.
- (18) Mariko, K. H., Kazuhiko, S., Sakae, K. "Relationship between photosynthesis and chlorophyll content during leaf senescence of rice seedlings." *Plant Cell Physio.*, **1987**, 28(7), 1321-1329.
- (19) Gilchrist, T. L. "Heterocyclic Chemistry." 2nd edi., Longman/Wiley, Harlow/Chichester, **1992**.
- (20) Faulkner, P. "Enzymic reduction of sugar phosphates in insect blood." *Polyhydric Alcohol Phosphate Dehydrogenase*, **1956**, 64, 436-441.
- (21) Siva, R. "Status of natural dyes and dye-yielding plants in India." *Curr. Sci.*, **2007**, 92(7), 916-925.
- (22) Wilderness, C. "Protection of poison strychnine." *Spring.*, **2009**, 4.
- (23) Runge, F. F. *Ann. Physk.*, **1834**, 31, 67.
- (24) Anderson, T. *Trans. Royal Soc., Edinburgh*, **1849**, 16, 123.
- (25) Brugnatelli, di. *Giornale di fisica, chemica, eec.*, **1818**, 38, 117.
- (26) Liebig, J. V., Wohler, F. "Untersuchungen uber die Natur der Harnsaure." **1838**.

GENERAL REMARKS

1. Melting points were recorded by open capillary method and are uncorrected.
2. Infrared spectra were recorded on SHIMADZU FTIR-8400 (Diffuse reflectance attachment) in the frequency range of 4000-400 cm^{-1} using KBr. Spectra were calibrated against the polystyrene absorption at 1610 cm^{-1} .
3. ^1H NMR spectra were recorded on BRUKER AVANCE II 400 spectrometer. Making a solution of samples in DMSO d_6 and CDCl_3 solvents using tetramethylsilane (TMS) as the internal standard unless otherwise mentioned, and are given in the δ scale.
4. Mass spectra were recorded on SHIMADZU GCMS-QP2010 spectrometer operating at 70 eV using direct injection probe technique.
5. Analytical thin layer chromatography (TLC) was performed on Merck-precoated silica gel-G F254 aluminium plates. Visualization of the spots on TLC plates was achieved either by exposure to iodine vapor or UV light.
6. The chemicals used for the synthesis of intermediates and end products were purchased from Spectrochem, Sisco Research Laboratories (SRL), Thomas-Baker, Sd fine chemicals and Loba chemie.
7. Ultrasonic interferometer was carried Mittal Enterprise, New Delhi, Model No. F-81) working at frequency of 2 MHz was used to determine sound velocity, with the uncertainty of 0.01%.
8. The structures and names of all the compounds given in the experimental section were generated using ACD ChemsSketch version 12.0.


Chapter -2

Synthesis and Characterization



Part-1

Synthesis of Pyrimidine Derivatives



Section-I


Synthesis of Dihydropyrimidinones

INTRODUCTION

Multicomponent reactions (MCRs) are of increasing importance in organic and medicinal chemistry. Over the last decade, industrial and academic researchers have made such powerful MCR strategies into one of the most efficient and cost-effective tools for combinatorial and parallel synthesis.¹ One prominent MCR that produces an interesting class of nitrogen heterocycles is the venerable Biginelli dihydropyrimidine synthesis.² Thus, multicomponent reaction is of much importance due to excellent pharmacological properties of dihydropyrimidines.

The nitrogen containing heterocycles are an important class of compounds, contributed to the society from biological and industrial point of view which helps to understand life processes³. Dihydropyrimidines are known to possess therapeutic and pharmacological properties² such as in vitro activity against unrelated DNA and RNA, viruses including polio herpes viruses, diuretic, antitumor, anti HIV, cardiovascular⁴.

Several alkaloids isolated from marine sources exhibit interesting biological properties^{5, 6}. These isolated alkaloids contain the dihydropyrimidine unit. Some of these isolated marine alkaloids containing dihydropyrimidinones (eg. betzelladine alkaloids) are known to be a potent HIVgp-120-CD4 inhibitors⁷⁻⁹. The compound nitractin(A) i.e., a dihydropyrimidine derivative exhibited anti viral activity¹⁰. 4-Aryl dihydropyrimidines e.g. nifedipine (B) are used as calcium channel modulars, which is also used for the treatment of cardiovascular diseases¹¹. Some of the analogues were screened as anti tumor agents. Pyrimidine -5-carboxamide of type (C) was reported to possess anti carcinogenic activity, anti inflammatory¹² and analgesic¹³ activities.



There are various methods for the synthesis of dihydropyrimidines. Many researchers have reported the use of different catalysts in the synthesis of dihydropyrimidines. Recently, various new methods for the synthesis of these compounds have been developed to improve the efficiency of the Biginelli reaction, such as use of heteropoly acids¹⁴, CuCl₂-H₂O¹⁵, l-proline¹⁶, praseodymium methanesulfonate¹⁷, chloroacetic acid¹⁸, etidronic acid¹⁹, zinc methanesulfonate²⁰ etc. The search for the new, readily available, and green catalysts is still being actively pursued.

Because of the great potential of room-temperature ionic liquids as environmentally benign media for catalytic processes, much attention has currently been focused on the organic reactions catalyzed with or in ionic liquids. Various organic reactions promoted with acid-base catalysts are now done in ionic liquids.²¹ In recent years, the acidic ionic liquids have also been used as catalysts for the Biginelli reaction²²⁻²⁴. Reddy and coworker²⁵ have reported the synthesis of dihydropyrimidines by using Zirconium chloride catalyst. Guo and salehi²⁶ have synthesized dihydropyrimidinones using magnesium bromide as a catalyst under solvent free condition.

Very recently, for novel Biginelli-like scaffold syntheses²⁷, the common open-chain b-dicarbonyl compounds has been used to cyclic b-diketones²⁸, b-ketolactones²⁹, cyclic b-diester³⁰ or b-diamides^{31, 32}, benzocyclic ketones and a-ketoacids³³. These reactions suffer from limitations such as low yields, very long reaction times, harsh reaction conditions and unrecoverable strong acids.

Thus, owing to the biological importance of dihydropyrimidinone derivatives, the present chapter describes the synthesis and characterization of some 1,4-dihydropyrimidinone derivatives.

EXPERIMENTAL

Synthesis of 4-(3,4-dimethoxyphenyl)-6-methyl-2-oxo-N-phenyl-1,2,3,4-tetrahydropyrimidine-5-carboxamide.

[A] Synthesis of 3, 4-dimethoxybenzaldehyde:

An aqueous solution (20 ml) of vaniline (0.01M) was refluxed at 100 °C for half an hour with stirring. To this solution, few drops of NaOH and 2 ml of dimethylsulphate (DMS) were added slowly and again the reaction mixture was stirred with refluxed for 3 to 3.5 hrs. Then, diethyl ether was added to the reaction mixture and solvent was allowed to evaporate. The crude product was isolated and crystallized from absolute ethanol.


[B] Synthesis of 3-oxo-N-phenylbutanamide:

A mixture of substituted aniline (0.01 M) and ethylacetoacetate (0.01 M) in 40 ml toluene was refluxed for 12 hours in presence of few drops of slurry of NaOH in water. The reaction was monitoring with TLC. The excess toluene was distilled out and the reaction mixture was taken in hexane and stirred with glass rod. The products were isolated in hexane and filtered, dried. The crude product was taken in aqueous NaOH solution and neutralized with dilute HCl. The recrystallisation was done in ethanol.


[C] Synthesis of 4-(3,4-dimethoxyphenyl)-6-methyl-2-oxo-N-phenyl-1,2,3,4-tetrahydropyrimidine-5-carboxamide:

A mixture of 3, 4 dimethoxy aldehyde (0.01 M), substituted diketones (0.01 M) and urea (0.015 M) in methanol was refluxed for 12 hours in presence of few drops of concentrated HCl as catalyst. The product was isolated and crystallized from ethanol. All the synthesized compounds were recrystallized from ethanol.

[A] Synthesis of 3, 4-dimethoxybenzaldehyde:



[B] Synthesis of 3-oxo-N-phenylbutanamide:



[C] Synthesis of 4-(3,4-dimethoxyphenyl)-6-methyl-2-oxo-N-phenyl-1,2,3,4-tetrahydropyrimidine-5-carboxamide:




Table 2.1.1: Physical constants of Dihydropyrimidenons.

| Sr. No. | Code | R | M.F. | M. Wt. (g/mol) | R _f * Value | M.P. °C | Yield % |
|---------|--------|---------------------|---|-------------------|---------------------------|------------|------------|
| 1 | RVG-1 | 4-OCH ₃ | C ₂₁ H ₂₃ N ₃ O ₅ | 397 | 0.61 | 168 | 53 |
| 2 | RVG-2 | 4-CH ₃ | C ₂₁ H ₂₃ N ₃ O ₄ | 381 | 0.54 | 152 | 57 |
| 3 | RVG-3 | 4-Cl | C ₂₀ H ₂₀ ClN ₃ O ₄ | 401 | 0.49 | 179 | 46 |
| 4 | RVG-4 | 2- CH ₃ | C ₂₁ H ₂₃ N ₃ O ₄ | 381 | 0.47 | 182 | 62 |
| 5 | RVG-5 | 3- OCH ₃ | C ₂₁ H ₂₃ N ₃ O ₅ | 397 | 0.54 | 164 | 45 |
| 6 | RVG-6 | 4-F | C ₂₀ H ₂₀ FN ₃ O ₄ | 385 | 0.71 | 160 | 59 |
| 7 | RVG-7 | 2,5 – di Cl | C ₂₀ H ₁₉ Cl ₂ N ₃ O ₄ | 436 | 0.37 | 171 | 72 |
| 8 | RVG-8 | 3-Cl | C ₂₀ H ₂₀ ClN ₃ O ₄ | 401 | 0.62 | 198 | 62 |
| 9 | RVG-9 | 3,4 – di Cl | C ₂₀ H ₁₉ Cl ₂ N ₃ O ₄ | 436 | 0.66 | 169 | 77 |
| 10 | RVG-10 | 3-Cl, 4-F | C ₂₀ H ₁₉ ClFN ₃ O ₄ | 419 | 0.52 | 145 | 78 |

*Chloroform : Methanol: 9:1

The various physical constants such as R_f value, melting point and percentage of yield for all the synthesized 1,4-dihydropyrimidinone derivatives are given in Table 2.1.1. The characterization was done by IR, ^1H NMR and mass spectra.

Infrared spectra:

The IR spectra were recorded by SHIMADZU-FTIR-8400 Spectrophotometer in the frequency range of $4000\text{--}400\text{ cm}^{-1}$ by KBr powder method. Figure 2.1.1 shows IR spectra of RVG-5. The IR spectral data for RVG-5 is given in Table 2.1.2. The spectral data for all other compounds are reported in Table 2.1.3.

^1H NMR Spectra:

The NMR spectra were recorded by BRUKER Spectrometer (400 MHz) using internal reference TMS and solvent $\text{CDCl}_3/\text{DMSO}$. Figure 2.1.2 shows NMR spectra of RVG-5. The spectral data for RVG-5 is given in Table 2.1.4.

Mass spectra:

The Mass spectra were recorded by GCMS-SHIMADZU-QP2010. Figure 2.1.3 shows mass spectra of RVG-5. The proposed mass fragmentation of the same compound is also given in Scheme 2.1.1.

Figure 2.1.1: IR spectra of 4-(3,4-dimethoxyphenyl)-N-(2-methoxyphenyl)-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide (RVG-5).




Table 2.1.2: IR spectral data of 4-(3,4-dimethoxyphenyl)-N-(2-methoxyphenyl)-6-methyl-2-oxo-1,2,3,4-tetrahydro pyrimidine-5-carboxamide (RVG-5).

| Type | Vibration mode | Frequency in cm^{-1} | |
|----------|--------------------|-------------------------------|-----------|
| | | Observed | Reported |
| Alkane | C-H str. (asym.) | 2941.54 | 2975-2900 |
| | C-H str. (sym.) | 2837.38 | 2880-2810 |
| | C-H def. (asym.) | 1454.02 | 1480-1435 |
| | C-H def. (sym.) | 1330.93 | 1985-1350 |
| Aromatic | C-H str. | 3080.42 | 3100-3000 |
| | C=C str. | 1518.03 | 1585-1480 |
| | C-H i.p. def. | 1128.39 | 1125-1090 |
| | C-H o.o.p. def. | 796.63 | 860-810 |
| Ketones | C=O str.(cyclic) | 1703.20 | 1740-1680 |
| | C=O str. (alip.) | 1670.41 | 1710-1650 |
| Nitrogen | C-N str. | 1249.91 | 1350-1200 |
| | N-H str. | 3275.24 | 3400-3200 |
| | N-H def. | 1523.82 | 1650-1500 |
| Ether | C-O-C str. (asym.) | 1161.19 | 1400-1000 |
| | C-O-C str. (sym.) | 1028.09 | 1075-1020 |

Table 2.1.3: IR spectral data of synthesized 1,4-Dihydropyrimidinones.

| Compounds | <i>IR ν, (cm⁻¹)</i> | | | | | |
|---------------|---|----------------|---------------|---------------|----------------|----------|
| | C=O (cyclic) | C=O (alip.) | N-H (Str.) | C=C (Str.) | C-H (asym.) | R |
| RVG-1 | 1710.27 | 1674.33 | 3342.20 | 1517.61 | 2912.31 | 1145.12 |
| RVG-2 | 1712.34 | 1674.38 | 3294.16 | 1516.54 | 2935.28 | 2879.30 |
| RVG-3 | 1711.30 | 1676.46 | 3279.17 | 1539.21 | 2946.46 | 759.59 |
| RVG-4 | 1716.16 | 1681.67 | 3261.14 | 1518.08 | 2943.42 | 2839.50 |
| RVG-6 | 1715.24 | 1664.59 | 3321.26 | 1537.03 | 2953.27 | 689.46 |
| RVG-7 | 1722.22 | 1673.51 | 3267.20 | 1529.55 | 2955.21 | 749.12 |
| RVG-8 | 1720.08 | 1659.52 | 3309.49 | 1520.43 | 2912.55 | 762.20 |
| RVG-9 | 1711.09 | 1666.45 | 3261.34 | 1514.37 | 2967.24 | 758.12 |
| RVG-10 | 1707.10 | 1678.51 | 3286.51 | 1502.36 | 2972.33 | 671, 754 |

Figure 2.1.2: ^1H NMR spectra of 4-(3,4-dimethoxyphenyl)-N-(2-methoxyphenyl)-6-methyl-2-oxo-1,2,3,4-tetrahydro pyrimidine-5-carboxamide (RVG-5).







Table 2.1.4: ^1H NMR spectral data of 4-(3,4-dimethoxyphenyl)-N-(2-methoxyphenyl)-6-methyl-2-oxo-1,2,3,4-tetrahydro pyrimidine-5-carboxamide (RVG-5).




| Singal No. | Signal Position (δ ppm) | Relative No. of Protons | Multiplicity | Inference | J (Hz) |
|------------|---------------------------------|-------------------------|--------------|----------------------|----------|
| 1 | 2.24 | 3H | singlet | $-\text{CH}_a$ | - |
| 2 | 3.57 | 3H | singlet | $-\text{OCH}_b$ | - |
| 3 | 3.87 | 6H | singlet | $-\text{OCH}_{cc'}$ | - |
| 4 | 5.57 | 1H | singlet | H_d | - |
| 5 | 6.72-6.93 | 6H | multiplat | $-\text{Ar-H}_{e-j}$ | - |
| 6 | 7.21 | 1H | singlet | $-\text{Ar-H}_k$ | - |
| 7 | 7.61 | 1H | singlet | $-\text{NH}_l$ | - |
| 8 | 8.08 | 1H | singlet | $-\text{NH}_m$ | - |
| 9 | 8.71 | 1H | singlet | $-\text{NH}_n$ | - |

Figure 2.1.3: Mass spectra of 4-(3,4-dimethoxyphenyl)-N-(2-methoxyphenyl)-6-methyl-2-oxo-1,2,3,4-tetrahydro pyrimidine-5-carboxamide (RVG-5).



Scheme 2.1.1: Proposed mass fragmentation of 4-(3,4-dimethoxyphenyl)-N-(2-methoxyphenyl)-6-methyl-2-oxo 1,2,3,4-tetrahydropyrimidine-5-carboxamide (RVG-5).




LITERATURE CITED

- (1) Bienayme, H., Hulme, C., Odon, G., Schmitt, P. "Maximizing synthetic efficiency: multi-component transformations lead the way." *Chem. Eur. J.* **2000**, 6, 3321-3329.
- (2) Biginelli, P. "The condensation reaction described by Biginelli". *Gazz. Chim. Ital.*, **1893**, 23, 360-416.
- (3) Mentel, M., Schmidt, A. M., Gorray, M., Eilbracht, P., Breinbauer, R. "Polystyrene Sulfonyl Chloride: A Highly Orthogonal Linker Resin for the Synthesis of nitrogen-containing Heterocycles." *Angewandte Chemie*, **2009**, 48(32), 5841-5844
- (4) Kappe, C. O. "100 Years of the Biginelli dihydropyrimidine synthesis." *Tetrah.*, **1993**, 49, 6937-6963.
- (5) Atwal, K. S., Rovnyak, G. C., O'Reilly, B. C., Schwartz, J. "Substituted 1,4-dihydropyrimidines. 3. Synthesis of selectivity functionalized 2-hetero-1,4-dihydropyrimidines." *J. Org. Chem.*, **1989**, 54, 5898.
- (6) Kappe, C. O., Fabian, W. M. F., Semones, M. A. "Conformational analysis of 4-aryl-dihydropyrimidine calcium channel modulators. A comparison of ab initio, semiempirical and x-ray crystallographic studies." *Tetrah.* **1997**, 53, 2803.
- (7) Patil, A. D., Kumar, N. V., Kokke, W. C., Bean, M. F., Freyer, A. J., De Brosse, C., Mai, S., Truneh, A., Faulkner, D. J., Crate, B., Breen, A. L., Hertzberg, R. P., Johnson, R. K., Westley, J. W., Potts, B. C. M. "Novel alkaloids from the sponge Batzella sp.: inhibitors of HIV gp120-human CD4 binding." *J. Org. Chem.*, **1995**, 60, 1182-1185.
- (8) Snider, B. B., Chen, J., Patil, A. D., Freyer, A. "Synthesis of the tricyclic portions of batzelladines A, B and D. Revision of the stereochemistry of batzelladines A and D." *Tetrah. Lett.*, **1996**, 37, 6977-6981.
- (9) Rama Rao, A. V., Gujar, M. K., Vasudevan, J. "An enantiospecific synthesis of the tricyclic guanidine segment of the anti-HIV marine alkaloid batzelladine A." *J. Chem. Soc., Chem. Commun.*, **1995**, (13), 1369-1374.
- (10) Antoni 'n, H., Ivan, V., Milena, M., Graciela, A., Robert, S., Lieve, N., Erik, De C., Jan, B. "6-[2-(Phosphonomethoxy)alkoxy]pyrimidines with Antiviral Activity." *J. Med. Chem.* **2002**, 45, 1918-1929.
- (11) Birgit, J., Tetiana, P., Kappe, C. O. "Design and synthesis of a conformationally rigid mimic of the dihydropyrimidine calcium channel modulator SQ 32,926." *Molecules.*, **2000**, 5(3), 227-239.

- (12) Kappe, C. O., Farber, G., "Thermolysis and photolysis of 6-diazidomethyl-1,2,3,4-tetrahydro-2-oxopyrimidine-5-carboxylates." *J. Chem. Soc., Perkin Trans.*, **1991**, (5), 1342-1344.
- (13) Sadanandam, Y. S., Shetty, M. M., Diwan, P. V. "Synthesis and biological evaluation of new 3,4-dihydro-6-methyl-5-N-methylcarbamoyl-4-(substituted phenyl)-2(1H)pyrimidinones and pyrimidinethiones." *Eur. J. Med. Chem.*, **1992**, 27, 87-92.
- (14) Romanelli, G. P., Sathicq, A. G., Autino, J. C., Baronetti, G., Thomas, H. J. "Solvent-free approach to 3,4-dihydropyrimidin-2(1H)-(thio)ones: Biginelli reaction catalyzed by a Wells–Dawson reusable heteropolyacid." *Synth. Commun.*, **2007**, 37(22), 3907–3916.
- (15) Xu, F., Wang, J., Tian, Y. "New procedure for one-pot synthesis of 3,4-dihydropyrimidin-2(1H)-ones by Biginelli reaction." *Synth. Commun.* **2008**, 38(8), 1299–1310.
- (16) Mabry, J., Ganem, B. "Studies on the Biginelli reaction: A mild and selective route to 3,4-dihydropyrimidin-2(1H)-ones via enamine intermediates." *Tetrah. Lett.*, **2006**, 47, 55–56.
- (17) Wang, M., Song, Z., Gong, H. "Praseodymium methanesulfonate-catalyzed one-pot synthesis of 3,4-dihydropyrimidin-2-(1H)-ones." *Prep. Bio. Biotech.*, **2008**, 38 (1), 105–114.
- (18) Yu, Y., Liu, D., Liu, C., Luo, G. "One-pot synthesis of 3,4-dihydropyrimidin-2(1H)-ones using chloroacetic acid as catalyst." *Biol. Med. Chem. Lett.*, **2007**, 17, 3508–3510.
- (19) Savant, M. M., Pansuriya, A. M., Bhuva, C. V., Kapuriya, N. P., Naliyapara, V. T. " Etidronic acid: a new and efficient catalyst for the synthesis of novel 5-nitro-3,4dihydropyrimidin-2(1H)-one." *Catal. Lett.*, **2009**, 132, 281-284.
- (20) Wang, M., Jiang, H., Wang, Z. "Biginelli condensation of aliphatic aldehydes catalysed by zinc methanesulfonate." *J. Chem. Res.*, **2005**, 691–693.
- (21) Suman, L. J., Jomy, K. J., Bir S. "Ionic liquid promoted an improved synthesis of 3,4-dihydropyrimidinones using [bmim]BF₄ immobilized Cu (II) acetylacetonate as recyclable catalytic system." *Cata. Lett.*, **2007**, 115, 1–2.
- (22) Fang, D., Luo, J., Zhou, X., Ye, Z., Liu, Z. "One-pot green procedure for Biginelli reaction catalyzed by novel task specific room-temperature ionic liquids." *J. Mol. Catal. Chem.*, **2007**, 274, 208–211.

- (23) Zheng, R., Wang, X., Xu, H., Du, J. "Bronsted acidic ionic liquid: An efficient and reusable catalyst for the synthesis of 3,4-dihydropyrimidin-2(1H)-ones." *Synth. Commun.*, **2006**, 36 (11), 1503-1513.
- (24) Liu, D., Gui, J., Zhu, X., Song, L., Sun, Z. "Synthesis and characterization of task-specific ionic liquids possessing two Bronsted acid sites." *Synth. Commun.*, **2007**, 37, 759-765.
- (25) Reddy, C. V., Mahesh, M., Raju, P. V. K., Babu, T. R., Reddy, V. V. N. "Zireohium (IV) chloride-catalyzed one-pot synthesis of 3,4-dihydropyrimidin-2(1H)-ones." *Tetrah. Lett.*, **2002**, 43, 2657-2659.
- (26) Salehi, H. Guo, Q. X. "A facile and efficient one-pot synthesis of dihydropyrimidinones catalyzed by magnesium bromide under solvent-free conditions." *Synth. Commun.*, **2004**, 34(1), 171-179.
- (27) Kappe, C. O. "Recent Advances in the Biginelli Dihydropyrimidine Synthesis. New Tricks from an Old Dog." *Acc. Chem. Res.*, **2000**, 33(12), 879-888.
- (28) Yarim, M., Sarac, S., Kilic, F. S., Erol, K. "Synthesis and in vitro calcium antagonist activity of 4-aryl-7,7-dimethyl/1,7,7-trimethyl-1,2,3,4,5,6,7,8-octahydroquinazoline-2,5-dione derivatives." *Farmaco*, **2003**, 58(1), 17-24.
- (29) Byk, G., Gettlieb, H. E., Herscovici, J., Mirkin, F. "New Regioselective Multicomponent Reaction: One Pot Synthesis of Spiro Heterobicyclic Aliphatic Rings." *J. Comb. Chem.*, **2000**, 2(6), 732-735.
- (30) Shaabani, A., Bazgir, A., Bijanzadeh, H. R. "A reexamination of Biginelli-like multicomponent condensation reaction: one-pot regioselective synthesis of spiro heterobicyclic rings." *Mol. Diver.*, **2004**, 8(2), 141-145.
- (31) Paluchowska, M. H., Bojarski, A. J., Bugno, R., Charakchieva, M. S. Wesolowska, A. "Modification of the structure of 4, 6-disubstituted 2-(4-alkyl-1-piperazinyl)pyridines: synthesis and their 5-HT_{2A} receptor activity." *Archiv der Pharm.*, **2003**, 336(2), 104-110.
- (32) Mamaev, V. P., Borovik, V. P. "Pyrimidines, formation of spiro compounds in reactions of barbituric acid with arylidenebisureas." *Seriya Khimicheskikh Nauk.*, **1968**, (5), 83-87.
- (33) Abelman, M. M., Smith, S. C., James, D. R. "Cyclic ketones and substituted α -keto acids as alternative substrates for novel Biginelli-like scaffold syntheses." *Tetrah. Lett.*, **2003**, 44(24), 4559-4562.



Section-II

Synthesis of Dihydropyrimidinethiones

INTRODUCTION

The chemistry of pyrimidinethiones and its derivatives have been studied since past century due to their pharmacological association with diverse pharmacological properties. Pyrimidine was first isolated by Gabriel and Colman in 1899. Though pyrimidine itself does not exist in nature but substituted pyrimidines containing pyrimidine moiety are found as a part of more complex systems and are widely distributed.

The synthesis of dihydropyrimidinethiones derivatives by the Biginelli reaction is important because of their pharmacological profiles¹⁻⁷. Further, various fused bicyclic systems can also be developed from this precursor⁸. Various modified building blocks, e.g. β -ketoesters, ureas/thioureas, and aldehydes, are used to synthesize molecules with targeted substitutions on a basic skeleton. A large number of catalysts⁹⁻¹⁵ are reported to catalyze the reaction efficiently and to provide higher yields. Different methods of synthesis, e.g., microwave-assisted synthesis¹⁶⁻¹⁸, solid phase synthesis¹⁹, and use of ionic liquids²⁰, have also been investigated.

The first primary synthesis of pyrimidinethione derivatives from aliphatic fragments was carried out by Frankland and Kolbs in 1848. Since then, many distinct primary synthetic methods have been devised.

Pyrimidinethiones represent one of the most active class of compounds possessing a wide spectrum of biological activities, such as antiviral and antitumor^{21, 22}, anti-inflammatory and analgesic²³, antifilarial²⁴, anticancer and herbicidal^{25, 26}, antineoplastic²⁷, anti HIV²⁸, antitubercular²⁹, antagonist³⁰ antimicrobial^{31, 32} antibacterial³³ etc.

Abdel-Rahman et al.³⁴ have synthesized pyrimidinethione derivatives and reported them as potent antimicrobial agent. Recently, Kern et al.³⁵ have reported the activity against orthopoxvirus infection of pyrimidinethione derivatives. Liu and co-workers³⁶ have reported the highly enantioselective organocatalytic thiopyrimidines. The inhibitory activities against HIV virus of some modified thiopyrimidine nucleoside have also been reported³⁷. The antimicrobial activity of some pyrimidinethiones has also been studied^{38, 39}.

Thus, the important role played by pyrimidinethiones nucleus for various physiological activities prompted us to synthesis pyrimidinethiones derivatives.

EXPERIMENTAL

Synthesis of 4-(3,4-dimethoxyphenyl)-6-methyl-N-phenyl-2-thioxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide

[A] Synthesis of 3, 4-dimethoxybenzaldehyde:

An aqueous solution (20 ml) of vaniline (0.01M) was refluxed at 100 °C for half an hour with stirring. To this solution, few drops of NaOH and 2 ml of dimethylsulphate (DMS) were added slowly and again the reaction mixture was refluxed for 3 to 3.5 hrs with stirring. Then, diethyl ether was added to the reaction mixture and solvent was allowed to evaporate. The crude product was isolated and crystallized from absolute ethanol.


[B] Synthesis of 3-oxo-N-phenylbutanamide:

A mixture of substituted aniline (0.01 M) and ethylacetoacetate (0.01 M) in 40 ml toluene was refluxed for 12 hours in presence of few drops of slurry of NaOH in water. The reaction was monitoring with TLC. The excess toluene was distilled out and the reaction mixture was taken in hexane and stirred with glass rod. The products were isolated in hexane and filtered, dried. The crude product was taken in aqueous NaOH solution and neutralized with dilute HCl. The recrystallisation was done in ethanol.


[C] Synthesis of 4-(3,4-dimethoxyphenyl)-6-methyl-N-phenyl-2-thioxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide:

A mixture of 3, 4 dimethoxy aldehyde (0.01 M), 3-oxo-N-phenylbutanamide (0.01 M) and thiourea (0.012 M) in methanol was refluxed for 12 hours in presence of few drops of concentrated HCl as catalyst. The product was isolated and crystallized from ethanol. All the synthesized compounds were recrystallized from ethanol.

[A] Synthesis of 3, 4-dimethoxybenzaldehyde:



[B] Synthesis of 3-oxo-N-phenylbutanamide:



[C] Synthesis of 4-(3,4-dimethoxyphenyl)-6-methyl-N-phenyl-2-thioxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide:




Table 2.2.1: Physical constants of Dihydropyrimidenethiones.

| Sr. No. | Code | R | M.F. | M. Wt. (g/mol) | R _f * Value | M.P. °C | Yield % |
|---------|--------|---------------------|---|-------------------|---------------------------|------------|------------|
| 1 | SRG-1 | 4-OCH ₃ | C ₂₁ H ₂₃ N ₃ O ₄ S | 413 | 0.49 | 211 | 63 |
| 2 | SRG-2 | 4-CH ₃ | C ₂₁ H ₂₃ N ₃ O ₃ S | 397 | 0.53 | 194 | 56 |
| 3 | SRG-3 | 4-Cl | C ₂₀ H ₂₀ ClN ₃ O ₃ S | 417 | 0.46 | 224 | 66 |
| 4 | SRG-4 | 2- CH ₃ | C ₂₁ H ₂₃ N ₃ O ₃ S | 397 | 0.49 | 232 | 61 |
| 5 | SRG-5 | 3- OCH ₃ | C ₂₁ H ₂₃ N ₃ O ₄ S | 413 | 0.62 | 215 | 75 |
| 6 | SRG-6 | 4-F | C ₂₀ H ₂₀ FN ₃ O ₃ S | 401 | 0.70 | 248 | 59 |
| 7 | SRG-7 | 2,5 - di Cl | C ₂₀ H ₁₉ Cl ₂ N ₃ O ₃ S | 452 | 0.59 | 251 | 70 |
| 8 | SRG-8 | 3-Cl | C ₂₀ H ₂₀ ClN ₃ O ₃ S | 417 | 0.82 | 219 | 65 |
| 9 | SRG-9 | 3,4 - di Cl | C ₂₀ H ₁₉ Cl ₂ N ₃ O ₃ S | 452 | 0.63 | 263 | 67 |
| 10 | SRG-10 | 3-Cl, 4-F | C ₂₀ H ₁₉ ClFN ₃ O ₃ S | 435 | 0.70 | 201 | 72 |

* Chloroform:Methanol:- 9:1

The various physical constants such as R_f value, melting point and percentage of yield for all the synthesized dihydropyrimidinethione derivatives are given in Table 2.2.1. The characterization was done by IR, ^1H NMR and mass spectra.

Infrared spectra:

The IR spectra were recorded by SHIMADZU-FTIR-8400 Spectrophotometer in the frequency range of $4000\text{--}400\text{ cm}^{-1}$ by KBr powder method. Figure 2.2.1 shows IR spectra of SRG-1. The IR spectral data for SRG-1 is given in Table 2.2.2. The spectral data for all other compounds are reported in Table 2.2.3.

^1H NMR Spectra:

The NMR spectra were recorded by BRUKER Spectrometer (400 MHz) using internal reference TMS and solvent $\text{CDCl}_3/\text{DMSO}$. Figure 2.2.2 shows NMR spectra of SRG-1. The spectral data for SRG-1 is given in Table 2.2.4.

Mass spectra:

The Mass spectra were recorded by GCMS-SHIMADZU-QP2010. Figure 2.2.3 shows mass spectra of SRG-1. The proposed mass fragmentation of the same compound is also given in Scheme 2.2.1.

Figure 2.2.1: IR spectra of 4-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-6-methyl-2-thioxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide (SRG-1).




Table 2.2.2: IR spectral data of 4-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-6-methyl-2-thioxo-1,2,3,4-tetrahydro pyrimidine-5-carboxamide (SRG-1).

| Type | Vibration mode | Frequency in cm ⁻¹ | |
|----------|----------------------|-------------------------------|-----------|
| | | Observed | Reported |
| Alkane | C-H str. (asym.) | 2937.68 | 2975-2900 |
| | C-H str. (sym.) | 2843.17 | 2880-2810 |
| | C-H def. (asym.) | 1452.45 | 1480-1435 |
| | C-H def. (sym.) | 1336.71 | 1985-1350 |
| Aromatic | C-H str. | 2995.55 | 3100-3000 |
| | C=C str. | 1514.17 | 1585-1480 |
| | C-H i.p. def. | 1134.18 | 1125-1090 |
| | C-H o.o.p. def. | 756.12 | 860-810 |
| Sulfur | C=S str.(cyclic) | 1134.18 | 1050-1200 |
| Ketones | C=O str. (aliphatic) | 1637.62 | 1710-1650 |
| Nitrogen | C-N str. | 1301.99 | 1350-1200 |
| | N-H str. | 3350 | 3400-3200 |
| | N-H def. | 1539. | 1650-1500 |
| Ether | C-O-C str. (asym.) | 1195.91 | 1400-1000 |
| | C-O-C str. (sym.) | 1031.95 | 1075-1020 |

Table 2.2.3: IR spectral data of synthesized Dihydropyrimidenethiones.

| Compounds | <i>IR ν, (cm^{-1})</i> | | | | | |
|---------------|--|-----------------------|-----------------------|-----------------------|------------------------|----------|
| | C=S (cyclic) | C=O (Str.) | N-H (Str.) | C=C (Str.) | C-H (asym.) | R |
| SRG-2 | 1146.21 | 1697.67 | 3369.11 | 1516.47 | 2933.23 | 2837.22 |
| SRG-3 | 1189.54 | 1660.71 | 3252.27 | 1500.06 | 2995.20 | 734.60 |
| SRG-4 | 1207.34 | 1699.39 | 3314.21 | 1524.08 | 2964.37 | 2861.54 |
| SRG-5 | 1147.58 | 1676.24 | 3217.54 | 1516.01 | 2935.48 | 1145.10 |
| SRG-6 | 1094.56 | 1702.57 | 3285.60 | 1522.67 | 2956.54 | 637.45 |
| SRG-7 | 1231.79 | 1688.19 | 3299.83 | 1531.57 | 2961.97 | 736.19 |
| SRG-8 | 1161.15 | 1710.49 | 3337.70 | 1529.33 | 2919.80 | 764.70 |
| SRG-9 | 1101.61 | 1689.56 | 3250.34 | 1543.24 | 2943.45 | 782.45 |
| SRG-10 | 1177.57 | 1691.34 | 3257.10 | 1527.49 | 2964.20 | 661, 743 |






Table 2.2.4: ^1H NMR spectral data of 4-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-6-methyl-2-thioxo-1,2,3,4-tetrahydro pyrimidine-5-carboxamide (SRG-1).




| Singal No. | Signal Position (δ ppm) | Relative No. of Protons | Multiplicity | Inference | J (Hz) |
|------------|---------------------------------|-------------------------|--------------|-------------------------|----------|
| 1 | 2.12 | 3H | singlet | $-\text{CH}_a$ | - |
| 2 | 2.27 | 3H | singlet | $-\text{OCH}_b$ | - |
| 3 | 3.74 | 3H | singlet | $-\text{OCH}_c$ | - |
| 4 | 3.77 | 3H | singlet | $-\text{OCH}_d$ | - |
| 5 | 6.91 | 3H | multiplet | Ar- $\text{H}_{ee'e''}$ | - |
| 6 | 7.03-7.05 | 2H | doublet | Ar- $\text{H}_{gg'}$ | |
| 7 | 7.43-7.45 | 2H | doublet | Ar- $\text{H}_{ff'}$ | |
| 8 | 5.43 | 1H | singlet | Ar- H_h | - |
| 9 | 9.29 | 1H | singlet | Ar- H_i | - |
| 10 | 9.51 | 1H | singlet | Ar- H_j | - |
| 11 | 9.84 | 1H | singlet | Ar- H_k | - |

Figure 2.2.3: Mass spectra of 4-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-6-methyl-2-thioxo-1,2,3,4-tetrahydro pyrimidine-5-carboxamide (SRG-1).



Scheme 2.2.1: Proposed mass fragmentation of 4-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-6-methyl-2-thioxo-1,2,3,4-tetrahydro pyrimidine-5-carboxamide (SRG-1).




LITERATURE CITED

- (1) McConnell, J. D., Bruskewitz, R., Walsh, P., Andriole, G., Lieber, M., Holtgrewe, H. L., Albertsen, P., Roehrborn, C. G., Nickel, J. C., Wang, D. Z., Taylor, A. M., Waldstreicher, J. "The effect of finasteride on the risk of acute urinary retention and the need for surgical treatment among men with benign prostatic hyperplasia. Finasteride Long-Term Efficacy and Safety Study Group." *N. Engl. J. Med.*, **1998**, 338, 557.
- (2) Roehrborn, C. G., Oesterling, J. E., Auerbach, S., Kaplan, S. A., Lloyd, L. K., Milam, D. F., Padley, R. J. *Urology*, **1996**, 47, 159.
- (3) Heible, J. P., Bylund, D. B., Clarke, D. E., Eikenburg, D. C., Langer, S. Z., Lefkowitz, R. J., Minneman, K. P., Ruffolo, R. R. International Union of Pharmacology X. Recommendation for Nomenclature of α_1 -Adrenoceptors: Consensus Update *Pharmacol. Rev.*, **1995**, 47, 267.
- (4) Forray, C., Bard, J. A., Wetzel, J. M., Chiu, G., Shapiro, E., Tang, R., Lepor, H., Hartig, P. R., Weinshank, R. L., Branchek, T. A., Gluchowski, C. The α_1 -adrenergic receptor that mediates smooth muscle contraction in human prostate has the pharmacological properties of the cloned human α_{1c} subtype. *Mol. Pharma.*, **1994**, 45, 703-708.
- (5) Barrow, J. C., Nantermet, P. G., Selnick, H. G., Glass, K. L., Rittle, K. E., Gilbert, K. F., Steele, T. G., Homnick, C. F., Freidinger, R. M., Ransom, R. W., Kling, P., Reiss, D., Broten, T. P., Schorn, T. W., Chang, R. S. L., O'Malley, S. S., Olah, T. V., Ellis, J. D., Barrish, A., Kassahun, K., Leppert, P., Nagarathnam, D., Forray, C. "In Vitro and in Vivo Evaluation of Dihydropyrimidinone C-5 Amides as Potent and Selective α_1 A Receptor Antagonists for the Treatment of Benign Prostatic Hyperplasia." *J. Med. Chem.*, **2000**, 43, 2703-2718.
- (6) Nagarathnam, D., Miao, S. W., Lagu, B., Chiu, G., Fang, J., Dhar, T. G. M., Zhang, J., Tyagarajan, S., Marzabadi, M. R., Zhang, F., Wong, W. C., Sun, W., Tian, D., Wetzel, J. M., Forray, C., Chang, R. S. L., Broten, T. P., Ransom, R. W., Schorn, T. W., Chen, T. B., O'Malley, S., Kling, P., Schneck, K., Bendesky, R., Harrell, C., M., Vyas, K. P., Gluchowski, C. "Design and Synthesis of Novel α_1 A Adrenoceptor-Selective Antagonists. 1. Structure-Activity Relationship in Dihydropyrimidinones." *J. Med. Chem.*, **1999**, 42(23), 4764-4777.
- (7) Atwal, K. S., Swanson, B. N., Unger, S. E., Floyd, D. M., Moreland, S., Hedberg, A., O'Reilly, B. C. "Dihydropyrimidine calcium channel blockers. 3. 3-Carbamoyl-4-aryl-1,2,3,4-tetrahydro -6-methyl-5-pyrimidine

- carboxylic acid esters as orally effective antihypertensive agents." *J. Med. Chem.*, **1991**, 34, 806-811.
- (8) Abdel-Aziz, S. A., Allimony, H. A., El-Shaaer, H. M., Ali, U. F., Abdel-Rahman, R. M. "Fused cyanopyrimidines. Part II. Synthesis and reactions of fused cyanopyrimidine derivatives as affecting enzymic agents." *Phos. Sul. Sili.*, **1996**, 113(1-4), 67-77.
- (9) Hu, E. H., Sidler, D. R., Dolling, U. "Unprecedented catalytic three component one-pot condensation reaction: An efficient synthesis of 5-alkoxycarbonyl-4-aryl-dihydropyrimidin-2(1H)-ones." *J. Org. Chem.*, **1998**, 63(10), 3454-3457.
- (10) Ranu, B. C., Hajra, A., Jana, U. "Indium(III) chloride-catalyzed one-pot synthesis of dihydropyrimidinones by a three-component coupling of 1,3-dicarbonyl compounds, aldehydes, and urea: an improved procedure for the Biginelli reaction." *J. Org. Chem.*, **2000**, 65, 6270-6272.
- (11) Bose, D. S., Liyakat, F., Mereyala, H. B. "Green Chemistry Approaches to the Synthesis of 5-Alkoxycarbonyl-4-aryl-3,4- dihydropyrimidin-2(1H)-ones by a Three-Component Coupling of One-Pot Condensation Reaction: Comparison of Ethanol, Water, and Solvent-free Conditions." *J. Org. Chem.*, **2003**, 68(2), 587-590.
- (12) Yijun, H., Fengyue, Y., Chengjian, Z. *J. Am. Chem. Soc.*, **2005**, 127, 16386.
- (13) Zhou, H., He, M., Liu, C., Luo, G. "One-pot synthesis of 3,4-dihydropyrimidin-2(1H)-ones using hydrochloric acid- β -cyclodextrin-combined catalysts." *Lett. Org. Chem.*, **2006**, 3(3), 225-227.
- (14) Sharma, S. D., Gogoi, P., Konwar, D. "A highly efficient and green method for the synthesis of 3,4-dihydropyrimidin-2-ones and 1,5-benzodiazepines catalyzed by dodecyl sulfonic acid in water." *Green Chem.*, **2007**, 9(2), 153-157.
- (15) Naozumi N., Ryosuke, B., Chika, N., Kazuaki, O., Machida, M. "A novel method for the synthesis of 4-thiopyrimidine nucleosides using hypervalent iodine compounds." *Org. Biomol. Chem.*, **2003**, 1, 3692-3697.
- (16) Stadler, A., Yousefi, B. H., Dallinger, D., Walla, P., Van der Eycken, E., Kaval, N., Kappe, C. O. "Scalability of microwave-assisted organic synthesis. From single-mode to multimode parallel batch reactors." *Org. Process Res. Dev.*, **2003**, 7(5), 707-716.

- (17) Yadav, J. S., Reddy, B. V. S., Reddy, E. J., Ramalingam, T. "Microwave-assisted efficient synthesis of dihydro pyrimidines: improved high yielding protocol for the Biginelli reaction." *J. Chem. Res.*, **2000**, 7, 354-355.
- (18) Gohain, M., Prajapati, D., Sandhu, J. S. "A novel Cu-catalysed three-component one-pot synthesis of dihydro-pyrimidin-2(1H)-ones using microwaves under solvent-free conditions." *Synlett*, **2004**, 2, 235-238.
- (19) Weiwei, L., Yulin, L. *J. Comb. Chem.*, **2005**, 7, 721.
- (20) Zong-Ting, W., Shu-Chao, W., Li-Wen, X. "Three new highly oxygenated diterpenoids from *excoecaria cochinchinensis* Lour." *Helv. Chimica Acta*, **2005**, 88(5), 986-973.
- (21) Dallinger, D., Stadler, A., Kappe, C. O. *Pure Appl. Chem*, **2004**, 76, 1017.
- (21) Holland, J. F., Robert, G., Paul, S., Howard, T. "Inhibition of Tumor Growth in Mice by a Series of 2-Substituted Thiopyrimidines." *Can. Res.*, **1958**, 18, 776-780.
- (22) Di, Marco, A., Gaetani, M. "Inhibiting action of some thiopyrimidines on the growth of transplant tumors." *Tumori*, **1956**, 42, 531-537.
- (23) Sondhi, S. M., Dinodia, M., Rani, R., Shukla, R., Raghubir, R. "Synthesis, anti-inflammatory and analgesic activity evaluation of some pyrimidine derivatives." *Ind. J. Chem.*, 2009, 48B, 273-281.
- (24) Howells, R. E., Tinsly, J., Devancy, E., Smith, G. *Acta. Tropica.*, **1982**, 38, 289.
- (25) Cocco, M. T., Congiu, C., Onnis, V., Piras, R. *Farmaco.*, **2001**, 56, 741.
- (26) Wili, P., Jie, C., Jun, W., Mingzhi, L., Xujian, X., Zhuanli, F., Shuomingshv, S. G., CN. 1, 348,985 (Cl. CO7D 239/46) 2000; *Chem. Abstr.*, **2003**, 139, 53027.
- (27) Petric, C. R., Cotton, H. B., Me-Kerman, P. A., *J. Med. Chem.*, **1988**, 28, 1010.
- (28) Patil, A. D., Kumar, N. V., Kokke, W. C., Bean, M. F., Freyer, A. J.; De Brosse, C., Mai, S., Truneh, A., Faulkner, D. J., Carte, B., Breen, A. L., Hertzberg, R. P., Johnson, R. K., Westley, J. W., Potts, B. C. M. *J. Org. Chem.*, **1995**, 60, 1182-1185.
- (29) Jani, M. K., Shah, B. R., Udavia, N. K. "Synthesis and antimicrobial activity of some pyrimidine-2-thiones." *Chem. Environ. Res.*, **1992**, 1, 345-349.
- (30) Christopherson, R. I., Seymour, K. K., Yeo, A. E. T. "Pyrimidine antagonists and antifolates as antimalarial drugs." *Adv. Exper. Med. Bio.*, **1998**, 431, 705-709.

- (31) Cocco, M. T., Congiu, C., Onnis, V. *Farmaco.*, **1995**, 50, 73.
- (32) Ladani, M. J., Tala, S. D., Akbari, J. D., Dhaduk, M. F., Joshi, H. S. "Synthesis and biological study of oxypyrimidines and thiopyrimidines of 2- (2,4-dichlorophenyl) imidazo[1,2-a] pyridin- 3 -carbaldehyde." *J. Ind. Chem. Soc.*, **2009**, 86(1), 104-108.
- (33) Desai, K. R., Mistry, R. N. "Microwave studies on the synthesis of some new heterocyclic chalcone and pyrimidine-2-thione derivatives and their antibacterial activity." *Asian J. Chem.*, **2004**, 16(1), 201-206.
- (34) Abdel-Rahman, A. A. H., Abdel-Megied, A. E. S., Mohamed, A. M. H., Kasem, E. R., Shabaan, M. T. "Synthesis and antimicrobial evaluation of some chalcones and their derived pyrazoles, pyrazolines, isoxazolines, and 5,6-dihydropyrimidine-2-(1H)-thiones." *Monatshefte fur Chemie.*, **2007**, 138, 889-897.
- (35) Kern, E. R., Mark, N. P., Debra, C. Q., Kathy, A. K., Tiwari, K. N., Joseph, A. M., John, A. S. "Activities of certain 5-substituted 4'-thiopyrimidine nucleosides against orthopoxvirus infections." *Antimicrob. Agent Chemother.*, **2009**, 53, 572-579.
- (36) Gong, L. Z., Nan, Li, Xiao-Hua C., Song, J., Shi-Wei L., Wu F., "Highly Enantioselective Organocatalytic Biginelli and Biginelli-Like Condensations: Reversal of the stereochemistry by tuning the 3,3'-disubstituents of phosphoric acids." *J. Amer. Chem. Soc.*, **2009**, 131, 15301-15310.
- (37) Attia, A. M., Sallam, M. A., Almehti, A. A., Abbasi, M. M. *Nucleosides Nucleotides*, **1999**, 18, 2307.
- (38) Desai, K. R. Mistry, R. N. "Microwave studies on the synthesis of some new heterocyclic chalcone and pyrimidine-2-thione derivatives and their antibacterial activity." *Asian J. Chem.*, **2003**, 15(1), 91-94.
- (39) El-Ghaffar, A., Nahed, F., Kassab, R. R. S., Soliman, Fekria, M. A. "Synthesis and reactions of new substituted pyrimidine thione derivatives as antimicrobial agents." *Al-Azhar Bull. Sci.*, **2000**, 11(1), 161-170.




Section-III

Synthesis of N-methylpyrimidines

INTRODUCTION

Pyrimidine (1) is the trivial name for 1,3-diazine: two meta oriental -CH units in benzene have been replaced by nitrogen atom.



Pyrimidine derivatives have attracted considerable attention because of their pharmacological properties¹⁻², including antiviral^{3, 4}, antitumor⁵, antibacterial^{6, 7}, and antihypertensive^{8, 9}, antileukemic¹⁰ effects. Thus, pyrimidines have been subjected to a large variety of structural modifications in order to synthesize derivatives with different biological properties. In recent years, pyrimidines represent a heterocyclic system with remarkable pharmacological efficiency which shows a very similar pharmacological profile of pyrimidine derivatives calcium channel modulators¹¹⁻¹⁵ e.g. Nifedipine. Today, such medicines are used for the treatment of cardiovascular diseases¹⁶.

The remarkable biological activities of pyrimidine ring are due to structure similarity to purine RNA and DNA structures¹⁷. e.g. uracil, thiamine. Various workers have reported the synthesis of 1-methyl pyrimidine and studied their biological activity.

Thus, due to biological potential of the N-methyl pyrimidines, the present work was undertaken to synthesize some novel N-methyl pyrimidine derivatives.

EXPERIMENTAL

Synthesis of 4-(3,4-dimethoxyphenyl)-6-methyl-2-oxo-N-phenyl-1,2,3,4-tetrahydropyrimidine-5-carboxamide.

[A] Synthesis of 3, 4-dimethoxybenzaldehyde:

As per Section - I [A].

[B] Synthesis of 3-oxo-N-phenylbutanamide:

As per Section – I [B].

[C] Synthesis of 6-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-3,4-dimethyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide (RDV-1):

A mixture of 3, 4 dimethoxy aldehyde (0.01 M), 3-oxo-N-phenylbutanamide (0.01 M) and N-methylurea (0.012 M) in methanol was refluxed for 12 hours in presence of few drops of con. HCl as catalyst. The product was isolated and crystallized from ethanol. All the synthesized compounds were recrystallized from ethanol.

REACTION SCHEME

Scheme [c]:




Table 2.3.1: Physical constants of N-methyl pyrimidenes.

| Sr. No. | Code | R | M.F. | M. Wt. (g/mol) | R _f * Value | M.P. °C | Yield % |
|---------|--------|---------------------|---|-------------------|---------------------------|------------|------------|
| 1 | RDV-1 | 4-OCH ₃ | C ₂₂ H ₂₅ N ₃ O ₅ | 411 | 0.68 | 254 | 52 |
| 2 | RDV-2 | 4-CH ₃ | C ₂₂ H ₂₅ N ₃ O ₄ | 395 | 0.49 | 219 | 61 |
| 3 | RDV-3 | 4-Cl | C ₂₁ H ₂₂ ClN ₃ O ₄ | 415 | 0.72 | 264 | 49 |
| 4 | RDV-4 | 2- CH ₃ | C ₂₂ H ₂₅ N ₃ O ₄ | 395 | 0.66 | 245 | 72 |
| 5 | RDV-5 | 3- OCH ₃ | C ₂₂ H ₂₅ N ₃ O ₅ | 411 | 0.59 | 261 | 69 |
| 6 | RDV-6 | 4-F | C ₂₁ H ₂₂ FN ₃ O ₄ | 399 | 0.51 | 251 | 76 |
| 7 | RDV-7 | 2,5 – di Cl | C ₂₁ H ₂₁ Cl ₂ N ₃ O ₄ | 450 | 0.76 | 244 | 47 |
| 8 | RDV-8 | 3-Cl | C ₂₁ H ₂₂ ClN ₃ O ₄ | 415 | 0.71 | 228 | 66 |
| 9 | RDV-9 | 3,4 – di Cl | C ₂₁ H ₂₁ Cl ₂ N ₃ O ₄ | 450 | 0.54 | 247 | 61 |
| 10 | RDV-10 | 3-Cl, 4-F | C ₂₂ H ₂₅ ClFN ₃ O ₄ | 449 | 0.62 | 256 | 65 |

* Chloroform : Methanol: 9:1

The various physical constants such as R_f value, melting point and percentage of yield for all the synthesized N-methylpyrimidine derivatives are given in Table 2.3.1. The characterization was done by IR, ^1H NMR and mass spectra.

Infrared spectra:

The IR spectra were recorded by SHIMADZU-FTIR-8400 Spectrophotometer in the frequency range of $4000\text{--}400\text{ cm}^{-1}$ by KBr powder method. Figure 2.3.1 shows IR spectra of RDV-1. The IR spectral data for RDV-1 is given in Table 2.3.2. The spectral data for all other compounds are reported in Table 2.3.3.

^1H NMR Spectra:

The NMR spectra were recorded by BRUKER Spectrometer (400 MHz) using internal reference TMS and solvent $\text{CDCl}_3/\text{DMSO}$. Figure 2.3.2 shows NMR spectra of RDV-1. The spectral data for RDV-1 is given in Table 2.3.4.

Mass spectra:

The Mass spectra were recorded by GCMS-SHIMADZU-QP2010. Figure 2.3.3 shows mass spectra of RDV-1. The proposed mass fragmentation of the same compound is also given in Scheme 2.3.1.

Figure 2.3.1: IR spectra of 6-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-3,4-dimethyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide (RDV-1)




Table 2.3.2: IR spectral data of 6-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-3,4-dimethyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide (RDV-1).

| Type | Vibration mode | Frequency in cm^{-1} | |
|----------|----------------------|-------------------------------|-----------|
| | | Observed | Reported |
| Alkane | C-H str. (asym.) | 2912.61 | 2975-2900 |
| | C-H str. (sym.) | 2817.84 | 2880-2810 |
| | C-H def. (asym.) | 1446.66 | 1480-1435 |
| | C-H def. (sym.) | 1313.57 | 1985-1350 |
| Aromatic | C-H str. | 3024.48 | 3100-3000 |
| | C=C str. | 1514.10 | 1585-1480 |
| | C-H i.p. def. | 1143.83 | 1125-1090 |
| | C-H o.o.p. def. | 796.63 | 860-810 |
| Ketones | C=O str.(cyclic) | 1715.92 | 1740-1680 |
| | C=O str. (aliphatic) | 1674.27 | 1710-1650 |
| Nitrogen | C-N str. | 1255.84 | 1350-1200 |
| | N-H str. | 3342.75 | 3400-3200 |
| | N-H def. | 1514.10 | 1650-1500 |
| Ether | C-O-C str. (asym.) | 1143.83 | 1400-1000 |
| | C-O-C str. (sym.) | 1024.24 | 1075-1020 |

Table 2.3.3: IR spectral data of synthesized N-methylpyrimidines.

| Compounds | <i>IR ν, (cm⁻¹)</i> | | | | | |
|---------------|---|----------------|---------------|---------------|----------------|---------|
| | C=O (cyclic) | C=O (alip.) | N-H (Str.) | C=C (Str.) | C-H (asym.) | R |
| RDV-2 | 1716.41 | 1659.20 | 3289.33 | 1524.23 | 2931.63 | 2835.47 |
| RDV-3 | 1710.57 | 1657.90 | 3257.32 | 1541.49 | 2984.20 | 757.49 |
| RDV-4 | 1711.12 | 1683.60 | 3238.29 | 1524.53 | 2944.29 | 2826.24 |
| RDV-5 | 1707.23 | 1668.05 | 3354.62 | 1511.17 | 2928.49 | 1133.24 |
| RDV-6 | 1707.54 | 1679.01 | 3359.54 | 1539.51 | 2956.52 | 688.11 |
| RDV-7 | 1716.68 | 1682.55 | 3275.61 | 1542.20 | 2929.49 | 756.29 |
| RDV-8 | 1706.10 | 1667.47 | 3361.12 | 1529.28 | 2937.40 | 771.57 |
| RDV-9 | 1715.37 | 1666.61 | 3281.19 | 1528.24 | 2961.19 | 749.24 |
| RDV-10 | 1709.58 | 1658.08 | 3266.08 | 1540.47 | 2983.24 | 676.17 |

Figure 2.3.2: ^1H NMR spectra of 6-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-3,4-dimethyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide (RDV-1).







Table 2.3.4: ^1H NMR spectral data of 6-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-3,4-dimethyl-2-oxo-1,2,3,4-tetrahydro pyrimidine-5-carboxamide (RDV-1).




| Singal No. | Signal Position (δ ppm) | Relative No. of Protons | Multiplicity | Inference | J (Hz) |
|------------|---------------------------------|-------------------------|--------------|-------------------------------|----------|
| 1 | 2.24 | 3H | singlet | $-\text{CH}_a$ | - |
| 2 | 2.76 | 3H | singlet | $-\text{CH}_b$ | - |
| 3 | 3.58 | 3H | triplet | $-\text{OCH}_c$ | - |
| 4 | 3.79 | 6H | singlet | $-\text{OCH}_{dd'}$ | - |
| 5 | 5.15 | 1H | singlet | $-\text{H}_h$ | - |
| 6 | 6.46-7.72 | 7H | multiplat | $-\text{Ar-H}_{e-g, kk' ll'}$ | - |
| 7 | 8.10 | 1H | singlet | $-\text{NH}_i$ | - |
| 8 | 8.87 | 1H | singlet | $-\text{NH}_j$ | - |

Figure 2.3.3: Mass spectra of 6-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-3,4-dimethyl-2-oxo-1,2,3,4-tetrahydro pyrimidine-5-carboxamide (RDV-1).




Scheme 2.3.1: Proposed mass fragmentation of 6-(3,4-dimethoxyphenyl)-N-(4-methoxyphenyl)-3,4-dimethyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide (RDV-1).



LITERATURE CITED

- (1) Kleeman, A., Engel, J. "Pharmazeut Wirkstoffe II." *Aufl. Thieme. Stuttgart, New York.* **1982**, 25.
- (2) Wagner, E., Poreba, K., Jakowicz, I., Balicka, D., Rutkowska, M., Kedzierska, G. L., Szelag. "Synthesis and pharmacological properties of derivatives of isoxazolo[4,3-d]pyrimidine. III." *Farmaco*, **1995**, 50(3), 183-187.
- (3) Maquoi, E., Sounni, N. E., Devy, L., Olivier, F., Frankenne, F., Krell, H. V., Gramms, F., Foidart, J. M., Noel, A. *Clin. Can. Res.*, **2004**, 10, 4038-4045.
- (4) Shaoman, Z., Julie, M. B., Katherine, Z. B., John, C. D., Earl, R. K., Elizabeth, G., Yung-Chi, C., Jiri, Z. "Synthesis and Antiviral Activity of (Z)- and (E)-2,2-Bis(hydroxymethyl)cyclopropylidene]methylpurines and -pyrimidines: Second-Generation Methylenecyclopropane Analogues of Nucleosides." *J. Med. Chem.* **2004**, 47, 566-575.
- (5) Krishna, K. K., Barbara, A. M., Csaba, H., Joseph, R. B. "Enhancement of Antitumor Activity of 2,4-Diamino-5-(3',4'-dichlorophenyl)-6-methylpyrimidine and Baker's Antifol (Triazinate) with Carboxypeptidase G₁¹" *Cancer Res.*, **1979**, 39, 3441-3445.
- (6) El-Sayed, H. A., Moustafa, A. H., Haikal, A. Z., Abdou, I. M., El-Ashry, E. S. H. "Synthesis and evaluation of antimicrobial activity of some pyrimidine glycosides." *Nucleosides, Nucleotides and Nucleic Acids*, **2008**, 27(9), 1061-1071.
- (7) Khadiga, M. G., Farag, T., Khairia, M. Y. "Synthesis and evaluation of some 2-, 4- and 2,4-disubstituted-6-methylpyrimidine derivatives for antimicrobial activity." *J. Ind. Chem. Soc.*, **1986**, LXIII, pp. 914-917.
- (8) Botros, S., Youssef, K. M., Issac, Z. "Synthesis of novel 1,4-disubstituted piperazines as potential antihypertensive agents." *Egy. J. Pharm. Sci.*, **1987**, 30, 419-427.
- (9) Badran, M. M., Youssef, K. M. "Novel piperazinyl-substituted-pyrimidines as possible anti-hypertensive and vasodilators." *Egy. J. Pharm. Sci.*, **1989**, 30, 444-462.
- (10) Ghoneim, K. M., El-Telbany, F. A., El-Enany, M., Youssef, K. M. "Synthesis of some novel 2-benzylthio-4-substituted amino-6-methylpyrimidines of expected antileukemic activity." *Ori. J. Chem.*, **1987**, 3(3), 312-318.
- (11) Alfredo, P., Ramon, A., Juan, J. V., Julio, A. B., Miguel, F. de C. J., Carlos, S., Jaime, G. P., Isabel, F., Julia, S. A. "Synthesis and structure of new

- pyrido[2,3-d]pyrimidine derivatives with calcium channel antagonist activity." *Tetrah.*, **1994**, 50(27), 8085-8098.
- (12) Sircar, I., Gregor, E. K., Anderson, K. R., Haleen, S. J., Shih, Y. H., Weishaar, R. E., Steffen, R. P., Pugsley, T. A., Taylor, M. D. "Calcium channel blocking and positive inotropic activities of ethyl 5-cyano-1,4-dihydro-6-methyl-2-[(phenylsulfonyl)methyl]-4-aryl-3-pyridinecarboxylate and analogs. Synthesis and structure-activity relationships." *J. Med. Chem.*, **1991**, 34(7), 2248-2260.
- (13) Guarneri, L., Patrizia, A., Marina, I., Elena, P., Carlo, T., Amedeo, L.; Rodolfo, T. *Arzneimittel Forschung*, **1996**, 46(10), 15.
- (14) Wetzel, J. M., Miao, S. W., Borden, L. A., Branhek, T. A., Gluchowski, C. "Discovery of .alpha.1a-Adrenergic Receptor Antagonists Based on the L-Type Ca²⁺ Channel Antagonist Niguldipine." *J. Med. Chem.*, **1995**, 38, 1579.
- (15) Sweet, F., Fissekis, J. D. "Synthesis of 3,4-dihydro-2(1H)-pyrimidinones and the mechanism of the Biginelli reaction." *J. Ame. Chem. Soc.*, **1973**, 95(26), 8741-8746.
- (16) Lundy, Amber; Lutfi, Nahla; Beckey, Cherylyn. "Review of nifedipine GITS in the treatment of high risk patients with coronary artery disease and hypertension." *Vascular Health Risk Management*, **2009**, 5, 429-440.
- (17) Bannwarth, W., Iaiza, P. "Triple helices of optimally capped duplex DNA (α - DNA) with homopyrimidine DNA and RNA at Neutral pH." *Hel. Chim. Acta*, **1999**, 82(11), 1806-1812.



Section-IV

Synthesis of Tetrahydropyrimidines

INTRODUCTION

The chemistry and the synthesis of tetrahydropyrimidine have been attracting widespread attention in recent years. The present popularity of these compounds is mainly due to their close structural relationship to the clinically important dihydropyridine calcium channel blockers¹⁻⁶.

Literature survey shows that lots of work has been done for tetrahydropyrimidines. Many researchers have been worked on QSAR study of tetrahydropyrimidines⁷⁻⁹. Zhang and co-workers have reported the synthesis of tetrahydropyrimidine via catalyst free multicomponent reaction¹⁰. One pot synthesis of these compounds has been reported by Akbar et al.¹¹ using zeolite as catalyst. Perumal and co-workers¹² have synthesized tetrahydropyrimidines using phosphotungstic acid. Using microwave irradiation, Roy and Bordoloi¹³ have synthesized some new tetrahydropyrimidines. Zhao and Liu¹⁴ have reported the synthesis of novel fluoroalkylated multifunctional tetra hydropyrimidines. The stereomutations of conformational atropisomers of hindered 1,2-diaryltetrahydropyrimidines has been reported by Gracia and co-worker¹⁵. Baldev et al¹⁶ have reported the thermal and microwave assisted synthesis of tetrahydropyrimidines. By ultrasonic irradiation, Muravyova¹⁷ et al have synthesized some new tetrahydropyrimidines.

Tetrahydropyrimidine is known as a versatile heterocyclic compound which has been subjected to a large variety of structural modifications in order to synthesize derivatives with different biological properties. Their various condensed derivatives are reported to possess calcium antagonist,¹⁸⁻²⁰ anti-inflammatory,²¹⁻²³ analgesic,²⁴ antitumor,^{25, 26} antidepressant,²⁷ antibacterial, and antifungal effects.²⁸⁻³⁰ Several synthetic approaches have been reported for the synthesis of fused heterocyclic pyrimidine derivatives.³¹⁻³³ These compounds act as muscarinic agonist in the rat central nervous system.^{34, 35} William and co-workers have reported the M1 receptor of tetrahydropyrimidines and their biochemical activity.³⁶ The antidepressant³⁷ and fungicidal³⁸ activity of some of these derivatives have also been reported. Upshall³⁹ has reported the nicotinic activity whereas Faust and co-workers⁴¹ have reported the antihypertensive activity of these compounds.

Thus, due to the importance of biological active tetrahydropyrimidines, the present chapter describes the synthesis and characterization of some tetrahydropyrimidine derivatives.

EXPERIMENTAL

Synthesis of 4-amino-2-hydroxy-6-phenyl-1,4,5,6-tetrahydropyrimidine-5-carbonitrile:

A mixture of substituted aldehyde (0.01 M), malenonitrile (0.01 M) and urea (0.012 M) in methanol was refluxed for 5 hours. The product was isolated and crystallized from ethanol. All the synthesized compounds were recrystallized from ethanol.

REACTION SCHEME




Table 2.4.1: Physical constants of Tetrahydropyrimidenes.

| Sr. No. | Code | R | M.F. | M. Wt. (g/mol) | R _f * Value | M.P. °C | Yield % |
|---------|--------|---|---|-------------------|---------------------------|------------|------------|
| 1 | SAR-1 | -C ₆ H ₅ | C ₁₁ H ₁₂ N ₄ O | 216 | 0.52 | 64 | 93 |
| 2 | SAR-2 | 4-Cl-C ₆ H ₄ | C ₁₃ H ₁₃ ClN ₄ O | 250 | 0.61 | 110 | 80 |
| 3 | SAR-3 | Cinnamaldehyde | C ₁₃ H ₁₄ N ₄ O | 242 | 0.66 | 146 | 94 |
| 4 | SAR-4 | 3-Cl-C ₆ H ₄ | C ₁₃ H ₁₃ ClN ₄ O | 250 | 0.49 | 150 | 70 |
| 5 | SAR-5 | 4-F-C ₆ H ₄ | C ₁₁ H ₁₁ FN ₄ O | 234 | 0.58 | 134 | 59 |
| 6 | SAR-6 | 4-OCH ₃ -C ₆ H ₄ | C ₁₂ H ₁₄ N ₄ O ₂ | 246 | 0.49 | 154 | 60 |
| 7 | SAR-7 | 3-NO ₂ -C ₆ H ₄ | C ₁₁ H ₁₁ N ₅ O ₃ | 261 | 0.71 | 184 | 77 |
| 8 | SAR-8 | Vaniline | C ₁₂ H ₁₄ N ₄ O ₃ | 262 | 0.57 | 115 | 50 |
| 9 | SAR-9 | Furfuraldehyde | C ₉ H ₁₀ N ₄ O ₂ | 206 | 0.50 | 80 | 76 |
| 10 | SAR-10 | 4-OH- C ₆ H ₄ | C ₁₁ H ₁₂ N ₄ O ₂ | 232 | 0.48 | 70 | 85 |

* Chloroform:Methanol:- 9:1

The various physical constants such as R_f value, melting point and percentage of yield for all the synthesized tetrahydropyrimidine derivatives are given in Table 2.4.1. The characterization was done by IR, ^1H NMR and mass spectra.

Infrared spectra:

The IR spectra were recorded by SHIMADZU-FTIR-8400 Spectrophotometer in the frequency range of $4000\text{--}400\text{ cm}^{-1}$ by KBr powder method. Figure 2.4.1 shows IR spectra of SAR-2. The IR spectral data for SAR-2 is given in Table 2.4.2. The spectral data for all other compounds are reported in Table 2.4.3.

^1H NMR Spectra:

The NMR spectra were recorded by BRUKER Spectrometer (400 MHz) using internal reference TMS and solvent $\text{CDCl}_3/\text{DMSO}$. Figure 2.4.2 shows NMR spectra of SAR-2. The spectral data for SAR-2 is given in Table 2.4.4.

Mass spectra:

The Mass spectra were recorded by GCMS-SHIMADZU-QP2010. Figure 2.4.3 shows mass spectra of SAR-2. The proposed mass fragmentation of the same compound is also given in Scheme 2.4.1.

Figure 2.4.1: IR spectra of 4-amino-6-(4-chlorophenyl)-2-hydroxy-1,4,5,6-tetrahydropyrimidine-5-carbonitrile (SAR-2).




Table 2.4.2: IR spectral data of 4-amino-6-(4-chlorophenyl)-2-hydroxy-1,4,5,6-tetrahydropyrimidine-5-carbonitrile (SAR-2).

| Type | Vibration mode | Frequency in cm^{-1} | |
|----------|--------------------|-------------------------------|-----------|
| | | Observed | Reported |
| Alkane | C-H str. (asym.) | 2935.76 | 2975-2900 |
| | C-H str. (sym.) | 2867 | 2880-2810 |
| | C-H def. (asym.) | 1440.87 | 1480-1435 |
| | C-H def. (sym.) | 1351 | 1985-1350 |
| Aromatic | C-H str. | 3007.12 | 3100-3000 |
| | C=C | 1516.10 | 1585-1480 |
| | C-H i.p. def. | 1141.90 | 1125-1090 |
| | C-H o.o.p. def. | 862.21 | 860-810 |
| Nitrile | C \equiv N str. | 2251.84 | 2210-2280 |
| Alcohol | O-H str. | 3594.12 | 3600-3450 |
| Nitrogen | C-N str. | 1317.43 | 1350-1200 |
| | N-H str. | 3294.53 | 3400-3200 |
| | N-H def. | 1599.04 | 1650-1500 |
| Ether | C-O-C str. (asym.) | 1141.90 | 1400-1000 |
| | C-O-C str. (sym.) | 1018.45 | 1075-1020 |

Table 1.3: IR spectral data of synthesized Tetrahydropyrimidenes.

| Compounds | <i>IR ν, (cm⁻¹)</i> | | | | | |
|---------------|---|---------------|------------------------|---------------|----------------|---------|
| | O-H (Str.) | N-H (Str.) | C \equiv N (str.) | C=N (Str.) | C-H (asym.) | R |
| SAR-1 | 3524.12 | 3389.12 | 2235.12 | 1684.21 | 2933.45 | - |
| SAR-3 | 3499.41 | 3412.27 | 2214.32 | 1654.87 | 2912.37 | 786 |
| SAR-4 | 3529.22 | 3437.54 | 2241.51 | 1627.31 | 2962.21 | 1543.34 |
| SAR-5 | 3535.47 | 3388.12 | 2220.54 | 1629.14 | 2945.39 | 746.34 |
| SAR-6 | 3564.10 | 3356.21 | 2268.34 | 1647.61 | 2941.24 | 651.27 |
| SAR-7 | 3584.32 | 3377.54 | 2245.76 | 1630.24 | 2954.67 | 1139.33 |
| SAR-8 | 3568.22 | 3364.30 | 2278.94 | 1655.84 | 2920.20 | 681.28 |
| SAR-9 | 3486.24 | 3394.61 | 2218.58 | 1634.67 | 2961.84 | - |
| SAR-10 | 3530.65 | 3408.27 | 2282.54 | 1672.29 | 2933.48 | 3527.99 |

Figure 2.4.2: ^1H NMR spectra of 4-amino-6-(4-chlorophenyl)-2-hydroxy-1,4,5,6-tetrahydropyrimidine-5-carbonitrile (SAR-2).







Table 2.4.4: ^1H NMR spectral data of 4-amino-6-(4-chlorophenyl)-2-hydroxy-1,4,5,6-tetrahydropyrimidine-5-carbonitrile (SAR-2).




| Singal No. | Signal Position (δ ppm) | Relative No. of Protons | Multiplicity | Inference | <i>J</i> (Hz) |
|------------|---------------------------------|-------------------------|--------------|-----------------------|---------------|
| 1 | 8.15 | 1H | singlet | -NH _a | - |
| 2 | 7.87-7.94 | 2H | singlet | -NH _b | - |
| 3 | 8.36 | 1H | singlet | -OH _c | - |
| 4 | 6.85-6.87 | 2H | doublet | -Ar-H _{dd'} | 6.80 |
| 5 | 7.04-7.07 | 2H | doublet | -Ar-H _{ee''} | 5.16 |
| 7 | 5.40 | 1H | singlet | -Ar-H _f | - |
| 8 | 7.45-7.64 | 2H | singlet | -Ar-H _{g,h} | - |

Figure 2.4.3: Mass spectra of 4-amino-6-(4-chlorophenyl)-2-hydroxy-1,4,5,6-tetrahydropyrimidine-5-carbonitrile (SAR-2).



Scheme 2.4.1: Proposed mass fragmentation of 4-amino-6-(4-chlorophenyl)-2-hydroxy-1,4,5,6-tetrahydropyrimidine-5-carbonitrile (SAR-2).




LITERATURE CITED

- (1) Vitolina, R., Kimenis, A. "Foridon, a new vasodilator and calcium antagonist." *Khimiko-Farmatsevticheskii Zhurnal*, **1989**, 23(7), 889-892.
- (2) Kastron, V. V., Vitolina, R., Khanina, E. L., Duburs, G., Kimenis, A., Kondratenko, N. V., Popov, V. I., Yagupolskii, L. M., Kolomeitsev, A. A. *U.S. Patent*, 4738965, **1988**.
- (3) Cho, H., Ueda, M., Shima, K., Mizuno, A., Hayashimatsu, M., Ohnaka, Y., Takeuchi, Y., Hamaguchi, M., Aisaka, K., Hidaka, T., Kawai, M., Takeda, M., Ishihara, T., Funahashi, K., Satoh, F., Morita, M., Noguchi, T. "Dihydropyrimidines: novel calcium antagonists with potent and long-lasting vasodilative and anti-hypertensive activity." *J. Med. Chem.*, **1989**, 32, 2399.
- (4) Rovnyak, G. C., Atwal, K. S., Hedberg, A., Kimball, S. D., Moreland, S., Gougoutas, J. Z., O'Reilly, B. C., Schwartz, J. M., Mary, F. "Dihydropyrimidine calcium channel blockers. 4. Basic 3-substituted-4-aryl-1,4-dihydropyrimidine-5-carboxylic acid esters. Potent antihypertensive agents." *J. Med. Chem.*, **1992**, 35(17), 3254-3263.
- (5) Ertan, M., Balkan, A., Sarac, S., Uma, S., Renaud, J. F., Rolland, Y. "Synthesis and calcium antagonistic activity of some new 2-thioxo-1,2,3,4-tetrahydropyrimidine derivatives." *Archiv der Pharmazie*, **1991**, 324(3), 135-139.
- (6) Ertan, M., Balkan, A., Sarac, S., Uma, S., Ruebseman, K., Renaud, J. F. "Synthesis and biological evaluations of some 2-thioxo-1,2,3,4-tetrahydropyrimidine derivatives." *Arzneimittel-Forschung*, **1991**, 41(7), 725-727.
- (7) Sawant, R. L., Bhatia, M. S. "Synthesis, screening and QSAR studies of 3-formyl-2-oxo-1,2,3,4-tetrahydropyrimidine analogues as antibacterial agents." *Int. J. Chem. Sci.*, **2008**, 6(1), 45-48.
- (8) Sawant, R. L., Bhatia, M. S. "Synthesis, screening and QSAR studies of 3-formyl-2-thioxo-1,2,3,4-tetrahydropyrimidine analogues as antibacterial agents." *Ori. J. Chem.*, **2008**, 24(1), 181-188.
- (9) Zanatta, N., Madruga, C. C., Marisco, P. C., da R., Luciana, S., Fernandes, L. da S., Flores, D. C., Flores, A. F. C., Burrow, R. A., Bonacorso, H. G., Martins, M. A. P. "Synthesis and structural study of a new series of 2-methylsulfanyl-tetrahydropyrimidines from α -alkoxyvinyl trihalomethyl ketones." *J. Heterocyc. Chem.*, **2008**, 45(1), 221-227.

- (10) Zhang, M., Jiang, H., Liu, H., Zhu, Q. "Convenient One-Pot Synthesis of Multisubstituted Tetrahydropyrimidines via Catalyst-Free Multicomponent Reactions." *Org. Lett.*, **2007**, 9(21), 4111-4113.
- (11) Akbar, M., Foroughifar, N., Karimi, G., Forougifar, N. "One-pot synthesis of tetrahydropyrimidines catalyzed by zeolite." *Synthesis Reactivity in Inorg., Metal-Org., Nano-Metal Chem.*, **2007**, 37(4), 279-282.
- (12) Sivaprasad, Ganesabaskaran; Perumal, Paramasivan T. "Synthesis of novel 4-pyrazolyl-2-oxo-1,2,3,4-tetrahydropyrimidines promoted by phosphotungstic acid." *J. Heterocycl. Chem.*, **2005**, 42(5), 863-866.
- (13) Roy, D. K., Bordoloi, M. "Synthesis of some substituted 2-oxo-1,2,3,4-tetrahydropyrimidines (3,4-dihydropyrimidin-2(1H)-ones) and 2-thioxo-1,2,3,4-tetrahydropyrimidines, catalyzed by tin(II) chloride dihydrate and tin(II) iodide under microwave irradiation." *Ind. J. Chem.*, **2006**, 45B(4), 1067-1071.
- (14) Zhao, Fu-Lu; Liu, Jin-Tao. "A novel and efficient synthesis of fluoroalkylated multifunctional 1,2,3,4-tetrahydropyrimidines." *J. Fluorine Chem.*, **2004**, 125(12), 1841-1845.
- (15) Garcia, M. B., Grilli, S., Lunazzi, L., Mazzanti, A., Orelli, L. R. "Stereomutations of conformational atropisomers of hindered 1,2-diaryltetrahydropyrimidines." *Eur. J. Org. Chem.*, **2002**, (23), 4018-4023.
- (16) Kumar, B., Kaur, B., Kaur, J., Parmar, A., Anand, R. D., Kumar, H. "Thermal/microwave assisted synthesis of substituted tetrahydropyrimidines as potent calcium channel blockers." *Ind. J. Chem.*, **2002**, 41B(7), 1526-1530.
- (17) Muravyova, E. A., Desenko, S. M., Musatov, V. I., Knyazeva, I. V., Shishkina, S. V. Shishkin, O. V., Chebanov, V. A. "Ultrasonic-Promoted Three-Component Synthesis of Some Biologically Active 1,2,5,6-Tetrahydropyrimidines." *J. Combinatorial Chem.*, **2007**, 9(5), 797-803.
- (18) Balkan, A., Ertan, M., Burgemeister, T. "Synthesis and structural evaluations of thiazolo[3,2-a]pyrimidine derivatives." *Archiv der Pharmazie*, **1992**, 325(8), 499-501.
- (19) Balkan, A., Uma, S., Ertan, M., Wiegrebe, W. "Thiazolo[3,2-a]pyrimidine derivatives as calcium antagonists." *Pharmazie*, **1992**, 47(9), 687-688.
- (20) Balkan, A., Tozkoparan, B., Ertan, M., Sara, Y., Ertekin, N. "New thiazolo[3,2-a] pyrimidine derivatives, synthesis and calcium antagonistic activities." *Bollettino chimico farmaceutico*, **1996**, 135(11), 648-652.
- (21) Doria, G., Passarotti, C., Sala, R., Magrini, R., Sberze, P., Tibolla, M., Ceserani, R., Arcari, G., Castello, R., Toti, D. "7-trans-(2-Pyridylethenyl)-


- 5H-thiazolo[3,2-a]pyrimidine-5-ones: synthesis and pharmacological activity. " *Il Farmaco; edizione scientifica*, **1985**, 40(12), 885-894.
- (22) Jeanneau-Nicolle, E., Benoit-Guyod, M., Namil, A., Leclerc, G. "New thiazolo[3,2-a]pyrimidine derivatives, synthesis and structure-activity relationships. " *Eur. J. Med. Chem.*, **1992**, 27(2), 115-120.
- (23) Tozkoparan, B., Ertan, M., Kelicen, P., Demirdamar, R. "Synthesis and anti-inflammatory activities of some thiazolo[3,2-a]pyrimidine derivatives." *Farmaco*, **1999**, 54(9), 588-593.
- (24) Baetz, J. "Analgesic 2,3-dihydrothiazolo[3,2-a]pyrimidin-5-ones. " Brit. Patent, **1973**, 3 pp. 1,306,558.
- (25) Dash, B., Patra, M., Mohapatra, P. K. "Thiazolopyrimidine derivatives." *J. Insti. Chem.*, **1980**, 52(2), 92-96.
- (26) Teijin Ltd., Japanese Patent, *Jpn. Kokai Tokkyo Koho JP*, 24590 83 24590, **1983**, 58.
- (27) Bigg, D. "Thiazole derivatives and their therapeutic use." Fr. Demande **1981**, 12.
- (28) Mohan, J., Chadha, V. K., Chaudhary, H. S., Sharma, B. D., Pujari, H. K., Mohapatra, L. N. "Heterocyclic systems containing bridgehead nitrogen atom. XIII. Antifungal and antibacterial activities of thiazole and thiazolidinone derivatives." *Ind. J. Experim. Bio.*, **1972**, 10(1), 37-40.
- (29) Dehuri, S. N., Nayak, A. "Studies on thiazolidinones. Part XII: Synthesis and antimicrobial activities of thiazolidinones and their derivatives from cyclic thioureas." *J. Ind. Chem. Soc.*, **1983**, 60(10), 970-974.
- (30) Champaneri, H. R., Modi, S. R., Naik, H. B. "Studies on pyrimidines. Part II. Synthesis and antibacterial activity of 7-(2-hydroxy-5-methylphenyl)-2-acetyl-5-phenyl-3-oxo-2,3,4,5-tetrahydrothiazolo[3,2-a]pyrimidines." *Asian J. Chem.*, **1994**, 6(3), 737-738.
- (31) Kappe, C. O., Roschger, P. "Synthesis and reactions of Biginelli-compounds. Part I." *J. Heterocyc. Chem.*, **1989**, 26(1), 55-64.
- (32) Chanda, K., Dutta, M. C., Vishwakarma, J. N. "A facile one-pot synthetic route to substituted fused tetrahydropyrimidines. Part 4. Synthesis of 1-(aralkyl/aryl)-3-(alkyl/aralkyl/aryl)-5-oxo-1,2,3,4,5,6,7,8-octahydroquinazolines and 1-(aralkyl/aryl)-3-(alkyl/aralkyl/aryl)-7,7-dimethyl-5-oxo-1,2,3,4,5,6,7,8-octahydroquinazolines." *Ind. J. Chem.*, **2006**, 45B(4), 1076-1079.
- (33) Mobinikhaledi, A., Foroughifar, N., Goodarzi, F. "Synthesis of some bicyclic thiazolo- and thiazepinopyrimidine derivatives." *Phosp., Sulfur Silicon Related Elements*, **2004**, 179(3), 507-512.

- (34) Messer, W. S., Jr.; Ojo, B. "Preparation of 1,4,5,6-tetrahydropyrimidines as muscarinic agonists." *PCT Int. Appl.*, CAN 129:330737, **1998**, 18.
- (35) Dunbar, P. G., Durant, G. J., Rho, T., Ojo, B., Huzl, J. J., Smith, D. A., El-Assadi, A. A., Sbeih, S., Ngur, D. O., Periyasamy, S., Hoss, W., Messer, W. S. "Design, Synthesis, and Neurochemical Evaluation of 2-Amino-5-(alkoxycarbonyl)-3,4,5,6-tetrahydropyridines and 2-Amino-5-(alkoxycarbonyl)-1,4,5,6-tetrahydropyrimidines as M1 Muscarinic Receptor Agonists." *J. Med. Chem.*, **1994**, 37(17), 2774-2782.
- (36) Messer, W. S., Jr., Dunbar, P. G., Rho, T., Periyasamy, S., Ngur, D., Ellerbrock, B. R., Bohnett, M., Ryan, K., Durant, G. J., Hoss, W. "Synthesis, biochemical activity and behavioral effects of a series of 1,4,5,6-tetrahydropyrimidines as novel ligands for M1 receptors." *Bioorg. Med. Chem. Lett.*, **1992**, 2(8), 781-6
- (37) Weinhardt, K., Wallach, M. B., Marx, M. "Synthesis and antidepressant profiles of phenyl-substituted 2-amino- and 2-[(alkoxycarbonyl)amino]-1,4,5,6-tetrahydropyrimidines." *J. Med. Chem.*, **1985**, 28(6), 694-8.
- (38) Dolman, H., Kuipers, J., Raven, C. W. "Fungicidal and plant-growth-regulating properties of 2-arylamino-1,4,5,6-tetrahydropyrimidines." *Naturwissenschaften*, **1981**, 68(3), 144.
- (39) Upshall, D. G. "Correlation of chick embryo teratogenicity with the nicotinic activity of a series of tetrahydropyrimidines." *Teratology*, **1972**, 5(3), 287-294.
- (40) Nickell, Louis G.; Gordon, Philip N.; Goenaga, Alvaro. "2-n-Alkylmercapto-1,4,5,6-tetrahydropyrimidines, chemotherapeutic agents for plant rusts." *Plant Disease Reporter*, **1961**, 45, 756-758.
- (41) Faust, J. A.; Yee, L. S.; Sahyun, M. Sahyun Labs., Santa Barbara, CA. "Antihypertensive agents: derivatives of 2-imidazoline and 1,4,5,6-tetrahydropyrimidine." *J. Org. Chem.*, **1961**, 26, 4044-4047.



Part-2

Synthesis of Benzothiazole Derivatives



Synthesis of Azomethines

INTRODUCTION

Benzothiazole derivatives are of particular interest within the realm of medicinal chemistry¹. Pyrazole ring system is of some practical importance, because many drugs and medicines contain a pyrazole ring system. As early as 1884 Knorr discovered the antipyretic (temperature reducing) action of a pyrazole derivative in human beings and due to its antipyretic property, he named the compound "Antipyrine".

Biologists' attention was drawn to this class of series when the pharmacological profile of Riluzole was discovered. Thereafter, benzothiazole derivatives have been studied extensively and found to have diverse chemical reactivity and broad spectrum of biological activity. These compounds possess selective analgesic², antiinflammatory³, antitumor⁴, antitubercular⁵, anticonvulsant⁶, diuretic⁷ and antimicrobial⁸ etc., properties.

It is well known that the introduction of fluorine atom into an organic molecule causes dramatic changes in its biological profile, mainly due to high electronegativity of fluorine, the strong carbon-fluorine bond and increased solubility in lipids⁹. In search for new biodynamic potent molecule, it was thought worthwhile to incorporate some additional heterocyclic moieties in the β -lactam nucleus and study their biological and pharmacological activity¹⁰. The literature survey prompted us to synthesize some new substituted fluoro benzothiazole, azomethines as target compounds.

The chemistry of the carbon-nitrogen double bond plays a vital role in the progress of chemistry science¹¹. azomethines compounds have been used as fine chemicals and medical substrates. azomethines are important class of ligands due to their synthetic flexibility, their selectivity and sensitivity towards the central metal atom, structural similarities with natural biological substances and also due to presence of imine group (N=CH-) which imports in elucidating the mechanism of transformation and rasemination reaction in biological system¹²⁻¹⁴. They are well known intermediates for the preparation of azetidinone, thiazolidinone, formazone, arylacetamide and many other derivatives. The transformation of azomethines in the secondary amines via reduction¹⁵ or Grignard reagents addition¹⁶ or the cycloaddition of these azomethines¹⁷, is well known but a series of less investigated reactions, such

as conversion to amides after the treatment with tri alkyl boranes and CO in the presence of cobalt carbonyl¹⁸, Darzens-type condensations¹⁹ or Wittig reaction with phosphorus ylides²⁰ have also been reported.

Azomethines has a wide range of applications such as corrosion inhibitor²¹, intermediates in various reactions^{22, 23}, in perfumery²⁴ etc. Some are known to be used in many potential drugs²⁵ and are known to possess broad spectrum of biological activities such as antiviral²⁶, anticancer²⁷, anti HIV^{28, 29}, antifungal³⁰, antiparasitic³¹, anti-inflammatory³², anticonvulsant³³, antitumor^{34, 35}, antioxidant³⁶, antiulcerogenic³⁷ etc. The antitubercular³⁸ or cytostatic³⁹ activity of azomethines has been known for long and a large number of azomethines have been tested as non terpenoid insectjuvenile hormone mimics⁴⁰.

Synthesis of azomethines is often carried out with acid-catalyzed and generally by refluxing the mixture of aldehyde (or ketone) and amine in organic medium⁴¹. However, with the assistance of microwave irradiation, it was found that the condensation reaction of salicylaldehyde and various aryl amines could proceed fast and efficiently without solvent. Sun et al⁴² have reported the rapid synthesis of azomethines without solvent under microwave irradiation. Desai and Mistry⁴³ have reported the microwave synthesis and biological evaluation of some azomethines. Various workers reported the synthesis of azomethines by ultrasound irradiation⁴⁴⁻⁴⁷.

Literature survey shows that a lot of work has been done related to azomethines and their metal complexes. Corbin and work⁴⁸ have reported the azomethines-zinc complex from benzthiazolines. Hamil and co-workers⁴⁹ have reported the cobalt and copper complexes of azomethines and their antibacterial studies. Rajavel and Krishnan⁵⁰ have reported the synthesis and characterization of oxovanadium(IV) complexes of the azomethines derived by the condensation of 2-aminobenzaldehyde with various diamines as 1,2-diaminoethane, 1,2-diaminopropane, 1,3-diaminopropane.

In recent years, a lot of work has been done by various workers to synthesize azomethines and further thiazolidinones, azetidinones etc from benzothiazole nucleus. Pai and Chavan⁵¹ have reported the synthesis and antibacterial activity of azetidinones via azomethines from benzothiazole moiety. Vijaykumar and co-workers⁵² have synthesized the azetidinones via

azomethines from thiazole nucleus. Synthesis and antimicrobial activity of crown ethers of azomethines type have also reported⁵³. Sie-tiong et al.⁵⁴ have reported the mesogenic properties of azomethines comprising benzothiazole moiety. Zahid and Claudiu⁵⁵ have also reported the antibacterial activity of copper and nickel complexes of benzothiazole derived azomethines.

Thus, with an effort to capitalize the biological potential of the heterocyclic system and to provide more interesting compounds for biological study, the present work undertaken to synthesize some azomethines bearing benzothiazole nucleus.

EXPERIMENTAL

Synthesis of 6-fluoro-N-[(Z)-phenylmethylidene]-1, 3-benzothiazol-2-amine

[A] Synthesis of 6-fluoro-1,3-benzothiazol-2-amine:

To a solution of p-flouro aniline (1 mole) in chlorobenzene, concentrated sulfuric acid (0.55 mole) was added drop wise over a period of 5 minutes. To this finely divided suspension of p-flouro aniline sulfate, sodium thiocyanate (1.1 mole) was added and the mixture was heated for 3 hours at 100° (inside temperature) in an oil bath. The solution, which now contains the thiourea, was cooled to 30°C and 1.34 moles of sulfuryl chloride was added over a period of 15 minutes, with care that the temperature does not exceed 50°C. The mixture was kept at 50°C for 2 hours (no further evolution of hydrogen chloride) and then the chlorobenzene is removed by filtration. The solid residue was then dissolved in hot water. It was further washed by a current of steam to remove remaining solvent. The resulting solid was re precipitated from alkaline solution. For this, the solid residue was dissolved in concentrated ammonium hydroxide (sp. gr. 0.90). To this alkaline solution, HCl (0.1 N) was added drop wise and stirred. The 6-fluoro-1,3-benzothiazol-2-amine is precipitated, filtered and washed with water. The crude product was isolated and crystallized from absolute ethanol.


[B] Synthesis of 6-fluoro-N-[(E)-(4-methylphenyl)methylidene]-2,3-dihydro-1,3-benzothiazol-2-amine:

An ethanolic solution of 6-flouro 2-amino benzothiazole (0.01 M), 4-methyl benzaldehyde (0.01 M) and glacial acetic acid (as catalyst) was refluxed for 10 hrs. The reaction was monitoring with TLC. The product was isolated and crystallized from absolute ethanol.


Similarly, other azomethines were synthesized.

REACTION SCHEME

[A] Synthesis of 6-methoxy 2-amino benzothiazole:



[B] 6-fluoro-N-[(E)-(4-methylphenyl)methylidene]-2,3-dihydro-1,3-benzothiazol-2-amine:



The various physical constants such as R_f value, melting point and percentage of yield for all the synthesized azomethines are given in Table 2.1.

Table 2.1: Physical constants of synthesized Azomethines.

| Sr. No. | Code | R | M.F. | M. Wt. (g/mol) | R _f * Value | M.P. °C | Yield % |
|---------|-------|------------------------------------|--|-------------------|---------------------------|------------|------------|
| 1 | BT-1 | 4-N(CH ₃) ₂ | C ₁₆ H ₁₄ FN ₃ S | 299 | 0.59 | 163 | 81 |
| 2 | BT-2 | 4-CH ₃ | C ₁₅ H ₁₁ FN ₂ S | 270 | 0.62 | 154 | 76 |
| 3 | BT-3 | 4-Cl | C ₁₄ H ₈ ClFN ₂ S | 290 | 0.48 | 202 | 58 |
| 4 | BT-4 | 3,4 di-OCH ₃ | C ₁₆ H ₁₃ FN ₂ O ₂ S | 316 | 0.66 | 176 | 73 |
| 5 | BT-5 | 4-OCH ₃ | C ₁₅ H ₁₁ FN ₂ OS | 286 | 0.46 | 189 | 64 |
| 6 | BT-6 | 4-F | C ₁₄ H ₈ F ₂ N ₂ S | 274 | 0.75 | 164 | 66 |
| 7 | BT-7 | 2-OCH ₃ | C ₁₅ H ₁₁ FN ₂ OS | 286 | 0.62 | 171 | 49 |
| 8 | BT-8 | 3-Cl | C ₁₄ H ₈ ClFN ₂ S | 290 | 0.67 | 213 | 47 |
| 9 | BT-9 | 4-NO ₂ | C ₁₄ H ₈ FN ₃ O ₂ S | 301 | 0.81 | 210 | 41 |
| 10 | BT-10 | 2,5-di Cl | C ₁₄ H ₇ Cl ₂ FN ₂ S | 325 | 0.73 | 190 | 67 |

* Hexane:Ethyl acetate - 7:3

The characterization was done by IR, ^1H NMR and mass spectra.

Infrared spectra:

The IR spectra were recorded by SHIMADZU-FTIR-8400 Spectrophotometer in the frequency range of $4000\text{--}400\text{ cm}^{-1}$ by KBr powder method. Figure 2.1 shows IR spectra of BT-2 whereas its IR spectral data is given in Table 2.2. The spectral data for all other compounds are reported in Table 2.3.

^1H NMR Spectra:

The NMR spectra were recorded by BRUKER Spectrometer (400 MHz) using internal reference TMS and solvent $\text{CDCl}_3/\text{DMSO}$. Figure 2.2 and Table 1.4 show NMR spectra and spectral data of BT-2.

Mass spectra:

The Mass spectra were recorded by GCMS-SHIMADZU-QP2010. Figure 2.3 shows mass spectra of BT-2. The proposed mass fragmentation of the same compound is also given in Scheme 2.1.

Figure 2.1: IR spectra of 6-fluoro-N-[(Z)-(4-methylphenyl)methylidene] - 1,3-benzothiazol-2-amine (BT-2).




Table 2.2: IR spectral data of 6-fluoro-N-[(Z)-(4-methylphenyl)methylidene]-1,3-benzothiazol-2-amine (BT-2).

| Type | Vibration mode | Frequency in cm^{-1} | |
|------------|--------------------|-------------------------------|-----------|
| | | Observed | Reported |
| Alkane | C-H str. (asym.) | 2914.54 | 2975-2900 |
| | C-H str. (sym.) | 2843 | 2880-2810 |
| | C-H def. (asym.) | 1462.09 | 1480-1435 |
| | C-H def. (sym.) | 1367 | 1985-1350 |
| Aromatic | C-H str. | 3039.91 | 3000-3100 |
| | C=C str. | 1539.25 | 1585-1480 |
| | C-H i.p. def. | 1147.38 | 1125-1090 |
| | C-H o.o.p. def. | 848.71 | 860-810 |
| Azomethine | N=C str. | 1641.48 | 1620-1580 |
| | C-N str. | 1257.63 | 1350-1200 |
| Ether | C-O-C str. (asym.) | 1195.91 | 1275-1200 |
| | C-O-C str. (sym.) | 1047.38 | 1075-1020 |
| Halogen | C-F | 1114.89 | 1400-1000 |

Table 2.3: IR spectral data of synthesized Azomethines.

| Compounds | <i>IR ν, (cm^{-1})</i> | | | | | |
|--------------|--|---------------|----------------|---------------|------------------------|---------|
| | C=N (Str.) | C-N (Str.) | C-F (Bend.) | C=C (Str.) | C-H (str. asym.) | R |
| BT-1 | 1608.69 | 1242.20 | 1145.75 | 1539.25 | 2912.61 | 1242.27 |
| BT-3 | 1621.27 | 1247.99 | 1128.49 | 1537.34 | 2933.83 | 756.21 |
| BT-4 | 1599.97 | 1238.34 | 1145.37 | 1508.37 | 2920.32 | 1165.32 |
| BT-5 | 1618.33 | 1257.63 | 1116.82 | 1533.46 | 2922.25 | 1187.12 |
| BT-6 | 1608.29 | 1246.06 | 1153.47 | 1545.03 | 2941.54 | 1129.17 |
| BT-7 | 1599.04 | 1259.56 | 1112.96 | 1500.67 | 2967.23 | 1177.37 |
| BT-8 | 1610.31 | 1242.23 | 1109.34 | 1519.96 | 2971.35 | 716.24 |
| BT-9 | 1602.45 | 1236.54 | 1137.45 | 1529.37 | 2915.64 | 624.31 |
| BT-10 | 1600.84 | 1257.68 | 1141.90 | 1536.34 | 2933.12 | 784.28 |

Figure 2.2: ^1H NMR spectra of 6-fluoro-N-[(Z)-(4-methylphenyl)methylidene]-1,3-benzothiazol-2-amine (BT-2).







Table 2.4: ^1H NMR spectral data of 6-fluoro-N-[(Z)-(4-methylphenyl)methylidene]-1,3-benzothiazol-2-amine (BT-2).




| Singal No. | Signal Position (δ ppm) | Relative No. of Protons | Multiplicity | Inference | J (Hz) |
|------------|---------------------------------|-------------------------|--------------|----------------------|----------|
| 1 | 2.57 | 3H | Singlet | $-\text{CH}_{3a}$ | - |
| 2 | 6.91-6.96 | 1H | Doublet | $-\text{CH}_b$ | - |
| 3 | 7.28-7.31 | 1H | Singlet | $-\text{CH}_c$ | - |
| 4 | 7.03 | 1H | Singlet | $-\text{CH}_d$ | - |
| 5 | 7.51-7.55 | 2H | Doublet | Ar- $\text{H}_{ee'}$ | 8.96 |
| 6 | 7.96-7.99 | 2H | Doublet | Ar- $\text{H}_{ff'}$ | 7.84 |
| 7 | 8.52 | 1H | Singlet | Ar- H_g | - |

Figure 2.3: Mass spectra of 6-fluoro-N-[(Z)-(4-methylphenyl)methylidene]-1,3-benzothiazol-2-amine (BT-2).



Scheme 2.1: Proposed mass fragmentation of 6-fluoro-N-[(Z)-(4-methylphenyl)methylidene]-1,3-benzothiazol-2-amine (BT-2).



LITERATURE CITED

- (1) Augustin, M., Richter, M., Salas, S. "Reactions with N-acylimino dithiocarbonic acid diesters." *J. fuer Praktische Chem.*, **1980**, 322(1), 55-68.
- (2) Pattan, S. R., Babu, S. N., Angadi, J. "Synthesis and biological activity of 5-[4-(7-substituted-6-fluorobenzothiazol-2-ylaminosulfonyl)benzylidene]-2,4-thiazolidinedione." *Ind. J. Heter. Chem.*, **2002**, 11(4), 333-334.
- (3) Palagiano, F., Arenare, L., De Caprariis, P., Grandolini, G., Ambrogi, V., Perioli, L. "Synthesis and SAR study of imidazo[2,1-b]benzothiazole acids and some related compounds with anti-inflammatory and analgesic activities." *Farmaco*, **1996**, 51(7), 483-491.
- (4) Murugan, V., Kumar, S., Prabu, K. S., Suresh, B., Reddy, V. M. "Synthesis of some benzothiazole derivatives as possible alkylating agents." *Ind. J. Heterocycl. Chem.*, **2004**, 14(2), 175-176.
- (5) Paranjape S. D., Bobade A. S., Khadse B. G. "Synthesis and evaluation of new triazolobenzothiazole derivatives as potential anti-tubercular agents." *Arzneimittel-Forschung*, **2001**, 51(11), 916-919.
- (6) Jimonet, P., Audiau, F., Barreau, M., Blanchard, J. C., Boireau, A., Bour, Y., Coleno, M. A., Doble, A., Doerflinger, G., Do, H. C., Donat, M. H., Duchesne, J. M., Ganil, P., Gueremy, C., Honore, E., Just, B., Kerphirique, R., Gontier, S., Hubert, P., Laduron, P. M., Le, B. J., Meunier, M., Miquet, J. M., Nemecek, C., Pasquet, M., Piot, O., Pratt, J., Rataud, J., Reibaud, M., Stutzmann, J. M., Mignani, S. "Riluzole Series. Synthesis and in Vivo "Antiglutamate" Activity of 6-Substituted- 2-benzothiazolamines and 3-Substituted -2 -imino-benzothiazolines." *J. Med. Chem.*, **1999**, 42(15), 2828-2843.
- (7) Singh, S., Singh, J., Ingle, M., Mishra, R., Khadikar, P. V. "A QSAR study on carbonic anhydrase inhibition: predicting logKi(CAI) by using (SO₂NH₂) NMR chemical shift as a molecular descriptor." *Arkivoc*, **2006**, 16, 1-15.
- (8) Srivastava S. K., Yadav R., Srivastava S. D. "Synthesis and biological activity of 4-oxothiazolidines and their 5-arylidenes." *Ind. J. Chem.*, **2004**, 43B(2), 399-405.
- (9) Desbois Michel. "Process for the preparation of 4-fluoroanilines from 4-halonitrobenzenes." *Eur. Pat. Appl.*, 248746, **1987**, 6.
- (10) Biological assays, *Indian Pharmacopoeia Published by Govt. of India*, **1996**, 2, 88.

- (11) Patai, S. "The Chemistry of the carbon-nitrogen double bond." *John Wiley & Sons Ltd.*, London, **1970**.
- (12) Cimerman, Z., Miljanic, S., Galic, N. "Schiff bases derived from aminopyridines as spectrofluorimetric analytical reagents." *Croatica Chemica Acta.*, **2000**, 73, 81-86.
- (13) Sari, N., Arslan, S., Logoglu, E., Sakiyan, L. "Antibacterial activities of some new amino acid-Schiff bases." *J. Sci.*, **2003**, 16(2), 283-288.
- (14) Spinu, C., Kriza, A. "Co(II), Ni(II) and Cu(II) complexes of bidentate Schiff bases." *Acta Chim Slov.*, **2000**, 47(2), 179-185.
- (15) Harada, K. "In the Chemistry of the Carbon-Nitrogen Double Bond." ed. S. Patai, *Wiley, New York*, **1970**, 276.
- (16) Harada, K. "In the Chemistry of the Carbon-Nitrogen Double Bond." ed. S. Patai, *Wiley, New York*, **1970**, 266.
- (17) Sandhu, J. S., Sain, B. "Some recent advances in the chemistry of imines, in particular cycloaddition reactions." *Heterocycles*, **1987**, 26(3), 777-818.
- (18) Howard, A., Shiyamalie, A. "Metal carbonyl catalysis of organoborane reactions. Cobalt carbonyl catalyzed reductive carbonylation of Schiff bases." *J. Org. Chem.*, **1982**, 47(18), 3593-3595.
- (19) Deyrup, J. A. "Darzens aziridine synthesis." *J. Org. Chem.*, **1969**, 34(9), 2724-2727.
- (20) Bestmann, H. J., Seng, F. "Reactions with phosphinealkylenes. X. Reaction of triphenylphosphinealkylenes with Schiff bases." *Tetrah.*, **1965**, 21(6), 1373-1381.
- (21) Upadhyay, R. K., Mathur, S. P. "Effect of Schiff's bases as corrosion inhibitors on mild steel in sulphuric acid." *E-J. Chem.*, **2007**, 4(3), 408-414.
- (22) Euler, H. V., Hasselquist, H. "The structure and stability of Schiff bases and their intermediates." *Arkiv foer Kemi.*, **1953**, 6, 287-292.
- (23) Schroeder, C., Preis, S., Link, K. P. "Mechanism of formation of warfarin from 4-hydroxy-coumarin and anilinobenzylacetone. Evidence for Schiff base intermediates showing enhanced reactivity in the Michael reaction." *Tetrah. Lett.*, **1960**, 13, 23-29.
- (24) Pilecki, M., Marczewski, Z., Gora, J. "Schiff bases and their application in perfumery." *Kosmetyki*, **1984**, 28(11-12), 223-227.
- (25) Klayman, D. L., Scovill, J. P., Bartosevich, J. F., Bruce, J. "2-Acetylpyridine thiosemicarbazones. 5. 1-[1-(2-Pyridyl)ethyl]-3-thiosemicarbazides as potential antimalarial agents." *J. Med. Chem.*, **1983**, 26(1), 35-39.

- (26) Aliasghar, J., Dariush, K., Erik, De C., Chanaz, S., Jean, M. B. "Synthesis, Antibacterial, Antifungal and Antiviral Activity Evaluation of Some New bis-Schiff Bases of Isatin and Their Derivatives." *Molecules*, **2007**, 12, 1720-1730.
- (27) Yan, X., Caifeng, B., Yuhua, F., Cindy, C., Xia, Z., Ping, D. Q. "L-glutamin Schiff base copper complex as a proteasome inhibitor and a apoptosis inducer in human cancer cell." *Int. J. Onco.*, **2008**, 33, 1073-1079.
- (28) Pandeya, S. N., Sriram, D., Nath, G., DeClercq, E. "Synthesis, antibacterial, antifungal and anti-HIV activities of Schiff and Mannich bases derived from isatin derivatives and N-[4-(4'-chlorophenyl) thiazol-2-yl] thiosemicarbazide." *Eur. J. Pharm. Sci.*, **1999**, 9(1), 25-31.
- (29) Sriram, D., Yogeeswari, P., Kumar, T. G. "Microwave-assisted synthesis, anti- HIV, and anti-YFV activities of Schiff bases of N-hydroxy-N1-aminoguanidinetosylate." *Ind. J. Pharm. Sci.*, **2005**, 67(4), 493-496.
- (30) Kaushal, S., Chahal, K. K. "Schiff bases of Khusilal-synthesis and their antifungal activity." *Civil Lines Pesto.*, **2008**, 32(1), 47-49.
- (31) Rathelot, P., Azas, N., El-Kashef, H., Delmas, F., Di Giorgio, C., Timon, D. P., Maldonado, J., Vanelle, P. "1,3-Diphenylpyrazoles: synthesis and antiparasitic activities of azomethine derivatives." *Eur. J. Med. Chem.*, **2002**, 37(8), 671-679.
- (32) Glenn, H. H., Leslie, D. W. "Schiff base derivatives of anti-inflammatory o-substituted hydroxylamines." *J. Pharm. Sci.*, 2006, 60(6), 925-927.
- (33) Verma, M., Pandeya, S. N., Singh, K. N., Stables, J. P. "Anticonvulsant activity of Schiff bases of isatin derivatives." *Acta Pharm.*, **2004**, 54, 49-56.
- (34) Zhou, X., Shao, L., Jin, Z., Liu, J. B., Dai, H., Fang, J. X. "Synthesis and antitumor activity of some Schiff bases derived from 2-aminothiazole derivatives." *Heteroatom Chem.*, **2007**, 18(1), 55-59.
- (35) Zhong, X., Yi, J., Sun, J., Wei, H. L., Liu, W. S., Yu, K. B. "Synthesis and crystal structure of some transition metal complexes with a novel bis-schiff base ligand and their antitumor activities." *Eur. J. Med. Chem.*, **2006**, 41(9), 1090-1092.
- (36) Pontiki, E., Litina, D. H., Chavviara, A. T. "Evaluation of anti-inflammatory and antioxidant activities of Copper (II) Schiff mono-base and Copper(II) Schiff base coordination compounds of dien with heterocyclic aldehydes and 2-amino-5-methyl-thiazole." *J. Enzyme Inhi. Med. Chem.*, **2008**, 23(6), 1011-1017.

- (37) Mohamed, M. I., Hapipah, M. A., Mahmood, A. A., Robinson, W. T. "Synthesis, structural characterization, and anti-ulcerogenic activity of schiff base ligands derived from tryptamine and 5-chloro, 5-nitro, 3,5-ditertiarybutyl salicylaldehyde and their nickel(II), copper(II), and zinc(II) complexes." *Polyhedron*, **2009**, 28(18), 3993-3998.
- (38) Muslin, L., Roth, W., Erlenmeyer, H. "Metallic ions and biological activity. XIII. The tuberculostatic activity of several Schiff bases." *Helvet. Chim. Acta*, **1953**, 36, 886-890.
- (39) Popp, F. D. "Synthesis of Potential Anticancer Agents. II. Some Schiff bases." *J. Org. Chem.*, **1961**, 26(5), 1566-1568 (1961).
- (40) Demillo, A. B., Redfern, R. E. "New insect juvenile hormone mimics: aromatic Schiff bases and related compounds against the large milkweed bug and yellow mealworm." *J. Agric. Food Chem.*, **1979**, 27(4), 760-762.
- (41) Sprung, M. M. "A summary of the reactions of aldehydes with amines." *Chem. Rev.*, **1940**, 26, 297-338.
- (42) Hai, J. Y., Sun, W. H., Long, L. Z., Zhi, M. "The Rapid Synthesis of Schiff-base without Solvent under Microwave Irradiation." *Chin. Chem. Lett.*, **2002**, 13(1), 3-6.
- (43) Mistry, K., Desai, K. R. "Microwave assisted rapid and efficient synthesis of nitrogen and sulfur containing heterocyclic compounds and their pharmacological evaluation." *Ind. J. Chem.*, **2006**, 45B, 1762-1766.
- (44) Yuye, Y. "Synthesis of Schiff bases under ultrasound irradiation." *Asian J. Chem.*, **2007**, 19(3), 2476-2478.
- (45) Zhou, X., Wang, N., Ma, X. "Synthesis of Schiff base derivatives under ultrasound irradiation conditions." *Xinan Shifan Daxue Xuebao, Ziran Kexueban*, **2006**, 31(3), 62-65.
- (46) Liang-Xue, L., Liu, J. D., Wu, A. B., Qin, S. X., Li, Q., Deng, L. Z., Huang, X. L. "Synthesis under ultrasonic catalysis and structural characterization of dinuclear Zn(II)-macroheterocyclic complex." *Youji Huaxue*, **2004**, 24(3), 297-299.
- (47) Baluja, S., Kachhadia, N., Solanki, A., Movaliya, J. "Sonochemical synthesis of some Schiff bases." *Int. J. Chem. Sci.*, **2009**, 7(2), 976-980.
- (48) Corbin, J. L., Work, D. E. "Nitrogen-Sulfur ligand systems via reduction of Schiff's base - Zinc complexes derived from benzothiazoline." *Can. J. Chem.*, **1974**, 52, 1054-1058.
- (49) Hamil, A. M., Khalifa, K. M., AL-Houni, A., El-ajaily, M. M. "Synthesis, spectroscopic investigation and antiactivity activity of Schiff base

- complexes of cobalt (II) and copper (II) ions." *Rasayan J. Chem.*, **2009**, 2(2), 261-266.
- (50) Rajavel, R., Krishnan, C. N. "Studies on vanadyl complexes of Schiff bases derived from 2-aminobenzaldehyde." *Ori. J. Chem.*, **1998**, 14(2), 313-316.
- (51) Chavan, A. A., Pai, N. R. "Synthesis and biological activity of N-substituted - 3-chloro-2-azetidinones." *Molecules*, **2007**, 12, 2467-2477.
- (52) Vijay Kumar, M. M. J., Jayadevaiah, K. V., Nagaraja, T. S., Bharathi, D. R., Shameer, H., Jayachandran, E., Sreenivasa, G. M. "Synthesis, characterization and antimicrobial activity of new N-substituted-3-chloro-2-azetidinones." *Arch Pharm. Sci. Res.*, **2009**, 1(1), 31-39.
- (53) Mustafa, Y., Askin, K., Basaran, D. "Synthesis and antimicrobial activity of new crown ethers of Schiff base type." *J. Serb. Chem. Soc.*, **2007**, 72(3), 215-224.
- (54) Sie-Tiong, H., Teck-Ming, K., Siew-Teng, O., Guan-Yeow, Y., Hong-Cheu, L. "Synthesis and mesogenic properties of new Schiff bases comprising benzothiazole moiety." *13th int. Electronics Confer. On Synthetic Org. Chem.*, Nov., **2009**.
- (55) Zahid, H. C., Claudiu, T. S. "Antibacterial Co(II) AND Ni(II) complexes of benzothiazole-derived Schiff bases." *Synthesis Reactivity in Inorg., Metal-Org., Nano-Metal Chem.*, **2002**, 32(8), 1445-1461.

Chapter -3

Physico Chemical Properties



Section-I

Acoustical Properties

INTRODUCTION

Ultrasound waves are sound waves with a frequency greater than the upper limit of human hearing. Although this limit varies from person to person, it is approximately 20 kHz (20,000 hertz) and thus, 20 kHz serves as a useful lower limit in describing ultrasound. Some animals such as dogs, cats, dolphins, bats, and mice have an upper frequency limit that is greater than that of the human ear and thus can hear ultrasound.

Literature survey shows that these ultrasound waves have a number of applications in different fields¹⁻⁶ due to their non destructive nature. In medicine, it is used in non surgical removal of kidney stones⁷, treatment of intracranial tumor⁸, imaging fetal development during pregnancy⁹. In industry, these waves are used for large-scale cleaning¹⁰, emulsification of cosmetics etc¹¹. It is also used in food industry¹². In engineering^{13, 14}, these waves are used for welding plastics¹⁵, cutting alloys¹⁶ etc.

The study of chemical effects of ultrasound is a rapidly growing research area¹⁷⁻²³. Now a days, lots of interest has been generated on the use of ultrasound irradiation in synthetic organic chemistry, which cause decrease of reaction time, increase of yield, lower reaction temperature, avoidance of phase transfer catalysis etc^{24, 25}. Various workers have reported some organic reactions through ultrasound irradiation. Srinivasan et al^{26, 27} and Antunes et al²⁸ have reported some reactions using these waves.

This technique has also been used to study different materials²⁹, liquid crystals³⁰, cleavage³¹ and structure of solids³²⁻³⁵. It is also useful to the synthesis of biodiesel³⁶. Some of the most important aspects of sonochemistry have been its applications in the modification of both organic and inorganic materials^{37, 38}.

In physical chemistry, ultrasonic velocity measurements have been used to study the nature of molecular interactions in various binary³⁹⁻⁴², and ternary⁴³⁻⁴⁷, and liquid mixtures⁴⁸⁻⁵¹. However, little work has been done for solutions of solid organic compounds⁵²⁻⁵⁸. In recent years, various chemists synthesized different pyrimidine derivatives due to their biological importance but in our knowledge, no one has studied their acoustical properties.

Thus, with a view to understand the molecular interactions in solutions of dihydropyrimidinones, the present chapter deals with studies of various

acoustical and thermodynamic parameters of the solutions of dihydropyrimidinones in DMF and DMSO solutions at 298.15 K, over a wide range of concentration.

EXPERIMENTAL

The selected solvents DMF and DMSO for the present study are distilled by the reported procedure⁵⁹. The synthesized compounds dihydropyrimidinones were recrystallized before use.

The densities and ultrasonic velocities of solvents and solutions of dihydropyrimidinones of different concentrations were measured at 298.15 K by using pycnometer and single frequency ultrasonic interferometer operating at 2 MHz, with the uncertainties of 0.0001 g/cm³ and 0.01% respectively.

Density measurements:

The weight of distilled water, pure solvents and solutions of dihydropyrimidines were measured by using pycnometer. The densities (ρ) were evaluated by using following equation:

$$\rho(\text{g/cm}^3) = \frac{(\text{wt. of solvent or solution})(\text{density of water})}{(\text{wt. of water})} \quad \dots (3.1.1)$$

Viscosity Measurements:

To determine the viscosity of solution, Ubbelohde viscometer⁶⁰ was used, which obeys Stoke's law⁶¹. The measured quantity of the distilled water / solvent / solution was placed in the viscometer, which was suspended in a thermostat at 298.15 K. The digital stopwatch, with an accuracy of ± 0.01 sec was used to determine flow time of solutions. Using the flow times (t) and known viscosity of standard water sample, the viscosity of solvent and solutions were determined according to equation:

$$\frac{\eta_1}{\eta_2} = \frac{t_1 \rho_1}{t_2 \rho_2} \quad \dots (3.1.2)$$

Sound velocity measurement:

Ultrasonic interferometer (Mittal Enterprise, New Delhi, Model No. F-81) working at frequency of 2 MHz was used to determine sound velocity.

The solvent / solution were filled in the measuring cell with quartz crystal and then micrometer was fixed. The circulation of water from the thermostat at 298.15 K was started and test solvent / solution in the cell is allowed to thermally equilibrate. The micrometer was rotated very slowly so as to obtain a maximum or minimum of anode current (n). A number of maximum reading of anode current were counted. The total distance (d) travel by the

micrometer for $n=10$, was read. The wave length (λ) was determined according to the equation (3.1.3).

$$\lambda = \frac{2d}{n} \quad \dots (3.1.3)$$

The sound velocity (U) of solvent and solutions were calculated from the wavelength and frequency (F) according to equation (3.1.4).

$$U = \lambda F \quad \dots (3.1.4)$$

where, $F = 2 \times 10^6$ Hertz.

RESULTS AND DISCUSSION

Table 3.1.1 shows the experimental data of density (ρ), viscosity and sound velocity (U) of pure solvents and solutions of synthesized dihydropyrimidinones in DMF and DMSO at 298.15 K.

From these experimental data, various acoustical parameters like isentropic compressibility (κ_s), intermolecular free length (L_f), molar compressibility (W), Rao's molar sound function (R_m), Vander Waals constant (b), relaxation strength (r), apparent molar compressibility (ϕ_k) etc., were evaluated using the following equations:

1. Isentropic compressibility:

Isentropic compressibility (κ_s) was evaluated by Krishnamurty et al.⁶²:

$$\kappa_s = \frac{1}{U^2 \rho} \quad \dots (3.1.5)$$

2. Intermolecular free path length:

The intermolecular free path length (L_f), was calculated by equation⁶³:

$$L_f = K_J \kappa_s^{1/2} \quad \dots (3.1.6)$$

where K_J is Jacobson constant ($=2.0965 \times 10^{-6}$)

3. Molar compressibility:

Molar compressibility (W) can be calculated by the following equation⁶⁴:

$$W = \left(\frac{M}{\rho} \right) \kappa_s^{-1/7} \quad \dots (3.1.7)$$

The apparent molecular weight (M) of the solution can be calculated as:

$$M = M_1 W_1 + M_2 W_2 \quad \dots (3.1.8)$$

where W_1 and W_2 are weight fractions of solvent and solute, respectively. M_1 and M_2 are the molecular weights of the solvent and compounds respectively.

Table 3.1.1: The density (ρ), ultrasonic velocity (U) and viscosity (η) of Dihydropyrimidinones in DMF and DMSO at 298.15 K

| Conc. M | Density ρ $g.cm^{-3}$ | Velocity $U. 10^{-5}$ $cm.s^{-1}$ | Viscosity $\eta.10^3$ poise | Density ρ $g.cm^{-3}$ | Velocity $U. 10^{-5}$ $cm.s^{-1}$ | Viscosity $\eta.10^3$ poise |
|---------------------------------|---|--|---|---|--|---|
| | DMF | | | DMSO | | |
| | RVG-1 | | | RVG-1 | | |
| 0.00 | 0.9453 | 1.4516 | 8.4812 | 1.0959 | 1.4824 | 21.4335 |
| 0.01 | 0.9494 | 1.4552 | 9.2498 | 1.0971 | 1.4856 | 22.0260 |
| 0.02 | 0.9506 | 1.4580 | 9.3589 | 1.0982 | 1.4868 | 22.3210 |
| 0.04 | 0.9531 | 1.4604 | 9.7825 | 1.0989 | 1.4916 | 22.7872 |
| 0.06 | 0.9555 | 1.4656 | 10.0641 | 1.0998 | 1.4936 | 23.2742 |
| 0.08 | 0.9577 | 1.4676 | 10.2132 | 1.1004 | 1.4964 | 24.2054 |
| 0.10 | 0.9584 | 1.4688 | 10.3525 | 1.1012 | 1.4976 | 24.7118 |
| | RVG-2 | | | RVG-2 | | |
| 0.01 | 0.9485 | 1.4640 | 8.9859 | 1.0971 | 1.4844 | 22.2629 |
| 0.02 | 0.9498 | 1.4648 | 9.0944 | 1.0987 | 1.4872 | 22.7605 |
| 0.04 | 0.9516 | 1.4660 | 9.3217 | 1.0992 | 1.4904 | 23.3450 |
| 0.06 | 0.9536 | 1.4668 | 9.5184 | 1.1005 | 1.4932 | 23.7469 |
| 0.08 | 0.9554 | 1.4676 | 9.7231 | 1.1017 | 1.4948 | 24.0041 |
| 0.10 | 0.9571 | 1.4680 | 9.9205 | 1.1023 | 1.4964 | 24.0916 |
| | RVG-3 | | | RVG-3 | | |
| 0.01 | 0.9488 | 1.4548 | 9.0517 | 1.0965 | 1.4840 | 22.4517 |
| 0.02 | 0.949 | 1.4568 | 9.2779 | 1.0973 | 1.4852 | 22.7831 |
| 0.04 | 0.952 | 1.4584 | 9.5748 | 1.0994 | 1.4860 | 23.3055 |
| 0.06 | 0.9527 | 1.4604 | 9.7267 | 1.1009 | 1.4876 | 23.7051 |
| 0.08 | 0.9543 | 1.4612 | 9.9053 | 1.101 | 1.4904 | 24.0499 |
| 0.10 | 0.9565 | 1.4628 | 10.1104 | 1.1026 | 1.4924 | 24.5650 |
| | RVG-4 | | | RVG-4 | | |
| 0.01 | 0.947 | 1.4536 | 9.5794 | 1.0974 | 1.4836 | 21.9817 |
| 0.02 | 0.9474 | 1.4556 | 9.7583 | 1.0982 | 1.4860 | 22.3051 |
| 0.04 | 0.9478 | 1.4572 | 9.8767 | 1.0992 | 1.4916 | 22.7033 |
| 0.06 | 0.9484 | 1.4588 | 10.0145 | 1.0996 | 1.4908 | 23.0670 |
| 0.08 | 0.9489 | 1.4604 | 10.1159 | 1.1010 | 1.4928 | 23.3952 |
| 0.10 | 0.9505 | 1.4624 | 10.3497 | 1.1027 | 1.4956 | 23.7239 |
| | RVG-5 | | | RVG-5 | | |
| 0.01 | 0.9494 | 1.4536 | 9.0608 | 1.0969 | 1.4860 | 21.9783 |
| 0.02 | 0.9517 | 1.4548 | 9.2320 | 1.0985 | 1.4876 | 22.3748 |
| 0.04 | 0.953 | 1.4568 | 9.6159 | 1.0996 | 1.4888 | 22.8601 |
| 0.06 | 0.9555 | 1.4580 | 9.8382 | 1.1003 | 1.4904 | 23.1175 |
| 0.08 | 0.9577 | 1.4592 | 9.9568 | 1.1017 | 1.4920 | 23.3330 |
| 0.10 | 0.9597 | 1.4612 | 10.1686 | 1.1026 | 1.4964 | 23.5928 |

| Conc. M | Density ρ g.cm⁻³ | Velocity U. 10⁻⁵ cm.s⁻¹ | Viscosity η.10³ poise | Density ρ g.cm⁻³ | Velocity U. 10⁻⁵ cm.s⁻¹ | Viscosity η.10³ poise |
|--------------------|--|--|---|--|--|---|
| | DMF | | | DMSO | | |
| | RVG-6 | | | RVG-6 | | |
| 0.00 | 0.9487 | 1.4572 | 9.1114 | 1.0959 | 1.4824 | 21.4335 |
| 0.01 | 0.9505 | 1.4608 | 9.3522 | 1.0972 | 1.4836 | 21.8904 |
| 0.02 | 0.9518 | 1.4636 | 9.4626 | 1.0982 | 1.4852 | 22.4203 |
| 0.04 | 0.9547 | 1.4656 | 9.6089 | 1.0995 | 1.4860 | 22.8885 |
| 0.06 | 0.9554 | 1.4672 | 9.7611 | 1.1007 | 1.4872 | 23.2733 |
| 0.08 | 0.9566 | 1.4684 | 10.2176 | 1.1011 | 1.4884 | 23.4584 |
| 0.10 | 0.9487 | 1.4572 | 9.1114 | 1.1028 | 1.4896 | 23.5611 |
| | RVG-7 | | | RVG-7 | | |
| 0.01 | 0.9485 | 1.4536 | 9.0957 | 1.0978 | 1.4848 | 21.9738 |
| 0.02 | 0.95 | 1.4552 | 9.2487 | 1.0984 | 1.4864 | 22.1992 |
| 0.04 | 0.9515 | 1.4564 | 9.3988 | 1.0998 | 1.4888 | 22.6825 |
| 0.06 | 0.9528 | 1.4596 | 9.6162 | 1.1004 | 1.4900 | 22.8767 |
| 0.08 | 0.9535 | 1.4608 | 9.8153 | 1.1011 | 1.4916 | 23.3176 |
| 0.10 | 0.9546 | 1.4628 | 9.9936 | 1.1028 | 1.4936 | 23.6556 |
| | RVG-8 | | | RVG-8 | | |
| 0.01 | 0.9505 | 1.4524 | 9.4348 | 1.0986 | 1.4860 | 22.2297 |
| 0.02 | 0.9523 | 1.4540 | 9.7076 | 1.0995 | 1.4880 | 22.5092 |
| 0.04 | 0.9558 | 1.4564 | 9.9335 | 1.1003 | 1.4896 | 22.8667 |
| 0.06 | 0.9581 | 1.4592 | 10.1655 | 1.1011 | 1.4908 | 23.3389 |
| 0.08 | 0.9587 | 1.4600 | 10.1857 | 1.103 | 1.4928 | 23.5840 |
| 0.10 | 0.9595 | 1.4616 | 10.6120 | 1.1044 | 1.4948 | 23.9550 |
| | RVG-9 | | | RVG-9 | | |
| 0.01 | 0.9508 | 1.4540 | 9.2370 | 1.0982 | 1.4872 | 22.3634 |
| 0.02 | 0.9524 | 1.4548 | 9.4858 | 1.0986 | 1.4888 | 22.7426 |
| 0.04 | 0.9558 | 1.4564 | 9.5900 | 1.0998 | 1.4900 | 23.1083 |
| 0.06 | 0.9576 | 1.4572 | 9.8876 | 1.1006 | 1.4916 | 23.5672 |
| 0.08 | 0.9605 | 1.4592 | 10.0484 | 1.1013 | 1.4932 | 23.8665 |
| 0.10 | 0.9617 | 1.4608 | 10.1376 | 1.1021 | 1.4952 | 24.3731 |
| | RVG-10 | | | RVG-10 | | |
| 0.01 | 0.9514 | 1.4548 | 8.9812 | 1.097 | 1.4836 | 21.8745 |
| 0.02 | 0.9536 | 1.4568 | 9.1308 | 1.0983 | 1.4848 | 22.1640 |
| 0.04 | 0.9558 | 1.4580 | 9.4032 | 1.0991 | 1.4856 | 22.3949 |
| 0.06 | 0.9578 | 1.4592 | 9.6990 | 1.0999 | 1.4880 | 22.7350 |
| 0.08 | 0.959 | 1.4616 | 9.8106 | 1.1007 | 1.4892 | 23.4778 |
| 0.10 | 0.9602 | 1.4628 | 10.0731 | 1.1013 | 1.4912 | 23.7483 |

4. Rao's molar sound function:

Rao's molar sound function (R_m) can be evaluated by an equation given by Bagchi et al.⁶⁵:

$$R_m = \left(\frac{M}{\rho} \right) U^{1/3} \quad \dots (3.1.9)$$

5. Van der Waals Constant:

Van der Waals constant (b) can be calculated as follows⁶⁶:

$$b = \frac{M}{\rho} \left\{ 1 - \left(\frac{RT}{MU^2} \right) \left[\sqrt{1 + \frac{MU^2}{3RT}} - 1 \right] \right\} \quad \dots (3.1.10)$$

where R is the gas constant ($=8.3143 \text{ JK}^{-1} \text{ mol}^{-1}$) and T is the absolute temperature.

6. Relaxation Strength:

The relaxation strength (r) can be calculated as follows⁶⁷:

$$r = 1 - \left[\frac{U}{U_\infty} \right]^2 \quad \dots (3.1.11)$$

where $U_\infty = 1.6 \times 10^5 \text{ cm/sec}$.

7. Solvation number:

The solvation number (S_n) can be calculated according to the equation⁶⁸

$$S_n = \frac{M_2}{M_1} \left[\frac{1 - \kappa_s}{\kappa_{s1}} \right] \left[\frac{100 - X}{X} \right] \quad \dots (3.1.12)$$

where X is the number of grams of solute in 100 gm of the solution. M_1 and M_2 are the molecular weights and κ_{s1} and κ_s are isentropic compressibility of solvent and solute respectively.

8. Apparent Molar Compressibility (ϕ_k):

The apparent molar compressibility (ϕ_k) of the solutions was calculated by the following equation⁶⁹:

$$\phi_K = \frac{(\rho_0 \kappa_s - \rho \kappa_s^0) 1000}{c \rho_0} + \frac{\kappa_s^0 M_2}{\rho_0} \quad \dots (3.1.13)$$

where ρ_0 and κ_s^0 are density and isentropic compressibility of pure solvent respectively, c is the concentration of the solution and M_2 is the molecular weight of the compound.

Some of these calculated parameters are given in Table 3.1.2 for the studied compounds in DMF and DMSO respectively. Figure 3.1.1 show the variation of ultrasound velocity (U) with concentration in DMF and DMSO solutions. It is observed that overall ultrasonic velocity (U) increases with concentration for all the compounds in both the solvents. The velocity depends on intermolecular free length (L_f). The velocity increases with decrease in L_f or vice versa. It is evident from Table 3.1.2 that L_f decreases continuously, which suggests that there is strong interaction between solvent (both DMF and DMSO) and compound molecules.

This is further supported by isentropic compressibility (κ_s) and relaxation strength (r) values. The variation of isentropic compressibility (κ_s) with concentration of these compounds is also shown Figure 3.1.2 for all the solutions in both the solvents. The nature of curves is opposite to those of velocity (Fig. 3.1.1.). Both isentropic compressibility (κ_s) and relaxation strength (r) (Table 3.1.3) are observed to decrease with concentration for all the compounds. The decrease in isentropic compressibility is attributed to the fact that the dihydropyrimidinones (DHPMs) molecules in dilute solutions are considerably ionized and these ions are firmly bound to surrounding solvent molecules. The orientation of solvent molecules around the ions is attributed to the influence of the electrostatic field of the ions, which lowers the compressibility of the DHPMs solutions⁷⁰.

Figure 3.1.3 shows the linear variation of molar compressibility (W) with concentration whereas Table 3.1.3 show the variation of molar sound function (R_m) and Vander Waals constant (b) with concentration. All the three parameters varies linearly with concentration and the correlation coefficients are in the range of 0.9991- 0.9999. This linear change of these parameters suggests the absence of complex formation in these systems.

Table 3.1.3: Some acoustical parameters of Dihydropyrimidinones derivatives in DMF and DMSO at 298.15 K.

| DMF | | | | | DMSO | | | |
|------------|-------------|--------|---|--|-------------|--------|---|--|
| Conc. M | L_f A° | r | $R_m \cdot 10^{-3}$ $\text{cm}^{-8/3} \cdot \text{s}^{-1/3}$ | b $\text{cm}^3 \cdot \text{mol}^{-1}$ | L_f A° | r | $R_m \cdot 10^{-3}$ $\text{cm}^{-8/3} \cdot \text{s}^{-1/3}$ | b $\text{cm}^3 \cdot \text{mol}^{-1}$ |
| RVG-1 | | | | | RVG-1 | | | |
| 0.00 | 0.1485 | 0.1769 | 4.0635 | 77.3176 | 0.1351 | 0.1416 | 4.5458 | 85.8912 |
| 0.01 | 0.1479 | 0.1728 | 4.1244 | 78.4103 | 0.1347 | 0.1379 | 4.5970 | 86.7962 |
| 0.02 | 0.1475 | 0.1696 | 4.1966 | 79.7326 | 0.1346 | 0.1365 | 4.6464 | 87.7052 |
| 0.04 | 0.1470 | 0.1669 | 4.3366 | 82.3471 | 0.1341 | 0.1309 | 4.7540 | 89.6395 |
| 0.06 | 0.1463 | 0.1609 | 4.4786 | 84.9431 | 0.1338 | 0.1286 | 4.8575 | 91.5510 |
| 0.08 | 0.1460 | 0.1587 | 4.6172 | 87.5326 | 0.1336 | 0.1253 | 4.9632 | 93.4838 |
| 0.10 | 0.1458 | 0.1573 | 4.7624 | 90.2604 | 0.1334 | 0.1239 | 5.0659 | 95.3932 |
| RVG-2 | | | | | RVG-2 | | | |
| 0.01 | 0.1470 | 0.1628 | 4.1301 | 78.3607 | 0.1348 | 0.1393 | 4.5910 | 86.7053 |
| 0.02 | 0.1469 | 0.1619 | 4.1936 | 79.5521 | 0.1345 | 0.1360 | 4.6351 | 87.4831 |
| 0.04 | 0.1466 | 0.1605 | 4.3232 | 81.9877 | 0.1342 | 0.1323 | 4.7322 | 89.2517 |
| 0.06 | 0.1464 | 0.1596 | 4.4504 | 84.3850 | 0.1338 | 0.1290 | 4.8251 | 90.9469 |
| 0.08 | 0.1461 | 0.1587 | 4.5776 | 86.7819 | 0.1336 | 0.1272 | 4.9168 | 92.6429 |
| 0.10 | 0.1460 | 0.1582 | 4.7041 | 89.1708 | 0.1334 | 0.1253 | 5.0112 | 94.3876 |
| RVG-3 | | | | | RVG-3 | | | |
| 0.01 | 0.1479 | 0.1733 | 4.1283 | 78.4930 | 0.1349 | 0.1397 | 4.5992 | 86.8676 |
| 0.02 | 0.1477 | 0.1710 | 4.2062 | 79.9362 | 0.1348 | 0.1384 | 4.6511 | 87.8255 |
| 0.04 | 0.1473 | 0.1692 | 4.3467 | 82.5768 | 0.1346 | 0.1374 | 4.7507 | 89.6901 |
| 0.06 | 0.1471 | 0.1669 | 4.4979 | 85.4093 | 0.1343 | 0.1356 | 4.8532 | 91.5929 |
| 0.08 | 0.1469 | 0.1660 | 4.6425 | 88.1393 | 0.1341 | 0.1323 | 4.9635 | 93.6143 |
| 0.10 | 0.1465 | 0.1641 | 4.7835 | 90.7845 | 0.1338 | 0.1300 | 5.0652 | 95.4911 |
| RVG-4 | | | | | RVG-4 | | | |
| 0.01 | 0.1482 | 0.1746 | 4.1269 | 78.4869 | 0.1349 | 0.1402 | 4.5889 | 86.6813 |
| 0.02 | 0.1480 | 0.1724 | 4.1958 | 79.7602 | 0.1346 | 0.1374 | 4.6360 | 87.5238 |
| 0.04 | 0.1478 | 0.1705 | 4.3329 | 82.3373 | 0.1341 | 0.1309 | 4.7334 | 89.2517 |
| 0.06 | 0.1476 | 0.1687 | 4.4689 | 84.8904 | 0.1341 | 0.1318 | 4.8266 | 91.0258 |
| 0.08 | 0.1474 | 0.1669 | 4.6052 | 87.4473 | 0.1338 | 0.1295 | 4.9180 | 92.7064 |
| 0.10 | 0.1470 | 0.1646 | 4.7354 | 89.8795 | 0.1335 | 0.1262 | 5.0083 | 94.3501 |
| RVG-5 | | | | | RVG-5 | | | |
| 0.01 | 0.1480 | 0.1746 | 4.1229 | 78.4103 | 0.1347 | 0.1374 | 4.5983 | 86.8122 |
| 0.02 | 0.1477 | 0.1733 | 4.1885 | 79.6371 | 0.1345 | 0.1356 | 4.6460 | 87.6807 |
| 0.04 | 0.1474 | 0.1710 | 4.3335 | 82.3564 | 0.1343 | 0.1342 | 4.7479 | 89.5799 |
| 0.06 | 0.1471 | 0.1696 | 4.4708 | 84.9431 | 0.1341 | 0.1323 | 4.8517 | 91.5067 |
| 0.08 | 0.1468 | 0.1683 | 4.6084 | 87.5326 | 0.1339 | 0.1304 | 4.9520 | 93.3641 |
| 0.10 | 0.1465 | 0.1660 | 4.7467 | 90.1192 | 0.1334 | 0.1253 | 5.0575 | 95.2595 |

| DMF | | | | | DMSO | | | |
|-------------------|--------------------|----------|---|------------------------------|--------------------|----------|---|------------------------------|
| Conc. <i>M</i> | L_f A° | <i>r</i> | $R_m \cdot 10^{-3}$ $cm^{-8/3} \cdot s^{-1/3}$ | b $cm^3 \cdot mol^{-1}$ | L_f A° | <i>r</i> | $R_m \cdot 10^{-3}$ $cm^{-8/3} \cdot s^{-1/3}$ | b $cm^3 \cdot mol^{-1}$ |
| RVG-6 | | | | | RVG-6 | | | |
| 0.00 | 0.1477 | 0.1769 | 4.0635 | 77.3176 | 0.1351 | 0.1416 | 4.5458 | 85.8912 |
| 0.01 | 0.1472 | 0.1705 | 4.1311 | 78.5014 | 0.1349 | 0.1402 | 4.5958 | 86.8116 |
| 0.02 | 0.1468 | 0.1664 | 4.2031 | 79.8055 | 0.1347 | 0.1384 | 4.6472 | 87.7519 |
| 0.04 | 0.1464 | 0.1632 | 4.3528 | 82.5954 | 0.1345 | 0.1374 | 4.7503 | 89.6815 |
| 0.06 | 0.1462 | 0.1609 | 4.4928 | 85.2122 | 0.1344 | 0.1360 | 4.8538 | 91.6107 |
| 0.08 | 0.1460 | 0.1591 | 4.6428 | 88.0245 | 0.1342 | 0.1346 | 4.9608 | 93.6051 |
| 0.10 | 0.1477 | 0.1577 | 4.7891 | 90.7735 | 0.1340 | 0.1332 | 5.0610 | 95.4719 |
| RVG-7 | | | | | RVG-7 | | | |
| 0.01 | 0.1481 | 0.1746 | 4.1219 | 78.3915 | 0.1348 | 1.0978 | 4.5896 | 86.6717 |
| 0.02 | 0.1478 | 0.1728 | 4.1867 | 79.5962 | 0.1346 | 1.0984 | 4.6379 | 87.5520 |
| 0.04 | 0.1476 | 0.1714 | 4.3206 | 82.1193 | 0.1343 | 1.0998 | 4.7325 | 89.2899 |
| 0.06 | 0.1471 | 0.1678 | 4.4568 | 84.6456 | 0.1341 | 1.1004 | 4.8291 | 91.0888 |
| 0.08 | 0.1470 | 0.1664 | 4.5936 | 87.2192 | 0.1339 | 1.1011 | 4.9256 | 92.8746 |
| 0.10 | 0.1467 | 0.1641 | 4.7286 | 89.7421 | 0.1337 | 1.1028 | 5.0173 | 94.5617 |
| RVG-8 | | | | | RVG-8 | | | |
| 0.01 | 0.1481 | 0.1760 | 4.1341 | 78.6460 | 0.1346 | 0.1374 | 4.6037 | 86.9151 |
| 0.02 | 0.1478 | 0.1742 | 4.2194 | 80.2387 | 0.1344 | 0.1351 | 4.6673 | 88.0759 |
| 0.04 | 0.1472 | 0.1714 | 4.3878 | 83.3962 | 0.1342 | 0.1332 | 4.7960 | 90.4724 |
| 0.06 | 0.1468 | 0.1683 | 4.5607 | 86.6268 | 0.1340 | 0.1318 | 4.9240 | 92.8619 |
| 0.08 | 0.1467 | 0.1673 | 4.7397 | 90.0092 | 0.1337 | 0.1295 | 5.0470 | 95.1396 |
| 0.10 | 0.1464 | 0.1655 | 4.9179 | 93.3600 | 0.1335 | 0.1272 | 5.1719 | 97.4507 |
| RVG-9 | | | | | RVG-9 | | | |
| 0.01 | 0.1479 | 0.1742 | 4.1343 | 78.6206 | 0.1345 | 0.1360 | 4.6067 | 86.9472 |
| 0.02 | 0.1477 | 0.1733 | 4.2197 | 80.2300 | 0.1344 | 0.1342 | 4.6721 | 88.1501 |
| 0.04 | 0.1472 | 0.1714 | 4.3878 | 83.3962 | 0.1342 | 0.1328 | 4.7987 | 90.5158 |
| 0.06 | 0.1470 | 0.1705 | 4.5613 | 86.6774 | 0.1340 | 0.1309 | 4.9273 | 92.9074 |
| 0.08 | 0.1466 | 0.1683 | 4.7286 | 89.8147 | 0.1338 | 0.1290 | 5.0561 | 95.3016 |
| 0.10 | 0.1463 | 0.1664 | 4.9037 | 93.1072 | 0.1336 | 0.1267 | 5.1846 | 97.6796 |
| RVG-10 | | | | | RVG-10 | | | |
| 0.01 | 0.1477 | 0.1733 | 4.1247 | 78.4231 | 0.1349 | 0.1402 | 4.6024 | 86.9361 |
| 0.02 | 0.1474 | 0.1710 | 4.2007 | 79.8323 | 0.1347 | 0.1388 | 4.6578 | 87.9603 |
| 0.04 | 0.1471 | 0.1696 | 4.3588 | 82.8142 | 0.1346 | 0.1379 | 4.7746 | 90.1483 |
| 0.06 | 0.1468 | 0.1683 | 4.5165 | 85.7878 | 0.1343 | 0.1351 | 4.8927 | 92.3297 |
| 0.08 | 0.1465 | 0.1655 | 4.6788 | 88.8204 | 0.1342 | 0.1337 | 5.0093 | 94.5048 |
| 0.10 | 0.1463 | 0.1641 | 4.8390 | 91.8377 | 0.1340 | 0.1314 | 5.1276 | 96.6930 |

Figure 3.1.1: Variation of ultrasonic velocity (U) of Dihydropyrimidinones with concentration in [A] DMF and [B] DMSO at 298.15 K.




Figure 3.1.2: Variation of Isentropic compressibility (κ_s) of Dihydropyrimidinones with concentration in [A] DMF and [B] DMSO at 298.15 K.





Figure 3.1.3: Variation of Molar compressibility (W) of Dihydropyrimidinones with concentration in [A] DMF and [B] DMSO at 298.15 K.



The interactions occurring in different solutions can also be confirmed by the solvation number, which is measure of structure forming or structure breaking tendency. Figure 3.1.4 show the variation of solvation number with concentration. For the studied compounds, S_n values are positive at low concentrations. The positive S_n is due to structure forming tendency whereas negative S_n suggests structure breaking tendency. In both the solvents, solvation number is found to increase with increasing concentration. However, in DMSO solutions, slight decrease in S_n is observed for RVG-2 and RVG-4 whereas for others there is increase at lower concentrations Thus, the overall finding suggests that the studied compounds exhibit solute-solvent interactions i.e. structure forming tendency of these compounds in both the solvents.

Figure 3.1.4: Variation of Solvation Number (S_n) of Dihydropyrimidinones with concentration in [A] DMF and [B] DMSO at 298.15 K.




Table 3.1.4: Bachem's, Gucker's and Masson's constants of Dihydro pyrimidinones in DMF and DMSO at 298.15 K.

| Compounds | $A \times 10^{11}$ dyn^{-1} cm^{-3} $\cdot \text{mol}^{-1}$ | $B \times 10^{11}$ dyn^{-1} $\text{cm}^{-1/2}$ $\cdot \text{mol}^{-3/2}$ | $\phi_K^\circ \times 10^8$ dyn^{-1} $\cdot \text{mol}^{-1}$ | $S_K \times 10^8$ dyn^{-1} $\text{cm}^{-3/2}$ $\cdot \text{mol}^{-3/2}$ | ϕ_v° $\text{cm}^3 \cdot \text{mol}^{-1}$ | S_v $\text{cm}^3 \cdot \text{mol}^{-1}$ |
|---------------|--|--|--|---|---|--|
| DMF | | | | | | |
| <i>RVG-1</i> | -4.629 | 8.7972 | -3.3025 | 8.8136 | -252.31 | 1804.3 |
| <i>RVG-2</i> | -4.8972 | 10.098 | -39784 | 11.617 | -113.15 | 817.66 |
| <i>RVG-3</i> | -3.7944 | 8.242 | -3.0021 | 10.745 | -134 | 1205.1 |
| <i>RVG-4</i> | -2.7633 | 6.2117 | -2.0558 | 10.839 | -6.4976 | 590.72 |
| <i>RVG-5</i> | -3.6667 | 7.7383 | -2.0841 | 6.5901 | -155.94 | 988.95 |
| <i>RVG-6</i> | -4.9163 | 10.2250 | -4.1618 | 12.6530 | -251.14 | 2563.6 |
| <i>RVG-7</i> | -3.6717 | 8.3331 | -1.7364 | 6.5383 | -124.77 | 1227.8 |
| <i>RVG-8</i> | -3.9725 | 8.3206 | -3.8210 | 12.6430 | -322.34 | 2859.3 |
| <i>RVG-9</i> | -4.0006 | 8.4545 | -2.0437 | 6.2675 | -252.31 | 1804.3 |
| <i>RVG-10</i> | -3.9775 | 7.874 | -4.1675 | 13.1040 | -268.76 | 2137.2 |
| DMSO | | | | | | |
| <i>RVG-1</i> | -2.742 | 5.6196 | -1.3065 | 5.0384 | -25.027 | 819.51 |
| <i>RVG-2</i> | -1.8766 | 2.5988 | -0.9653 | 3.5489 | -0.5184 | 359.72 |
| <i>RVG-3</i> | -1.3760 | 2.4847 | -0.0474 | 1.8949 | -7.6917 | 419.02 |
| <i>RVG-4</i> | -1.3568 | 1.3155 | -0.6134 | 2.9043 | -57.559 | 1632.2 |
| <i>RVG-5</i> | -2.8254 | 7.1072 | -2.1338 | 9.1884 | -10.126 | 450.73 |
| <i>RVG-6</i> | -1.4248 | 2.7656 | -0.0934 | 2.2183 | -36.20 | 922.44 |
| <i>RVG-7</i> | -2.2937 | 5.1411 | -1.5367 | 6.7903 | -50.07 | 1229.5 |
| <i>RVG-8</i> | -2.1401 | 3.7012 | -0.1466 | 1.4902 | -29.19 | 514.72 |
| <i>RVG-9</i> | -2.218 | 4.1876 | -0.6994 | 3.7762 | -6.98 | 492.23 |
| <i>RVG-10</i> | -1.2705 | 1.9472 | -0.1342 | 2.8083 | -0.52 | 495.86 |

The isentropic compressibility of all the solutions was also fitted to the following Bachem's relation⁷¹:

$$\kappa_s = \kappa_{s1} + AC + BC^{3/2} \quad \dots (3.1.14)$$

where A and B are constants, C is the molar concentration of DHPMs, and κ_s and κ_{s1} are the adiabatic compressibility of the solution and solvent respectively. The constants A and B have been determined from the intercept and slope of the plots of $(\kappa_s - \kappa_{s1})/C$ vs. $C^{1/2}$

Further, apparent molar compressibility and apparent molar volume of solutions are fitted to Gucker's⁷² and Masson's⁷³ relations:

$$\phi_k = \phi_k^\circ + S_k C^{1/2} \quad \dots (3.1.15)$$

$$\phi_v = \phi_v^\circ + S_v C^{1/2} \quad \dots (3.1.16)$$

where ϕ_k° and ϕ_v° are limiting apparent molar compressibility and limiting apparent molar volume, respectively. S_k and S_v are parameters for solute-solvent interactions. The values of ϕ_k° , ϕ_v° , and constants S_k and S_v , have been obtained from the intercept and slope of the plots of ϕ_k vs $C^{1/2}$ and ϕ_v vs $C^{1/2}$. All these constants are given in Table 3.1.4.

It is observed that for all the compounds in both the solvents, A values are negative whereas B values are positive. The negative A and positive B further confirms solute-solvent interactions in the studied system. This is further confirmed by negative ϕ_k° and ϕ_v° values.

The positive S_k and S_v confirms the existence of solute-solvent interactions in the studied systems.

Thus, it is concluded that DMF and DMSO solutions of dihydropyrimidinones exhibit solute-solvent interactions.

LITERATURE CITED

- (1) Iwasaki, T., Lindberg, B., Meier, H. "The effect of ultrasonic treatment on individual wood fibers." *Svensk Pappersstidning*, **1962**, 65, 795-816.
- (2) Mehrotra, K. N., Upadhyaya, S. K. "Acoustical studies of calcium soaps." *Acoustics Lett.*, **1987**, 11, 66-71.
- (3) Singh, R. K., Singh, R. P., Singh, M. P. "Acoustical characterization of nanostructured metal." *Inter. J. Nanosci.*, **2008**, 7, 315-323.
- (4) Lum, I., Huang, H., Chang, B. H., Mayer, M., Du, D., Zhou, Y. "Effects of superimposed ultrasound on deformation of gold." *J. App. Phy.*, **2009**, 105, 1-5.
- (5) Perincek, S., Uzgur, A. E., Duran, K., Dogan, K., Dogan, A., Korlu, A. E., Bahtiyari, I. M., Emel, A. "Design parameter investigation of industrial size ultrasound textile treatment bath." *Ultra. Sonochem.*, **2009**, 16, 184-189.
- (6) Savaroglu, G., Aral, E. "Acoustical parameters of cow's milk added hydrogen peroxide and sodium bicarbonate at different temperatures." *J. Food Eng.*, **2007**, 79, 287-292.
- (7) George, E. B. William, H. B. "Ultrasonic destruction of kidney stones" *West J. Med.*, **1984**, 140, 227-232.
- (8) Fasano, V. A., Zeme, S., Frego, L., Gunetti, R. "Ultrasonic aspiration in the surgical treatment of intracranial tumors." *J. Neurosur. Sci.*, **1981**, 25, 35-40.
- (9) Busch, W., Staffeldt, K. G. "early diagnosis of molar pregnancy by ultrasonic investigation and HCG [Human chorionic gonadotropin] excretion." *Fertil. Steril., Proc. World. Congr. 7th*, **1973**, 831.
- (10) Clementson, J. J., Mond, D. "Ultrasonic cleaning with fluorocarbon solvent/water systems." *Ultra. Ind. Conf. Pap.*, **1969**, 40-43.
- (11) Burfeind, D. B. "New ultrasound technology touted for facial and neck tissue tightening". *Dermatology Nursing*, **2007**, 5, 451-454.
- (12) McClements, D. J. "Ultrasonics in Food Processing" *Food Processing Recent Developments*, **1995**, 2, 59-70.
- (13) Massimiliano, P., Bruno, L., Antonio, B. "Ultrasonic assessment of wear-induced modifications in engineering contacts" *17th Int. Conf. on Wear of Materials.*, **2009**, 267, 1117-1122.


- (14) Coffey, J. M., Oates, G., Whittle, M. J. "The Future of Ultrasonic Examination in Engineering" *Phil. Trans. R. Soc. Lond. A.*, **1979**, 292, 285-298.
- (15) Mori, E., Kaneko, S., Gakumazawa, M., Okawa, Y. "Ultrasonic welding of plastics." *Ultra. Int. Conf. Proc.*, **1973**, 6, 16-19.
- (16) Li, P., Yost, W. Cantrell, J. H., Salama, K. "Dependence of acoustic nonlinearity parameter on second phase precipitates of aluminum alloys." *Ultra. Symp. Proc.*, **1985**, 2, 1113-1115.
- (17) Suslick, K. S, "Ultrasound: Its Chemical, Physical, and Biological Effects." New York: VCH, **1988**.
- (18) Mason, T. J., Lorimer, J. P. "Sonochemistry: Theory, Applications and Uses of Ultrasound in Chemistry." *Chichester, UK: Harword.* **1988**.
- (19) Van Eldik, R., Hubbard, C. C., "Chemistry under extreme or non-classical conditions." *New York: Wiley & Sons.* **1997**.
- (20) Suslick, K. S., Crum, L. A. "In Encyclopedia of Acoustics, ed." *New York, Wiley-Interscience.*, **1997**, 1, 271-282.
- (21) Luche, J. L. "Synthetic organic sonochemistry." *NewYork, Plenum.*, **1998**.
- (22) Suslick, K. S. "In Kirk-Othmer Encyclopedia of Chemical Technology." *New York: Wiley & Sons, 4th ed.* **1998**, 26, 517-541.
- (23) Crum, L. A., Mason, T. J., Reisse, J. L., Suslick, K. S. "Sonochemistry and Sonoluminescence." *Proc. NATO Adv. Study Inst. Ser. C. Netherlands, Kluwer*, **1999**, 524.
- (24) Tramontana, M., Gachagan, A., Hayward, G., Nordon, A., Littlejohn, D. "Ultrasonic monitoring of heterogeneous chemical reactions." *Proce. - IEEE Ultra. Sympo.*, **2006**, 2, 910-913.
- (25) Price, G. J. "Current trends in Sonochemistry" *Royal Soc. Chem.*, Cambridge., **1992**.
- (26) Srinivasan, K. V., Rajagopal, R., Jarikote, V. "Ultrasound promoted Suzuki cross-coupling reactions in ionic liquid at ambient conditions." *Chem. Commun.*, **2001**, 17, 1544-1545.
- (27) Srinivasan, K. V., Rajagopal, R., Deshmukh, R. R. "Ultrasound promoted C-C bond formation: Heck reaction at ambient conditions in room temperature ionic liquids." *Chem. Commun.*, **2002**, 6, 616-617.
- (28) Andréa L. F. S., Lucyane, C. S., Bianca, L. O., Antunes, O. A. C. "Microwave- and ultrasound-assisted Suzuki-Miyaura cross-coupling reactions catalyzed by Pd/PVP." *Tetrah. Lett.*, **2008**, 49, 3895-3898.
- (29) Kamioka, H. "Effects of ultrasonic bonding materials on velocity and attenuation of sound in red lauan wood." *Jap. J. App. Phy.*, **1988**, 27,

- 188-191.
- (30) Langevin, D., Bouchiat, M. A. "Light scattering from the free surface of a nematic liquid crystal." *J. de Phy.*, **1972**, 33, 77-80.
- (31) Zhilin, W., Bernd, O., Annegret, S. "Ultrasonic Cleavage of Thioethers" *J. Phys. Chem. A*, **2005**, 109, 3762-3766.
- (32) Akhtar, Y. "Interaction studies of L-proline and L-glutamine with some electrolytes in aqueous media." *J. Ind. Council Chem.*, **2004**, 21, 43-47.
- (33) Agrawal, P. B., Narwade, M. L. "Prediction of viscosity and ultrasonic behavior of substituted flavone, isoxazole and pyrazole in 70% acetone water mixture." *Acta Clen. Ind., Chem.*, **2002**, 28, 163-166.
- (34) Shashikant, I. A., Rajput, P. R., Narwade, M. L. "Studies on acoustic properties of some substituted pyrazole, isoxazole and pyrazoline in dioxane at 303 K." *Ind. J. Chem., Sec. A*, **2005**, 44, 2495-2497.
- (35) Ravichandran, S., Ramanathan, K. "Ultrasonic study and allied properties of cholesterol in chloroform solutions at 294 K." *J. Pure and Appl. Ultra.*, **2006**, 28, 40-45.
- (36) Abhishek, K., Thirugnanasambandam, S., Vijayanand, S. M. "Physical Mechanism of Ultrasound-Assisted Synthesis of Biodiesel" *Ind. Eng. Chem. Res.*, **2009**, 48, 534-544.
- (37) Mehrotra, K. N., Rawat, M. K. "Ultrasonic and viscometric studies of manganese dicaprate in organic solvents." *Ind. J. Pure Appl. Phy.*, **1991**, 29, 131-133.
- (38) Warminska, D. S., Anna, P., Hanna, K., Waclaw, G. "Adiabatic Compressibilities of Divalent Transition-Metal Perchlorates and Chlorides in *N,N*-Dimethylacetamide and Dimethylsulfoxide" *J. Chem. Eng. Data*, **2009**, 54, 745-751.
- (39) Labowski, M. A. "Acoustic study of critical nitrobenzene-n-heptane mixture throughout a wide range of frequencies." *Acou. Lett.*, **1979**, 3, 37-42.
- (40) Pandharinath, S. N., Mehdi H. "Ultrasonic velocity and adiabatic compressibility of monochloroacetic acid in aqueous ethanol at various temperatures" *J. Chem. Eng. Data*, **1988**, 33, 165-169.
- (41) Hagen, R., Behrends, R., Kaatze, U. "Acoustical Properties of Aqueous Solutions of Urea:Reference Data for the Ultrasonic Spectrometry of Liquids" *J. Chem. Eng. Data*, **2004**, 49, 988-991.
- (42) Nithya, R., Nithiyanantham, S., Mullainathan, S., Rajasekaran, M. "Ultrasonic Investigation of Molecular Interactions in Binary Mixtures at 303 K" *E-J. Chem.* **2009**, 6, 138-140.

- (43) Daridon, J. L., Lagourette, B., Lagrabette, A. "Acoustic determination of thermodynamic properties of ternary liquid mixtures up to 150 MPa." *Phy. Chem. Liq.*, **1999**, 37, 137-160.
- (44) Ali, A., Tiwari, K., Nain, A. K., Chakravorty, V. "Ultrasonic study of molecular interactions in ternary mixtures of dimethyl sulphoxide (1) + carbon tetrachloride (2) + aromatic hydrocarbons (3) at 308.15 K." *Ind. J. Phy., B*, **2000**, 74, 351-355.
- (45) Vijayalakshmi, K., Lalitha, V., Gandhimathi, R. "Acoustic and excess parameter studies on interaction in ternary liquid mixtures." *J. Pure Appl. Ultra.*, **2006**, 28, 154-156.
- (46) Shipra, B., Falguni, K. "A comparative study of ultrasonic velocity and allied parameters of ternary systems". *J. Pure Appl. Ultras.*, **2000**, 22, 82-89.
- (47) Awasthi, A., Rastogi, M., Gupta, M., Shukla, J. P. "Ultrasonic investigation of molecular association in ternary mixtures." *J. Mol. Liq.*, **1999**, 80, 77-88.
- (48) Oswal, S. L., Patel, N. B. "Speed of Sound, Isentropic Compressibility, Viscosity, and Excess Volume of Binary Mixtures. 2. Alkanenitriles + Dimethylformamide, + Dimethylacetamide, and + Dimethylsulfoxide" *J. Chem. Eng. Data*, **1995**, 40, 845-849.
- (49) Oswal, S. L., Patel, N. B. "Speed of Sound, Isentropic Compressibility, Viscosity, and Excess Volume of Binary Mixtures. 1. Alkanenitriles with Alkyl Acetates." *J. Chem. Eng. Data*, **1995**, 40, 840-844.
- (50) Sivaraman, A., Tiwari, M. K., Kartha, P. E. S., Chandrasekhar, P. "High pressure ultrasonic velocity measurements in critical binary liquid system cyclohexane + acetic anhydride." *Proce. Nuclear Phy. Solid State Phy. Symp.* **1976**, 19, 494-497.
- (51) Murali, K. P., Ranjith, K. B., Sathyanarayana, B., Satyanarayana, N. "Densities, Viscosities, and Speeds of Sound of Binary Liquid Mixtures of Sulfolane with Ethyl Acetate, *n*-Propyl Acetate, and *n*-Butyl Acetate at Temperature of (303.15, 308.15, and 313.15) K." *J. Chem. Eng. Data*, **2009**, 54, 1069-1072.
- (52) Nii, S., Matsuura, K., Fukazu, T., Toki, M., Kawaizumi, F. "A novel method to separate organic compounds through ultrasonic atomization." *Chem. Eng. Res. Des.*, **2006**, 84, 412-415.
- (53) Shipra B., Shah, A. R., Parsania, P. H. "Ultrasonic velocity studies of Dioxane solutions of 1,1-Bis(4-hydroxy phenyl) cyclohexane and 4,4'-dihydroxy diphenyl sulphone at 30 °C." *Asian J. Chem.*, **1996**, 8, 736-738.

- (54) Shipra, B., Rajkotia, K. M., Parsania, P. H. "Studies on acoustical properties of poly(4,4-cyclopentylidene diphenylene toluene-2,4 disulphonate) in different solvents at 30 °C." *Eur. Polym. J.*, **1997**, 33, 1005-1007.
- (55) Shipra, B., Parsania, P. H. "Ultrasonic Studies on acoustical properties of diclofenac in dioxane-water system at 30 °C." *J. Pure Appl. Ultra.*, **1997**, 19, 36-39.
- (56) Shipra, B. "Studies of acoustical properties of methyl salicylate in different solvents." *Asian J. Chem.*, **2001**, 13, 71-75.
- (57) Shipra, B., Swati, O. "Ultrasonic Studies of Some Derivatives of Sulphonamide in Dimethyl formamide." *Fluid Phase Equi.*, **2002**, 200, 11-18.
- (58) Godvani, N, Movaliya, J., Baluja, S. "Acoustical studies of some derivatives of 1,5-benzodiazepines at 298.15 K." *Russ. J. Phys. Chem.*, **2009**, 83(13), 2223-2229.
- (59) Riddick, J. A., Bunger, W. B., Sakano, T. "Organic Solvents: Physical Properties and methods of purification." Fourth Edition., Techniques of Chemistry, **II**, A Wiley-Interscience Publication, John Wiley, New York **1986**.
- (60) Ubbelohde, L. "Theory of the Friction of Lubricated Machine Parts." *Petroleum.*, **1912**, 7, 882-889.
- (61) Robinson, R. A., Stokes, R. H. "Electrolyte Solutions", *Butterworths Scientific Publications* (London), **1955**.
- (62) Sastry, G. L., Sastry, V. K. S., Krishnamurty, B. "Ultrasonic parameters in mixed salt solutions." *Ind. J. Pure Appl. Phys.*, **1968**, 6, 637-638.
- (63) Jacobson, B. *Nature* (London), **1954**, 173, 772.
- (64) Wada, Y. "The relation between compressibility and molal volume of organic liquids." *J. Phy. Soci. Jap.*, **1949**, 4, 280-283.
- (65) Sendhilnathan, S., Bagchi, S., Nema, S. K., Singh, R. P. "Ultrasonic and rheological investigations of solid propellant binders." *Euro. Poly. J.*, **1989**, 25, 441-444.
- (66) Vigoureux, P. "Ultrasonics", *Chapman and Hall*, London **1952**.
- (67) Johri, G. K., Misra, R. C. "Phase-transition study and ultrasonic investigations in tert-butyl alcohol (TBA)." *Ind. J. Phys., A.*, **1985**, 59A(5), 482-8.
- (68) Passynsky, A. *Acta. Phy. Chem.*, **1943**, 22, 317.
- (69) Surayanarayana, C. V. and Kuppasami, K. *J. Acoust. Soc. Ind.*, **1976**, 4, 75.

- (70) Prakash, S., Icinaporia, F. M., Pandey. J. D. "Structural Study of the Complex Barium Citrate by Ultrasonic Waves." *J. phys. Chem.*, **1964**, 68, 3078-3080.
- (71) Bachem, C. Z. *Electrochem.*, **1935**, 41, 570.
- (72) Gucker, F. T. *Chem. Rev.*, **1993**, 64, 111.
- (73) Masson, D. O. "Solute molecular volumes in relation to solvation and ionization." *Philos. Mag.*, **1929**, 8, 218-235.



Section-II

Density and Refractive Index

INTRODUCTION

The refractive index of a substance is defined as the ratio of the velocity of light in vacuum to its velocity in the given medium. The refractive index of a material is the most important property of any optical system that uses refraction.

The refractive index plays a vital role in many branches of physics, biology and chemistry. Knowledge of the refractive index of aqueous solutions of salts and biological agents is of crucial importance in applications of evanescent wave techniques in biochemistry¹. Chemical modifications may also be detected by measurements of refractive index². Among the many possible applications is the control of adulteration of liquids³.

It is used to calculate the focusing power of lenses, and the dispersive power of prisms. Since, refractive index is a fundamental physical property of a substance, it is often used to identify a particular substance, confirm its purity, or measure its concentration. For a solution of sugar, the refractive index can be used to determine the sugar content. A refractometer is the instrument used to measure refractive index.

Different methods have been developed to measure the refractive index of liquids. The most common type of refractometer measures the refractive index of the samples by detecting the critical angle of total reflection⁴. Many kinds of interferometric methods for determining the refractive index of materials have also been developed^{5, 6}.

Many workers have reported refractive index of oils^{7, 8}, amino acid⁹, protein¹⁰, sugar^{11, 12}, liquid crystals^{13, 14}, various gases¹⁵, other materials^{16, 17} etc. Further, much work has been done in liquid mixtures¹⁸⁻³⁰ but scanty work has been reported for the solutions of organic³¹⁻³³, inorganic³⁴⁻³⁶ and polymeric materials³⁷⁻³⁹.

Thus, in the present section, the refractive index of solutions of dihydropyrimidinones has been measured in dimethylformamide and dimethyl sulphoxide at 298.15 K. From the experimental data, the density and refractive index of the compounds have been evaluated.

EXPERIMENTAL

The solvents DMF and DMSO were purified by fractionally distillation by the reported method³⁹. For each compound, a series of solutions of different concentrations were prepared in these solvents.

The density and refractive index of pure solvents and solutions were measured by using pycnometer and Abbe refractometer respectively at 298.15 K. The temperature was maintained by circulating water through jacket around the prisms of refractometer from an electronically controlled thermostatic water bath (NOVA NV-8550 E). The uncertainty of temperature was ± 0.1 °C and that of density and refractive index was ± 0.0001 g/cm³ and 0.0005 respectively.

RESULTS AND DISCUSSION

The density of solution (ρ_{12}) is related to densities of the solvent, solute and their weight fractions g_1 and g_2 according to the equation:

$$\frac{1}{\rho_{12}} = \frac{g_1}{\rho_1} + \frac{g_2}{\rho_2} \quad \dots (3.2.1)$$

where ρ_{12} is the density of solution and ρ_1 and ρ_2 are the densities of solvent and solute respectively. Tables 3.2.1 and 3.2.2 show the experimental values of densities and refractive index for all the studied solutions.

The density of these compounds was determined from the slope of the plot of $1/g_1\rho_{12}$ verses g_2/g_1 . Figure 3.2.1 shows the plot of $1/g_1\rho_{12}$ verses g_2/g_1 for RVG-1 in DMF and DMSO respectively. The inverse of slope gives the value of ρ_2 . The densities of all the compounds (ρ_2) evaluated from such plots are given in Table 3.2.3 in DMF and DMSO solutions. Further, the density of compounds were evaluated by using the following equation (3.2.2),

$$\rho = KM/N_A \sum \Delta V_i \quad \dots (3.2.2)$$

where ρ is the density of the compound, K is packing fraction (0.599), M is the molecular weight of the compound, N_A is the Avogadro's number and ΔV_i is the volume increment of the atoms and atomic groups present in the compound. The densities of all the studied compounds have been evaluated and are reported in Table 3.2.3. The calculated volume increment ΔV_i for different atomic groups are given in Table 3.2.4.

Comparison of densities evaluated from graphs and those calculated from eq. (3.2.2) showed that calculated values are different from those evaluated graphically. Further, for the same compound, density in the two solvents is different. This suggests that solvent plays an important role. In solutions, molecular interactions exist which differ in different solvents. These interactions differ due to different substitutions in compounds. The presence of these interactions has also observed in ultrasonic studies which are discussed in section I of chapter 2. Due to these interactions, there may be some changes in volume which affects density.

Table 3.2.1: The density (ρ_{12}) and refractive index (n) of Dihydro pyrimidinones in DMF at 298.15K.

| Conc. <i>M</i> | ρ_{12} <i>g.cm⁻³</i> | <i>n</i> | ρ_{12} <i>g.cm⁻³</i> | <i>n</i> |
|---------------------------|--|-----------------|--|-----------------|
| | RVG-1 | | RVG-6 | |
| 0.00 | 0.9453 | 1.4218 | 0.9453 | 1.4218 |
| 0.01 | 0.9494 | 1.4261 | 0.9487 | 1.4219 |
| 0.02 | 0.9506 | 1.4274 | 0.9505 | 1.4227 |
| 0.04 | 0.9531 | 1.4291 | 0.9518 | 1.4233 |
| 0.06 | 0.9555 | 1.4308 | 0.9547 | 1.4249 |
| 0.08 | 0.9577 | 1.4321 | 0.9554 | 1.4259 |
| 0.10 | 0.9584 | 1.4336 | 0.9566 | 1.4273 |
| | RVG-2 | | RVG-7 | |
| 0.01 | 0.9485 | 1.4237 | 0.9485 | 1.4222 |
| 0.02 | 0.9498 | 1.4248 | 0.9500 | 1.4229 |
| 0.04 | 0.9516 | 1.4257 | 0.9515 | 1.4236 |
| 0.06 | 0.9536 | 1.4275 | 0.9528 | 1.4241 |
| 0.08 | 0.9554 | 1.4283 | 0.9535 | 1.4256 |
| 0.10 | 0.9571 | 1.4308 | 0.9546 | 1.4269 |
| | RVG-3 | | RVG-8 | |
| 0.01 | 0.9488 | 1.4223 | 0.9505 | 1.4219 |
| 0.02 | 0.9490 | 1.4231 | 0.9523 | 1.4224 |
| 0.04 | 0.9520 | 1.4235 | 0.9558 | 1.4230 |
| 0.06 | 0.9527 | 1.4241 | 0.9581 | 1.4236 |
| 0.08 | 0.9543 | 1.4249 | 0.9587 | 1.4241 |
| 0.10 | 0.9565 | 1.4265 | 0.9595 | 1.4249 |
| | RVG -4 | | RVG-9 | |
| 0.01 | 0.9470 | 1.4219 | 0.9508 | 1.4224 |
| 0.02 | 0.9474 | 1.4226 | 0.9524 | 1.4232 |
| 0.04 | 0.9478 | 1.4231 | 0.9558 | 1.4239 |
| 0.06 | 0.9484 | 1.4238 | 0.9576 | 1.4252 |
| 0.08 | 0.9489 | 1.4249 | 0.9605 | 1.4268 |
| 0.10 | 0.9505 | 1.4255 | 0.9617 | 1.4281 |
| | RVG -5 | | RVG-10 | |
| 0.01 | 0.9223 | 1.4225 | 0.9514 | 1.4221 |
| 0.02 | 0.9238 | 1.4234 | 0.9536 | 1.4227 |
| 0.04 | 0.9248 | 1.4243 | 0.9558 | 1.4234 |
| 0.06 | 0.9268 | 1.4253 | 0.9578 | 1.4241 |
| 0.08 | 0.9285 | 1.4258 | 0.9590 | 1.4248 |
| 0.10 | 0.9305 | 1.4283 | 0.9602 | 1.4256 |

Table 3.2.2: The density (ρ_{12}) and refractive index (n) of Dihydropyrimidinones in DMSO at 298.15K.

| Conc. <i>M</i> | ρ_{12} <i>g.cm⁻³</i> | <i>n</i> | ρ_{12} <i>g.cm⁻³</i> | <i>n</i> |
|---------------------------|--|-----------------|--|-----------------|
| | RVG-1 | | RVG-6 | |
| 0.00 | 1.0959 | 1.4716 | 1.0959 | 1.4716 |
| 0.01 | 1.0971 | 1.4719 | 1.0972 | 1.4732 |
| 0.02 | 1.0982 | 1.4729 | 1.0982 | 1.4741 |
| 0.04 | 1.0989 | 1.4733 | 1.0995 | 1.4747 |
| 0.06 | 1.0998 | 1.4739 | 1.1007 | 1.4753 |
| 0.08 | 1.1004 | 1.4745 | 1.1011 | 1.4764 |
| 0.10 | 1.1012 | 1.4779 | 1.1028 | 1.4781 |
| | RVG-2 | | RVG-7 | |
| 0.01 | 1.0971 | 1.4722 | 1.0978 | 1.4726 |
| 0.02 | 1.0987 | 1.4729 | 1.0984 | 1.4734 |
| 0.04 | 1.0992 | 1.4732 | 1.0998 | 1.4743 |
| 0.06 | 1.1005 | 1.4739 | 1.1004 | 1.4751 |
| 0.08 | 1.1017 | 1.4746 | 1.1011 | 1.4764 |
| 0.10 | 1.1023 | 1.4759 | 1.1028 | 1.4779 |
| | RVG-3 | | RVG-8 | |
| 0.01 | 1.0965 | 1.4728 | 1.0986 | 1.4726 |
| 0.02 | 1.0973 | 1.4733 | 1.0995 | 1.4735 |
| 0.04 | 1.1002 | 1.4741 | 1.1003 | 1.4743 |
| 0.06 | 1.1009 | 1.4749 | 1.1011 | 1.4758 |
| 0.08 | 1.1010 | 1.4762 | 1.1030 | 1.4771 |
| 0.10 | 1.1026 | 1.4779 | 1.1044 | 1.4784 |
| | RVG -4 | | RVG-9 | |
| 0.01 | 1.0974 | 1.4728 | 1.0982 | 1.4719 |
| 0.02 | 1.0982 | 1.4735 | 1.0986 | 1.4725 |
| 0.04 | 1.0992 | 1.4748 | 1.0998 | 1.4731 |
| 0.06 | 1.0996 | 1.4762 | 1.1006 | 1.4739 |
| 0.08 | 1.1010 | 1.4778 | 1.1013 | 1.4746 |
| 0.10 | 1.1027 | 1.4783 | 1.1021 | 1.4756 |
| | RVG -5 | | RVG-10 | |
| 0.01 | 1.0969 | 1.4719 | 1.0970 | 1.4719 |
| 0.02 | 1.0985 | 1.4723 | 1.0983 | 1.4725 |
| 0.04 | 1.0996 | 1.4729 | 1.0991 | 1.4731 |
| 0.06 | 1.1003 | 1.4741 | 1.0999 | 1.4739 |
| 0.08 | 1.1017 | 1.4749 | 1.1007 | 1.4746 |
| 0.10 | 1.1036 | 1.4756 | 1.1013 | 1.4756 |

Figure 3.2.1: The variation of $1/g_1\rho_{12}$ with g_2/g_1 for RVG-1 in [A] DMF and [B] DMSO at 298.15 K.









Table 3.2.3: Experimental and calculated densities of Dihydropyrimidinones in DMF and DMSO Solutions at 298.15 K.

| <i>Compounds</i> | Density (g.cm^{-3}) calculated from Figure 3.2.1 | | Density (g.cm^{-3}) Calculated from Eq ⁿ . 3.2.2 |
|------------------|---|-------------|--|
| | <i>DMF</i> | <i>DMSO</i> | |
| <i>RVG-1</i> | 1.3746 | 1.2347 | 1.0526 |
| <i>RVG-2</i> | 1.3226 | 1.3201 | 1.0749 |
| <i>RVG-3</i> | 1.2560 | 1.3168 | 1.1127 |
| <i>RVG-4</i> | 1.0622 | 1.3006 | 1.0749 |
| <i>RVG-5</i> | 1.1847 | 1.3355 | 1.0526 |
| <i>RVG-6</i> | 1.2962 | 1.3029 | 1.1319 |
| <i>RVG-7</i> | 1.2596 | 1.2999 | 1.1127 |
| <i>RVG-8</i> | 1.3454 | 1.3186 | 1.1588 |
| <i>RVG-9</i> | 1.3263 | 1.2204 | 1.1588 |
| <i>RVG-10</i> | 1.3939 | 1.1790 | 1.1695 |

Table 3.2.4: Volume increments of some atoms and groups of atoms.

| Atoms or Atomic group | Volume Increments (A°) ³ | Atoms or Atomic group | Volume Increments (A°) ³ |
|-----------------------|-------------------------------------|-----------------------|-------------------------------------|
| | 10.2 | | 19.35 |
| | 7.84 | | 9.2 |
| | 9.0 | | 2.89 |
| | 3.61 | | 7.46 |
| | 11.36 | | 0.9 |
| | 14.10 | | 14.7 |
| | 10.47 | | 5.093 |
| | 11.40 | | 5.62 |
| | 10.39 | | 2.67 |

| Atoms or Atomic group | Volume Increments (A°) ³ | Atoms or Atomic group | Volume Increments (A°) ³ |
|---|-------------------------------------|--|-------------------------------------|
|  | 23.5 |  | 26.3 |
|  | 11.65 |  | 8.3 |

Further, the molar refraction of a pure liquid $(MRD)_1$ were calculated by the following equation:

$$(MRD)_1 = \left[\frac{n^2 - 1}{n^2 + 1} \right] \frac{M}{\rho} \quad \dots (3.2.3)$$

where n , M and ρ are refractive index, molecular weight and density of pure liquid respectively.

For solutions, following equation was used to determine molar refraction.

$$(MRD)_{12} = \left[\frac{n_{12}^2 - 1}{n_{12}^2 + 1} \right] \left[\frac{X_1 M_1 + X_2 M_2}{\rho_{12}} \right] \quad \dots (3.2.4)$$

where n_{12} and ρ_{12} are refractive index and density of solution respectively. X_1 and X_2 are the mole fractions and M_1 and M_2 are the molecular weight of the solvent and solute respectively.

The plots of $(MRD)_{12}$ verses concentration for dihydropyrimidines series in DMF and DMSO are given in Figures 3.2.2 and 3.2.3 respectively. It is evident from these figures that $(MRD)_{12}$ increase with the increase in concentration. From the values of the molar refraction of solution and pure solvent, molar refraction of solid compounds were determined by following equation:

$$(MRD)_{12} = X_1 (MRD)_1 + X_2 (MRD)_2 \quad \dots (3.2.5)$$

From the density and molar refraction data, the refractive indexes of all the compounds were calculated from eq. (3.2.3). The molar refraction $(MRD)_2$ and refractive index of all the compounds are reported in Table 3.2.5 for 0.1 M solution.

It is evident from Table 3.2.5 that both $(MRD)_2$ and refractive index of compounds are different in each solvent. This again confirms different inter molecular interactions in different solvents. In some solvents, aggregation or hydrogen bonding takes place whereas in others, breakage of bonds takes place.

Figure 3.2.2: The plots of molar refraction $(MRD)_{12}$ against concentration of Dihydropyrimidinones in DMF solutions at 298.15 K.





Figure 3.2.3: The plots of molar refraction $(MRD)_{12}$ against concentration of Dihydropyrimidinones in DMSO solutions at 298.15 K.





Table 3.2.5: Molar refraction (MRD_2) and refractive index (n) of 0.1M solution of Dihydropyrimidinones in DMF and DMSO at 298.15 K.

| Compounds | Solvents | | | |
|---------------|-------------|--------|-------------|--------|
| | DMF | | DMSO | |
| | (MRD_2) | n | (MRD_2) | n |
| RVG-1 | 133.61 | 1.4336 | 129.88 | 1.4779 |
| RVG-2 | 118.08 | 1.4308 | 102.81 | 1.4759 |
| RVG-3 | 102.37 | 1.4265 | 117.35 | 1.4779 |
| RVG-4 | 107.93 | 1.4255 | 114.02 | 1.4783 |
| RVG-5 | 181.09 | 1.4283 | 102.03 | 1.4756 |
| RVG-6 | 106.33 | 1.4273 | 117.85 | 1.4781 |
| RVG-7 | 105.32 | 1.4269 | 112.75 | 1.4779 |
| RVG-8 | 95.20 | 1.4249 | 124.21 | 1.4784 |
| RVG-9 | 106.18 | 1.4281 | 131.64 | 1.4787 |
| RVG-10 | 92.44 | 1.4256 | 113.55 | 1.4756 |

Further, different compounds have different substitutions with the same central moiety. So, $(MRD)_2$ and refractive index values in Table 3.2.5 suggest that substitutions affect molar refractions to a larger extent than the refractive index, which changes only slightly.

LITERATURE CITED

- (1) Zocchi, G. "Mechanical measurement of the unfolding of a protein." *Europhys. Let.*, **1996**, 35, 633-638.
- (2) Barbara, B., Fatme, M., Susan, E. Y., Dennis, T. C., Xavier, L., Michael, J. C. "Effect of physico-chemical modification on the immunogenicity of Haemophilus influenzae type b oligosaccharide-CRM197 conjugate vaccines." *Vaccine*, **2001**, 19, 3189-3200.
- (3) Angelis, De, M., Nicola, De, S., Ferraro, P., Finizio, A., Pierattini, G. "Liquid refractometer based on interferometric fringe projection." *Optics Comm.*, **2000**, 175, 315-321.
- (3) Chandra, B. P., Bhaiya, C. "A simple, accurate alternative to minimum deviation method of determining the refractive index of liquids." *Ame. J. Phys.*, **1983**, 51, 160-161.
- (4) EI-Kashef, H., Hassan, G. E., EI-Ghazaly, I. "Mach-Zehnder optical system as a sensitive measuring instrument." *App. Opt.*, **1994**, 33, 3540-3544.
- (5) Heavens, O. S., Ditchburn, R. W. "Interferometry in Insight into Optics." (John Wiley & Sons Ltd. Chichester West Sussex UK, **1991**, 193-202.
- (6) Chatwal, G., Anand, S. "Instrumental methods of chemical analysis" *Himalaya publishing house, 2nd edi.*, **1984**, 365.
- (7) Rhodes, F. H., Goldsmith, H. E. "Variation of Refractive Index of China Wood Oil with Temperature." *Ind. Eng. Chem.*, **1923**, 15, 786-786.
- (8) Mitsuhiro, F., Tadashi, Y., Satoshi, M., Yasuhiro, O. "Concentration measurement of refrigerant/refrigeration oil mixture by refractive index." *Int. J. Refrig.*, **2004**, 27, 346-352.
- (9) Ali, A., Khan, S., Hyder, S., Nain, A. K. "Volumetric, viscometric and refractive index study of amino acids in mixed solvents at 308.15 K." *Phys. Chem. Liq.*, **2006**, 44(6), 655-662.
- (10) Husband, F. A., Garrood, M. J., Mackie, A. R., Burnett, G. R., Wilde, P. J. "Adsorbed Protein Secondary and Tertiary Structures by Circular Dichroism and Infrared Spectroscopy with Refractive Index Matched Emulsions" *J. Agric. Food Chem.*, **2001**, 49, 859-866.
- (11) Barroso, M., Pineiro, M., Serra, M. E. Silva, Seixas de M. J. "Refractometry in determination of sugar and Aspartame contents in commercial beverages." *Quimica (Lisboa, Portugal)*, **2003**, 91, 61-65.
- (12) Ko, J. H., Cheong, W. J. "Simultaneous quantitative determination of mono-, di-, tri-, tetra-, and penta-saccharides in yogurt products by a simple HPLC system with a refractive index detector." *Bull. Korean Chem. Soc.*, **2001**, 22, 123-126.

- (13) Murakami, N., Yoshimi, H. "Liquid crystal cell substrate for liquid crystal display device and refractive indices thereof." *Jpn. Kokai Tokkyo Koho.*, **2009**, 26, 31-38.
- (14) Sim, J. S., Baek, G. R., Won, Y. C., Park, Y. S., Park, I. G., Jeon, S. H. "Optically anisotropic films with controlled refractive index in the thickness direction for liquid crystal displays." *Repub. Korean Kongkae Taeho Kongbo.*, **2009**, 6, 26-31.
- (15) Clergent, Y., Durou, C., Laurens, M. "Refractive Index Variations for Argon, Nitrogen, and Carbon Dioxide at $\lambda = 632.8$ nm (He-Ne Laser Light) in the Range 288.15 K, T 323.15 K, $0 < p < 110$ kPa." *J. Chem. Eng. Data.*, **1999**, 44, 197-199.
- (16) Burokur, S. N., Sellier, A., Kante, B., Lustrac, De, A. "Symmetry breaking in metallic cut wire pairs metamaterials for negative refractive index. *Appl. Phys. Lett.*, **2009**, 94(20), 201111-201113.
- (17) Wang, J., Hutchins, M., Woo, K., Matayabas, C., Konish, T. "Halogen-free, radiation-curable, high refractive index materials. *JCT Coatings Tech.*, **2009**, 6, 44-49.
- (18) Aleksandar, Z. T., Bojan, D. D., Dusan, K. G., Radojkovic, N. "Use of mixing rules in predicting refractive indexes and specific refractivities for some binary liquid mixtures." *J. Chem. Eng. Data.*, **1994**, 39, 404-404.
- (19) Chen, S., Lei, Q., Fang, W. "Density and Refractive Index at 298.15 K and Vapor-Liquid Equilibria at 101.3 kPa for Four Binary Systems of Methanol, *n*-Propanol, *n*-Butanol, or Isobutanol with *N*-Methylpiperazine" *J. Chem. Eng. Data*, **2002**, 47, 811-815.
- (20) Gomez-Diaz, D., Mejuto, J. C., Navaza, J. M. "Physicochemical Properties of Liquid Mixtures. 1. Viscosity, Density, Surface Tension and Refractive Index of Cyclohexane + 2,2,4-Trimethylpentane Binary Liquid Systems from 25 C to 50 C." *J. Chem. Eng. Data.*, **2001**, 46, 720-724.
- (21) Baluja, S., Pandaya, N., Kachhadia, N. and Solanki, A. "Refractive index: theoretical evaluation in binary liquid mixtures." *J. Ultra Sci. Phy. Sci.*, **2006**, 18, 247-250.
- (22) Pandey, J. D., Jain, P., Vyas, V. "Speed of sound, viscosity, and refractive index of multicomponent systems: analysis of experimental data from the Bertrand-Acree-Burchfield equation." *Can. J. Chem.*, **1994**, 72, 2486-2492.
- (23) Toti, U. S., Kariduraganavar, M. Y., Aralaguppi, M. I., Aminabhavi, T. M. "Density, Viscosity, Refractive Index, and Speed of Sound of Ternary Systems: Polystyrene in 1,4-Dioxane + Tetrahydrofuran Mixtures at

- (298.15, 303.15, and 308.15) K." *J. Chem. Eng. Data*, **2000**, 45, 920-925.
- (24) Gomez-Diaz, D., Mejuto, J. C., Navaza, J. M., Rodriguez-Alvarez, A. "Effect of Composition and Temperature upon Density, Viscosity, Surface Tension, and Refractive Index of 2,2,4- Trimethylpentane + Cyclohexane + Decane Ternary Liquid Systems." *J. Chem. Eng. Data.*, **2003**, 48, 231-235.
- (25) Proutiere, A., Megnassan, E., Hucteau, H. "Reply to Comment on "Refractive Index and Density Variations in Pure Liquids. A New Theoretical Relation." *J. Phys. Chem.*, **1994**, 98, 4769-4769.
- (26) Resa, J. M., Gonzalez, C. and Juez M. "Density, refractive index and speed of sound for mixtures of ethyl acetate with 2-butanol and 3-methyl-1-butanol. Vapor-liquid equilibrium of the ethyl acetate + 3-methyl-1-butanol system." *Fluid Phase Equi.*, **2004**, 217, 175-180.
- (27) Bruce, L., Merle, R. and Heinz, W. "Change of Index of Refraction with Concentration in Solutions of Water and Normal Butyl Alcohol at 30 °C." *J. Phys. Chem.*, **1941**, 45, 340-343.
- (28) Minc, S. "Refraction index of aliphatic alcohol + formamide mixtures at the solution-air interface determined by the ellipsometric method." *J. Electroana. Chem.*, **1982**, 140, 121-130.
- (29) Segade, L., Jimenez de Llano, J., Dominguez-Perez, M., Cabeza, O., Cabanas, M. and Jimenez, E. "Density, Surface Tension, and Refractive Index of Octane + 1-Alkanol Mixtures at $T = 298.15$ K." *J. Chem. Eng. Data.*, **2003**, 48, 1251-1255.
- (30) Resa, J. M., Gonzalez, C. Diez, E., Concha, R. G. and Iglesias, M. "Mixing properties of isopropyl acetate + aromatic hydrocarbons at 298.15 K: Density, refractive index and isentropic compressibility." *Korean J. Chem. Eng.*, **2009**, 21, 1015-1025.
- (31) Benjamin, J. B., Ernest, G. "Molar refraction as an index of proton transfer: An estimate of the acid strength of p-toluenesulfonic acid." *J. Ame. Chem. Soc.*, **1961**, 83, 2956-2956.
- (32) Liu, J., Nakamura, Y., Shibasaki, Y., Ando, S., Ueda, M. "High Refractive Index Polyimides Derived from 2,7-Bis(4-amino phenylenesulfanyl) thianthrene and Aromatic Dianhydrides." *Macromol.*, **2007**, 40, 4614-4620.
- (33) Tanio, N., Irie, M. "Refractive index of organic photochromic dye-amorphous polymer composites." *Jap. J. Appl. Phy.*, **1994**, 33, 3942-3946.

- (34) Sangwal, K., Kucharczyk, W. "Relationship between density and refractive index of inorganic solids." *J. Phys. D. Appl. Phys.*, **1987**, 20, 522-525.
- (35) Koutnik, V. "Refractive index of inorganic pigments." *Chemicky Prumysl.*, **1964**, 14, 604-606.
- (36) Yakowitz, M. L., Jorgensen, P. S. "Refractive Index of Strontium Nitrate" *Ind. Eng. Chem. Anal.*, **1937**, 9, 204-204.
- (37) Lu, C., Guan, C., Liu, Y., Cheng, Y., Yang, B. "PbS/Polymer Nanocomposite Optical Materials with High Refractive Index." *Chem. Mater.*, **2005**, 17, 2448-2454.
- (38) Mohsen-Nia, M., Modarress, H., Rasa, H. "Measurement and Modeling of Density, Kinematic Viscosity, and Refractive Index for Poly(ethylene Glycol) Aqueous Solution at Different Temperatures." *J. Chem. Eng. Data*, **2005**, 50, 1662-1666.
- (39) Assaid, I., Bosc, D., Hardy, I. "Improvements of the Poly(vinyl cinnamate) Photoresponse in Order to Induce High Refractive Index Variations." *J. Phys. Chem. B.*, **2004**, 108, 2801-2806.



Section-III

Conductance

INTRODUCTION

Conductance is the ease with which current flows through a conductor. In solutions, ions conduct electricity so the conductance of such electrolytic solutions depends on the concentration of the ions and also on the nature of the ions present (through their charges and mobilities). The conductance behavior as a function of concentration is different for strong and weak electrolytes. Both strong and weak electrolytes will be studied at a number of dilute concentrations¹⁻⁵, and the ionization constant for a weak electrolyte can be calculated from the data obtained⁶⁻⁸.

Conductivity measurement has widespread use in industrial applications that involve the detection of contaminants in water and concentration measurements⁹. Conductance measurements have been used for the determination of the equilibrium constants, degree of hydrolysis, dissociation constants and other properties of electrolyte solutions. Since all ions present in the solution contribute to the conductance process, conductance is an additive property of a solution depending on all the ions present. In acid-base titrations, this technique is also useful to determine relative strength of the two weak acids or bases, degree of hydrolysis, basicity of organic acid etc.

Literature survey shows that conductance of many organic, inorganic, polymers, rare earth, amino acid, vitamin etc. have been measured¹⁰⁻²². Further, the conductivity measurements have been used in liquid mixtures such as binary and ternary²³⁻²⁸. Many workers have been reported the conductance of organic synthetic compounds²⁹⁻³⁵. However, the conductivity data was not available for 1, 4 dihydropyrimidines.

Thus, in the present section conductance of all the synthesized 1, 4 dihydropyrimidines was measured in DMF and DMSO solutions at 298.15 K, over a wide range of concentration.

EXPERIMENTAL

The solvents DMF and DMSO were purified by fractionally distillation by the method reported in the literature³⁶.

The solutions of different concentrations were prepared for each compound in DMF and DMSO and the conductance of each solution was measured by using Equip-tronics Conductivity Meter (Model No. 664) having cell constant 0.96 cm^{-1} at 298.15 K. The measured conductance was corrected by subtracting the conductance of pure solvent.

RESULTS AND DISCUSSION

The measured conductance (k) of each solution after correction was used to determine the specific conductance (κ), which is then used for the calculation of equivalent conductance (λ_c).

The equations used for calculating specific conductance (κ) and equivalent conductance (λ_c) are:

$$\kappa = k\theta \quad \dots (3.3.1)$$

$$\lambda_c = 1000 \frac{\kappa}{C} \quad \dots (3.3.2)$$

where θ is the cell constant ($= 0.96 \text{ cm}^{-1}$) and c is the concentration (g.equi./lit.) of solution.

Tables 3.3.1 and 3.3.2 show the equivalent conductance of all the studied compounds in DMF and DMSO solutions at 298.15 K along with measured conductance (k). The variation of conductance with concentration for these compounds in both the solvents is given in Figures 3.3.1 and 3.3.2. For studied compounds, conductivities are observed to be less in DMSO than those in DMF. Further, for all the studied systems, conductance increases with concentration.

The equivalent conductance (λ_c) is plotted against \sqrt{C} for all studied compounds and is shown in Figures 3.3.3 and 3.3.4. In both DMF and DMSO solutions, usually λ_c increases with dilution. But for certain compounds in DMSO solutions, λ_c do not increase continuously but bend downward at low concentrations giving rise to maximum. This typical behavior may be due to interactions within the molecule thereby causing constriction within the molecule or due to association between solute with solvent molecules. Similar behavior was observed by Prasad et al³⁷ while studying conductance of polyions in DMSO solutions. Further, it is evident from these figures that some compounds behave as strong electrolytes whereas most of them exhibited weak electrolytic behavior. The behavior is different in both the solvents.

For the systems where λ_c values increase with dilution, λ_0 values were evaluated by extrapolation. These values are compared with those determined by an alternate procedure using the following equation:

Table 3.3.1: The Conductance (k) and equivalent conductance (λ_c) of Dihydropyrimidines in DMF at 298.15 K.

| Conc. M | $k.10^5$ mho | λ_c mho.cm ² .equi. ⁻¹ | $k.10^5$ mho | λ_c mho.cm ² .equi. ⁻¹ | $k.10^5$ mho | λ_c mho.cm ² .equi. ⁻¹ | $k.10^5$ mho | λ_c mho.cm ² .equi. ⁻¹ | $k.10^5$ mho | λ_c mho.cm ² .equi. ⁻¹ |
|--------------|-----------------|---|-----------------|---|-----------------|---|-----------------|---|-----------------|---|
| | RVG-1 | | RVG-2 | | RVG-3 | | RVG-4 | | RVG-5 | |
| 0.000 | 0.20 | - | 0.20 | - | 0.20 | - | 0.20 | - | 0.20 | -#DIV/0! |
| 0.001 | 0.39 | 1.7376 | 0.32 | 1.0656 | 0.40 | 1.8912 | 0.42 | 2.0448 | 0.30 | 0.9312 |
| 0.002 | 0.46 | 1.2336 | 0.39 | 0.8976 | 0.53 | 1.5844 | 0.64 | 2.1024 | 0.35 | 0.7200 |
| 0.004 | 0.67 | 1.1280 | 0.53 | 0.7824 | 0.84 | 1.5192 | 0.99 | 1.8912 | 0.49 | 0.6744 |
| 0.006 | 0.86 | 1.0432 | 0.68 | 0.7632 | 1.20 | 1.5952 | 1.41 | 1.9296 | 0.59 | 0.6224 |
| 0.008 | 0.99 | 0.9456 | 0.86 | 0.7896 | 1.49 | 1.5432 | 1.82 | 1.9392 | 0.69 | 0.5844 |
| 0.010 | 1.18 | 0.9370 | 1.07 | 0.8314 | 1.84 | 1.5706 | 2.19 | 1.9066 | 0.81 | 0.5856 |
| 0.020 | 2.14 | 0.9293 | 1.67 | 0.7037 | 3.42 | 1.5437 | 5.70 | 2.6381 | 1.15 | 0.4541 |
| 0.040 | 3.83 | 0.8702 | 3.06 | 0.6854 | 5.56 | 1.2854 | 10.20 | 2.3990 | 1.92 | 0.4118 |
| 0.060 | 5.19 | 0.7978 | 4.10 | 0.6234 | 7.23 | 1.1242 | 14.80 | 2.3354 | 2.36 | 0.3450 |
| 0.080 | 6.18 | 0.7171 | 4.95 | 0.5695 | 7.82 | 0.9139 | 18.80 | 2.2315 | 2.75 | 0.3055 |
| 0.100 | 7.03 | 0.6553 | 5.70 | 0.5276 | 8.52 | 0.7983 | 22.60 | 2.1500 | 3.20 | 0.2876 |
| | RVG-6 | | RVG-7 | | RVG-8 | | RVG-9 | | RVG-10 | |
| 0.001 | 0.31 | 1.0176 | 0.42 | 2.0736 | 0.32 | 1.0656 | 0.26 | 0.5280 | 0.55 | 3.2736 |
| 0.002 | 0.41 | 1.0032 | 0.59 | 1.8480 | 0.44 | 1.1424 | 0.30 | 0.4512 | 0.78 | 2.7696 |
| 0.004 | 0.55 | 0.8352 | 0.98 | 1.8672 | 0.62 | 1.0008 | 0.37 | 0.4008 | 1.14 | 2.2344 |
| 0.006 | 0.77 | 0.8976 | 1.34 | 1.8176 | 0.79 | 0.9408 | 0.44 | 0.3808 | 1.48 | 2.0432 |
| 0.008 | 0.98 | 0.9336 | 1.75 | 1.8552 | 0.98 | 0.9336 | 0.52 | 0.3828 | 1.84 | 1.9632 |
| 0.010 | 1.23 | 0.9850 | 2.09 | 1.8106 | 1.13 | 0.8890 | 0.61 | 0.3850 | 2.15 | 1.8722 |
| 0.020 | 2.09 | 0.9053 | 3.63 | 1.6445 | 2.16 | 0.9389 | 0.83 | 0.3010 | 3.21 | 1.4429 |
| 0.040 | 3.51 | 0.7934 | 6.24 | 1.4486 | 3.60 | 0.8150 | 1.12 | 0.2198 | 4.22 | 0.9638 |
| 0.060 | 4.62 | 0.7066 | 9.09 | 1.4218 | 4.74 | 0.7258 | 1.32 | 0.1786 | 5.96 | 0.9210 |
| 0.080 | 5.72 | 0.6619 | 11.13 | 1.3111 | 4.98 | 0.5731 | 1.56 | 0.1627 | 7.31 | 0.8527 |
| 0.100 | 6.41 | 0.5958 | 13.44 | 1.2707 | 5.30 | 0.4892 | 1.77 | 0.1503 | 9.22 | 0.8655 |

Table 3.3.2: The Conductance (k) and equivalent conductance (λ_c) of Dihydropyrimidines in DMSO at 298.15 K.

| Conc. <i>M</i> | $k.10^5$ mho | λ_c mho.cm ² .equi. ⁻¹ | $k.10^5$ mho | λ_c mho.cm ² .equi. ⁻¹ | $k.10^5$ mho | λ_c mho.cm ² .equi. ⁻¹ | $k.10^5$ mho | λ_c mho.cm ² .equi. ⁻¹ | $k.10^5$ mho | λ_c mho.cm ² .equi. ⁻¹ |
|-------------------|-----------------|---|-----------------|---|-----------------|---|-----------------|---|-----------------|---|
| | RVG-1 | | RVG-2 | | RVG-3 | | RVG-4 | | RVG-5 | |
| 0.000 | 0.016 | - | 0.016 | - | 0.016 | - | 0.016 | - | 0.016 | - |
| 0.001 | 0.134 | 1.0848 | 0.124 | 0.9888 | 0.191 | 1.6320 | 0.117 | 0.9216 | 0.291 | 2.5920 |
| 0.002 | 0.237 | 1.0368 | 0.177 | 0.7488 | 0.286 | 1.2720 | 0.197 | 0.8448 | 0.524 | 2.4144 |
| 0.004 | 0.425 | 0.9696 | 0.288 | 0.6408 | 0.497 | 1.1424 | 0.382 | 0.8664 | 0.980 | 2.3016 |
| 0.006 | 0.652 | 1.0096 | 0.375 | 0.5664 | 0.678 | 1.0512 | 0.548 | 0.8432 | 1.350 | 2.1264 |
| 0.008 | 0.871 | 1.0200 | 0.473 | 0.5424 | 0.870 | 1.0188 | 0.760 | 0.8868 | 1.670 | 1.9788 |
| 0.010 | 1.010 | 0.9494 | 0.520 | 0.4790 | 1.070 | 1.0070 | 1.020 | 0.9590 | 2.040 | 1.9382 |
| 0.020 | 1.560 | 0.7387 | 0.770 | 0.3595 | 1.750 | 0.8299 | 1.680 | 0.7963 | 3.570 | 1.7035 |
| 0.040 | 2.700 | 0.6430 | 1.320 | 0.3118 | 3.070 | 0.7318 | 3.020 | 0.7198 | 3.780 | 0.9022 |
| 0.060 | 3.770 | 0.5998 | 1.780 | 0.2814 | 4.080 | 0.6494 | 4.160 | 0.6622 | 4.070 | 0.6478 |
| 0.080 | 5.370 | 0.6419 | 2.260 | 0.2687 | 5.010 | 0.5987 | 5.340 | 0.6383 | 4.530 | 0.5411 |
| 0.100 | 6.330 | 0.6057 | 2.890 | 0.2754 | 5.910 | 0.5653 | 6.810 | 0.6517 | 4.680 | 0.4473 |
| | RVG-6 | | RVG-7 | | RVG-8 | | RVG-9 | | RVG-10 | |
| 0.001 | 0.027 | 0.0525 | 0.023 | 0.0172 | 0.022 | 0.0077 | 0.030 | 0.0740 | 0.023 | 0.0163 |
| 0.002 | 0.038 | 0.0748 | 0.028 | 0.0301 | 0.029 | 0.0335 | 0.045 | 0.1036 | 0.027 | 0.0262 |
| 0.004 | 0.060 | 0.0830 | 0.036 | 0.0316 | 0.039 | 0.0385 | 0.084 | 0.1361 | 0.039 | 0.0378 |
| 0.006 | 0.082 | 0.0867 | 0.045 | 0.0337 | 0.050 | 0.0421 | 0.121 | 0.1436 | 0.051 | 0.0423 |
| 0.008 | 0.104 | 0.0888 | 0.053 | 0.0345 | 0.062 | 0.0441 | 0.154 | 0.1433 | 0.061 | 0.0429 |
| 0.010 | 0.124 | 0.0886 | 0.159 | 0.1187 | 0.078 | 0.0490 | 0.214 | 0.1660 | 0.110 | 0.0765 |
| 0.020 | 0.356 | 0.1441 | 0.243 | 0.0955 | 0.137 | 0.0499 | 0.282 | 0.1122 | 0.192 | 0.0735 |
| 0.040 | 0.567 | 0.1174 | 0.385 | 0.0783 | 0.248 | 0.0488 | 0.371 | 0.0753 | 0.301 | 0.0602 |
| 0.060 | 0.785 | 0.1095 | 0.554 | 0.0764 | 0.361 | 0.0487 | 0.497 | 0.0682 | 0.438 | 0.0598 |
| 0.080 | 0.883 | 0.0927 | 0.718 | 0.0749 | 0.451 | 0.0462 | 0.620 | 0.0644 | 0.562 | 0.0582 |
| 0.100 | 0.996 | 0.0839 | 0.910 | 0.0765 | 0.522 | 0.0431 | 0.801 | 0.0671 | 0.756 | 0.0632 |

Figure 3.3.1: The variation of Conductance with concentration for Dihydropyrimidines in DMF at 298.15 K.





Figure 3.3.2: The variation of Conductance with concentration for Dihydropyrimidines in DMSO at 298.15 K.





Figure 3.3.3: The variation of equivalent conductance with \sqrt{C} for Dihydropyrimidines in DMF at 298.15 K.







Figure 3.3.4: The variation of equivalent conductance with \sqrt{c} for Dihydropyrimidines in DMSO at 298.15 K.



$$k = k_0 + \lambda_0 c + c\phi_{(c)} \quad \dots (3.3.3)$$

where k and k_0 are the electrolytic conductivity of the solutions and solvent respectively. c is the equivalent concentration and the function $\phi_{(c)}$ denotes the effect of interionic interactions. The limiting conductivity can be determined accurately from the slope, dk/dc of plot of k verses c , provided other derivatives (dk_0/dc) and $d[c\phi_{(c)}]/dc$ in differential form of equation (3.3.3) are neglected as compared to λ_0 , which can be determined from differential form of equation (3.3.3) is

$$\frac{dk}{dc} = \frac{dk_0}{dc} + \lambda_0 + \frac{d[c\phi_{(c)}]}{dc} \quad \dots (3.3.4)$$

Table 3.3.3 shows the λ_0 values for all studied compounds along with those determined by extrapolation. For the system where λ_c decreases at low concentrations, λ_0 could not be evaluated. Comparison of λ_0 values in Table 3.3.5 show that for most of the systems, values are in good agreement. However, for some cases, deviations are significant suggesting thereby that equations (3.3.3) and (3.3.4) are not valid for these systems. The reason for the deviation may be due to the fact that in equation (3.3.3), the term for interionic interactions³⁷ i.e., are neglected, which are actually present in the studied solutions, as discussed in section 1 (acoustical properties).

Table 3.3.3: The limiting equivalent conductance (λ_0) of all the compounds in DMF and DMSO at 298.15 K.

| Compound Code | λ_0 $\text{mho.cm}^2.\text{equi.}^{-1}$ from graph | $\lambda_0 10^3$ $\text{mho.cm}^2.\text{equi.}^{-1}$ from eq. (3.3.4) | λ_0 $\text{mho.cm}^2.\text{equi.}^{-1}$ from graph | $\lambda_0 10^3$ $\text{mho.cm}^2.\text{equi.}^{-1}$ from eq. (3.3.4) |
|------------------|--|---|--|---|
| | DMF | | DMSO | |
| RVG -1 | 1.10 | 0.94 | 1.10 | 1.01 |
| RVG -2 | 1.14 | 0.83 | 0.83 | 0.49 |
| RVG -3 | 2.01 | 1.59 | 1.10 | 1.01 |
| RVG -4 | - | 1.98 | 0.86 | 0.97 |
| RVG -5 | 0.80 | 0.59 | 2.56 | 2.1 |
| RVG -6 | 1.13 | 1.00 | - | 0.108 |
| RVG -7 | 1.95 | 1.89 | - | 0.045 |
| RVG -8 | 1.12 | 0.92 | - | 0.060 |
| RVG -9 | 0.68 | 0.39 | - | 0.193 |
| RVG -10 | 2.42 | 2.01 | - | 0.056 |

LITERATURE CITED

- (1) Barthel, J., Neueder, G. R., Schmid, A. "Electrolyte solutions for technology-new aspects and approaches." *Pure Appl. Chem.*, **1999**, 71, 1705-1715.
- (2) Barthel, J., Neueder, R., Feuerlein, F., Iberl, L. "Conductance of electrolytes in ethanol solutions from -45 to 25 °C." *J. Sol. Chem.*, **1983**, 12, 449-471.
- (3) Barthel, J., Graml, H., Neueder, R., Turq, P., Bernard, O. "Electrolyte conductivity from infinite dilution to saturation current topics in solution chemistry." *Res. Trends.*, **1994**, 2, 106-112.
- (4) Falkenhagen, H., Williams, J. W. "The frequency dependence of the electrical conductance of solutions of strong electrolytes." *J. Phy. Chem.*, **1929**, 33, 1121-1134.
- (5) Wang, K., Sun, X. "A method for calculating thermal conductivity of electrolytic non-aqueous solutions." *Huaxue Gongcheng (Xi'an, China)*. **2002**, 30, 62-64.
- (6) Linde, W. V., Northcott, D., Redmond, W., Robertson, R. E. "Basic dissociation constant for ethylamine by a convenient conductance method." *Can. J. Chem.*, **1969**, 47, 279-285.
- (7) Goffredi, M., Shedlovsky, T. "Studies of Electrolytic Conductance in Alcohol-Water Mixtures. The Ionization Constant of Acetic Acid in 1-Propanol - Water Mixtures at 15, 25, and 35 °C." *J. Phy. Chem.* **1967**, 71, 4436-4442.
- (8) Siow, K. S., Ang, K. P. "Thermodynamics of ionization of 2,4- and 2,6-dinitrophenol in dimethylsulfoxide" *Can. J. Chem.* **1980**, 58, 970-973.
- (9) Levitt, L. S., "Determination of the dissociation constant and limiting equivalent conductance of a weak electrolyte from conductance measurements on the weak electrolyte." *Ana. Chim. Acta.*, **1981**, 125, 219-220.
- (10) Latini, G., Grifoni, R. C., Passerini, G., Tascini, S. "An enhanced prediction method for thermal conductivity of organic compounds and their mixtures in the saturated liquid state." *Thermal Conduc.* **2005**, 26, 479-488.
- (11) Mayahi, M. F., Hussain, H. A. "Solvent effects on the conductivity of organic compounds." *J. Ind. Chem. Soc.*, **1973**, 50, 34-36.

- (12) Carlton, D. M., McCarthy, D. K., Genz, R. H. "The effect of structure on the electrical conductivity of organic compounds. Polyazophenylenes." *J. Phy. Chem.*, **1964**, 68, 2661-2665.
- (13) George, M., Ritzoulis, G. "Electrolytic Conductances of Tetrabutyl ammonium Tetraphenylborate in Propylene Carbonate + Acetonitrile Mixtures at (25, 30, and 35) °C" *J. Chem. Eng. Data*, **1996**, 41, 326-330.
- (14) Al-Najar, A. A., Khadim, M. H. A., Ebraheem, A. M. "Conductance of Lithium and Sodium Perchlorates and of Some Quaternary Ammonium Halides and Perchlorates in a Propylene Carbonate + Tetrahydrofuran Mixture at 25 °C" *J. Chem. Eng. Data*, **1992**, 37, 443-445.
- (15) Teo, F. A. G., Hector, R. G., Carol, C. S., Maria, E. T. "Density, Refractive Index, Viscosity, and Electrical Conductivity in the Na₂CO₃ + Poly(ethylene glycol) + H₂O System from (293.15 to 308.15) K" *J. Chem. Eng. Data*, **2004**, 49, 1254-1257.
- (16) Spedding, F. H., Joseph, A. R., Victor, W. S. "Electrical Conductances of Some Aqueous Rare Earth Electrolyte Solutions at 25 °C. II. Rare Earth Chlorides." *J. Chem. Eng. Data*, **1974**, 19, 373-378.
- (17) Srivastava, A. K., Shankar, S. L. "Ionic Conductivity in Binary Solvent Mixtures. 4. Dimethylsulfoxide + Water at 25 °C." *J. Chem. Eng. Data*, **2000**, 45, 92-96.
- (18) Rewick, R. T., Dolder, M. P. "Conductometric Analysis of Oleum" *Anal. Chem.*, **1972**, 44, 1096-1097.
- (19) Dev, S., Ismail, K. "Electrical Conductance of Sodium Dodecyl Sulfate in Calcium Nitrate Tetrahydrate + Acetamide Melts" *J. Chem. Eng. Data*, **2001**, 46, 574-576.
- (20) Rogac, B. M., Neueder, R., Barthel, J., Apelblat, A. "Conductivity studies on aqueous solutions of stereoisomers of Tartric acid and Tatrates- Part 3. Acidic Tatrates." *J. Sol. Chem.*, **1998**, 27, 299-307.
- (21) Nandi, D., Hazra, D. K., "Conductances of alkali metal chlorides and bromides in aqueous binary mixture of 2-methoxyethanol at 298.15 K." *J. Ind. Chem. Soc.*, **2003**, 80, 21-29.
- (22) Werner, M., Baars, A., Werner, F., Eder, C., Delgado, A. "Thermal conductivity of aqueous sugar solutions under high pressure." *Int. J. Thermo.*, **2007**, 28, 1427.
- (23) Aloys, J. D., Popovych, O. "Electrolytic Conductance and Ionic Association of Several Salts in Ethanol-Water Mixtures." *J. Chem. Eng. Data*, **1969**, 14, 156-160.

- (24) Champeney, D. C., Kaddour, F. O. "Conductance of KCl-Glycerol Solutions: Test of the Fuoss Paired Ion Model." *Phys. Chem. Liq.*, **1987**, 16, 239-247.
- (25) Yecid, P. J., Taboada, M. E., Flores, E. K., Galleguillos, H. R. "Density, Viscosity, and Electrical Conductivity in the Potassium Sulfate + Water + 1-Propanol System at Different Temperatures." *J. Chem. Eng. Data*, **2009**, 54, 1932-1934.
- (26) Wu, B., Zhang, Y. M., Wang, H. P. "Volumetric Properties and Conductivities of 1-Butyl-3-methylimidazolium Tetrafluoroborate + Sucrose + Water Mixtures." *J. Chem. Eng. Data*, **2009**, 54, 1430-1434.
- (27) Niazi, M. S. K. "Conductivities and Thermodynamic Dissociation Constants for Chloroacetic Acid in Binary Mixed Solvent Systems at 298.15 K." *J. Chem. Eng. Data*, **1993**, 38, 527-530.
- (28) Radhika, V., Manikyamba, P. "Conductance and Solvation Behavior of Quinolinium Dichromate in Binary Mixtures of Water with N,N-Dimethyl Formamide." *J. Chem. Eng. Data*, **2008**, 53, 2766-2769.
- (29) Baluja, S. "Physicochemical studies of some Schiff bases derived from 6-ethylbenzene-1,3-diol." *E- J. Chem.*, **2004**, 1, 199-205.
- (30) Kabir-ud-Din, A. A., Mohammed, D. A., Naqvi, A. Z., Akram, M. "Conductometric study of antidepressant drug-cationic surfactant mixed micelles in aqueous solution, Colloids and Surfaces." *Biointerfaces*. **2008**, 64, 65-69.
- (31) Parvatalu, P., Srivastava. A. K. "Ionic Conductivity in Binary Solvent Mixtures. 6. Behavior of Selected 1:1 Electrolytes in 80 mass % Propylene Carbonate + p-Xylene at 25 °C." *J. Chem. Eng. Data*, **2003**, 48, 608-611.
- (32) Campbell, T. G., Urbach, F. L. "Electronic, circular dichroism, and proton magnetic resonance spectral studies of the nickel(II) complexes of some neutral, tetradentate Schiff base ligands derived from 1,3-diamines." *Inorg. Chem.*, **1973**, 12, 1840-1846.
- (33) Liu, J., Masuda, Y., Sekido E. "A conductance study of macrocyclic Schiff base metal(II) complexes in methanol." *Bull. Chem. Soc. Jap.*, **1990**, 63, 2516-2520.
- (34) Carlen, P. L., Gurevich, N., Polc, P. "Low-dose benzodiazepine neuronal inhibition: enhanced Ca^{2+} mediated K^{+} conductance." *Brain res.*, **1983**, 271, 358-364.
- (35) Baluja, S. Kasundra, P., Vekariya, N. "Physicochemical studies of some azomethines of 5-amino isophthalic acid in solutions of DMF and DMSO at 308.15 K." *Int. J. Chem. Sci.*, 2009, 7, 533-538.

- (36) Riddick, J. A., Bunger, W. B., Sakano, T. "Organic Solvents-Physical Properties and methods of purification." Fourth Edition. Techniques of Chemistry, II, *A Wiley-Interscience Publication, John Wiley, New York, 1986.*
- (37) Prasad, B. B., Singh, M., Kumar, A., Easo, S. "Electrolytic conductivity of crystal violet based quaternary ammonium polyelectrolytes in N, N-dimethylformamide and dimethyl sulphoxide." *Can. J. Chem.*, **1997**, 75, 414-422.



Section-IV

Solubility

INTRODUCTION

The extensive information on the thermodynamic properties of organic compounds is needed not only their use in many industrial processes but also for the advancement of theoretical developments through an understanding of the intermolecular forces in solution for structure-property relationship. Solubility data is important information in drug discovery, drug formulation and crystallization-based separation investigations.

The solubility of drug is essential information in drug discovery and is important in the preparation of liquid drug formulation stages in the pharmaceutical industry^{1, 2}. There are many methods for solubilization of drugs including cosolvency, surface active agents, salt information, complexation, hydrotropism, crystal engineering and preparation of soluble prodrug³⁻⁵.

The process of dissolving is a process which involves the breaking and making of bonds and that involves energy which is in the form of heat.

Thus, the process of dissolution is accompanied by the heat change i.e., enthalpy change (ΔH_{sol}). If the heat is absorbed, the process is called endothermic. In this case, ΔH_{sol} is positive. If the heat is evolved i.e., process is exothermic and the ΔH_{sol} values will be negative.

This enthalpy change is known as heat of solution, which is the change in heat content when one mole of substance is dissolved in specific quantity of solvent at a given temperature.

The molar heat of solution and melting temperature of a substance can be determined from the solubility measurements⁶. In recent years many researchers have been worked on solubility of some natural compounds^{7, 8}, some organic⁹⁻¹¹, polymers^{12, 13}, amino acids^{14, 15}, drugs¹⁶⁻¹⁹, vitamins²⁰, ionic liquids^{21, 22}, and inorganic compounds²³⁻²⁵ by various methods²⁶⁻³³. In our laboratory, heat of solution of some synthesized heterocyclic compounds and drugs has also been determined³⁴⁻³⁶. From the solubility data, some thermodynamic parameters have also been evaluated.

In the present work, the solubility for some dihydropyrimidinones derivatives was determined in N, N-dimethylformamide (DMF) and dimethylsulfoxide (DMSO) at different temperatures (298.15 to 318.15) K. Further, some thermodynamic parameters such as enthalpy, Gibb's free energy and entropy of different solutions have also been evaluated from the solubility data.

EXPERIMENTAL

The heat of solution of dihydropyrimidines has been studied in DMF and DMSO. These solvents were purified and fractionally distilled prior to use by the method reported in the literature³⁷.

The solubilities were measured by a gravimetric method³⁸. For each measurement, an excess mass of dihydropyrimidinone was added to a known mass of solvent. Then, the equilibrium cell was heated to a constant temperature with continuous stirring. After, at least 3 h (the temperature of the water bath approached constant value, then the actual value of the temperature was recorded), the stirring was stopped and the solution was kept still for 2 h. A portion of this solution was filtered and by a preheated injector, 2 ml of this clear solution was taken in another weighted measuring vial (m_0). The vial was quickly and tightly closed and weighted (m_1) to determine the mass of the sample ($m_1 - m_0$). Then, the vial was covered with a piece of filter paper to prevent dust contamination. Then, the vial was placed in at room temperature to evaporate the solvent. After the solvent in the vial had completely evaporated, the vial was dried and reweighed (m_2) to determine the mass of the constant residue solid ($m_2 - m_0$). All the weights were taken using an electronic balance (Mettler Toledo AB204-S, Switzerland) with an accuracy of ± 0.0001 g. Thus, the solid concentration of the sample solution of mole fraction, x , could be determined from equation 1.

$$x = \frac{(m_2 - m_0) / M_1}{(m_2 - m_0) / M_1 + (m_1 - m_2) / M_2} \quad \text{.. (3.4.1)}$$

where M_1 is the molar mass of dihydropyrimidinone and M_2 is the molar mass of the solvent.

RESULTS AND DISCUSSION

The solubility (x) of synthesized compounds in the studied solvents is given in Tables 3.4.1 and 3.4.2. It is evident from these Tables that the solubility increases with temperature in both the solvents. Figure 3.4.1 shows the variation of mole fraction solubility (x) against temperature for RVG-1 in DMF and DMSO. Comparison of solubility of these compounds in DMF and DMSO shows that overall solubility is greater in DMF than that in DMSO. The dielectric constant and dipole moment of DMF (36.71, 3.86) are greater than that of DMSO (46.6, 3.9). Thus, these properties of solvent play an important role on the solubility, as expected.

The temperature dependence solubility in solvents is described by the modified Apelblat equation^{39, 40}

$$\ln x = A + B/(T/K) \quad \dots (3.4.1)$$

where x is the mass fraction solubility of compounds; T is the absolute temperature and A , and B are the coefficients. The values of these coefficients are given in Table 3.4.3. Using these values of A and B , calculated solubilities x_c were evaluated and are reported in Tables 3.4.1 and 3.4.2.

The relative deviations (RD) between the experimental and calculated values of solubilities are also calculated by equation 3.4.2 and are given in Tables 3.4.1 and 3.4.2.

$$Relative\ Deviation = \left(\frac{x - x_c}{x} \right) \quad \dots (3.4.2)$$

Further, relative average deviations (ARD) and root-mean-square deviations ($rmsd$) were calculated by equations 3.4.3 and 3.4.4 and are listed in Tables 3.4.5 and 3.4.6.

$$ARD = \frac{1}{N} \sum_i^N \left(\frac{x_i - x_c}{x_i} \right) \quad \dots (3.4.3)$$

$$rmsd = \left[\sum_{i=1}^N \frac{(x_{ci} - x_i)^2}{N-1} \right]^{1/2} \quad \dots (3.4.4)$$

where N is the number of experimental points.

Table 3.4.1: The experimental solubility (x), calculated solubility (x_c) and relative deviation (RD) of Dihydropyrimidinones in DMF at different temperatures.

| <i>Temp.</i> <i>K</i> | <i>x. 10²</i> | <i>x_c.10²</i> | <i>100 RD</i> | <i>x. 10²</i> | <i>x_c.10²</i> | <i>100 RD</i> |
|--------------------------|--------------------------|-------------------------------------|---------------|--------------------------|-------------------------------------|---------------|
| | RVG-1 | | | RVG-6 | | |
| 298.15 | 2.3942 | 2.4144 | -0.8458 | 2.5976 | 2.6249 | -1.0504 |
| 303.15 | 2.6012 | 2.5817 | 0.7047 | 2.9504 | 2.8821 | 2.3022 |
| 308.15 | 2.8144 | 2.7605 | 1.9148 | 3.3087 | 3.1645 | 4.3570 |
| 313.15 | 2.9711 | 2.9518 | 0.6128 | 3.5513 | 3.4746 | 2.1238 |
| 318.15 | 3.1325 | 3.1563 | -0.7603 | 3.7837 | 3.8151 | -0.8288 |
| | RVG-2 | | | RVG-7 | | |
| 298.15 | 2.0191 | 2.0246 | -0.2766 | 2.3994 | 2.4983 | -4.1223 |
| 303.15 | 2.2218 | 2.2465 | -1.1938 | 2.8305 | 2.8154 | 0.5165 |
| 308.15 | 2.4283 | 2.4927 | -2.6490 | 3.2556 | 3.1727 | 2.5475 |
| 313.15 | 2.7426 | 2.7658 | -0.9423 | 3.5743 | 3.5753 | -0.1491 |
| 318.15 | 3.0545 | 3.0689 | -0.4727 | 3.8799 | 4.0291 | -3.8439 |
| | RVG-3 | | | RVG-8 | | |
| 298.15 | 3.0658 | 3.1884 | -3.9961 | 2.7580 | 2.8242 | -2.4018 |
| 303.15 | 3.4312 | 3.4212 | 0.2564 | 3.0324 | 3.0502 | -0.6669 |
| 308.15 | 3.7873 | 3.6711 | 3.0691 | 3.3096 | 3.2943 | 0.4628 |
| 313.15 | 3.9324 | 3.9392 | -0.2331 | 3.5325 | 3.5579 | -0.7915 |
| 318.15 | 4.0719 | 4.2268 | -3.8062 | 3.7518 | 3.8427 | -2.4220 |
| | RVG-4 | | | RVG-9 | | |
| 298.15 | 3.2160 | 3.1933 | 0.7077 | 2.8882 | 2.8844 | 0.1294 |
| 303.15 | 3.5228 | 3.5114 | 0.2434 | 3.1615 | 3.2005 | -1.2028 |
| 308.15 | 3.8182 | 3.8613 | -1.1290 | 3.4357 | 3.5512 | -3.3619 |
| 313.15 | 4.2631 | 4.2461 | 0.3274 | 3.9025 | 3.9404 | -1.0350 |
| 318.15 | 4.7017 | 4.6691 | 0.6930 | 4.3692 | 4.3721 | -0.0681 |

| | RVG-5 | | | RVG-10 | | |
|---------------|--------------|--------|--------|---------------|--------|---------|
| 298.15 | 2.4148 | 2.3842 | 1.2646 | 3.0024 | 2.9135 | 2.9635 |
| 303.15 | 2.6904 | 2.6561 | 1.2597 | 3.2716 | 3.2343 | 1.0907 |
| 308.15 | 2.9593 | 2.9590 | 0.0114 | 3.5355 | 3.5905 | -1.5564 |
| 313.15 | 3.3421 | 3.2964 | 1.3060 | 4.0513 | 3.9860 | 1.5805 |
| 318.15 | 3.7220 | 3.6723 | 1.3365 | 4.5547 | 4.4250 | 2.8476 |

Table 3.4.2: The experimental solubility (x), calculated solubility (x_c) and relative deviation (RD) of Dihydropyrimidinones in DMSO at different temperatures.

| Temp. K | $x \cdot 10^2$ | $x_c \cdot 10^2$ | 100 RD | $x \cdot 10^2$ | $x_c \cdot 10^2$ | 100 RD |
|--------------------|----------------------------------|------------------------------------|---------------|----------------------------------|------------------------------------|---------------|
| | RVG-1 | | | RVG-6 | | |
| 298.15 | 2.2628 | 2.2440 | 0.8274 | 2.0230 | 2.0159 | 0.3471 |
| 303.15 | 2.3440 | 2.3216 | 0.5293 | 2.1122 | 2.0784 | 1.5994 |
| 308.15 | 2.4262 | 2.4019 | 1.0019 | 2.2013 | 2.1427 | 2.6623 |
| 313.15 | 2.5095 | 2.4850 | 0.9760 | 2.2285 | 2.2091 | 0.8727 |
| 318.15 | 2.5937 | 2.5709 | 0.8793 | 2.2957 | 2.2775 | 0.7942 |
| | RVG-2 | | | RVG-7 | | |
| 298.15 | 2.2130 | 2.2481 | -1.5850 | 1.9252 | 1.9241 | 0.0595 |
| 303.15 | 2.2867 | 2.2958 | -0.3961 | 2.0087 | 1.9797 | 1.4445 |
| 308.15 | 2.2805 | 2.3445 | -2.8086 | 2.0422 | 2.0369 | 0.2586 |
| 313.15 | 2.3589 | 2.3943 | -1.4980 | 2.1093 | 2.0958 | 0.6404 |
| 318.15 | 2.4174 | 2.4451 | -1.1454 | 2.1694 | 2.1563 | 0.6030 |
| | RVG-3 | | | RVG-8 | | |
| 298.15 | 2.1397 | 2.1818 | -1.9633 | 2.0242 | 2.0346 | -0.5148 |
| 303.15 | 2.2548 | 2.2685 | -0.6080 | 2.0901 | 2.1060 | -0.7603 |
| 308.15 | 2.2699 | 2.3587 | -3.9145 | 2.1661 | 2.1799 | -0.6404 |
| 313.15 | 2.3984 | 2.4525 | -2.2550 | 2.2394 | 2.2564 | -0.7623 |
| 318.15 | 2.5185 | 2.5500 | -1.2532 | 2.3227 | 2.3356 | -0.5575 |
| | RVG-4 | | | RVG-9 | | |
| 298.15 | 2.5097 | 2.4869 | 0.9096 | 2.1708 | 2.1619 | 0.4112 |
| 303.15 | 2.5813 | 2.5460 | 1.3666 | 2.2611 | 2.2378 | 1.0308 |
| 308.15 | 2.6329 | 2.6066 | 0.9997 | 2.3214 | 2.3163 | 0.2174 |
| 313.15 | 2.6902 | 2.6685 | 0.8055 | 2.4119 | 2.3976 | 0.5928 |
| 318.15 | 2.7675 | 2.7320 | 1.2842 | 2.4986 | 2.4818 | 0.6727 |

| | RVG-5 | | | RVG-10 | | |
|---------------|--------------|--------|---------|---------------|--------|---------|
| 298.15 | 2.1226 | 2.1778 | -2.6028 | 2.2280 | 2.2455 | -0.7820 |
| 303.15 | 2.2540 | 2.2498 | 0.1858 | 2.2978 | 2.3185 | -0.8990 |
| 308.15 | 2.3853 | 2.3241 | 2.5673 | 2.3782 | 2.3938 | -0.6567 |
| 313.15 | 2.4012 | 2.4008 | 0.0147 | 2.4663 | 2.4717 | -0.2192 |
| 318.15 | 2.4171 | 2.4801 | -2.6088 | 2.5123 | 2.5520 | -1.5813 |

Table 3.4.3: Coefficient A and B of equation 3.4.1, Relative Average Deviation (ARD), and root Mean Square Deviation (rmsd) of Dihydropyrimidinones in DMF and DMSO.

| Compounds | A | B | 10⁴ rmsd | 100 ARD |
|------------------|----------|----------|----------------------------|----------------|
| DMF | | | | |
| RVG-1 | -7.71 | 0.0134 | 0.36 | 0.32 |
| RVG-2 | -10.10 | 0.0208 | 0.45 | -1.10 |
| RVG-3 | -7.65 | 0.0141 | 4.21 | -0.94 |
| RVG-4 | -9.10 | 0.0190 | 0.29 | 0.16 |
| RVG-5 | -10.17 | 0.0216 | 0.51 | 1.03 |
| RVG-6 | -9.21 | 0.0187 | 2.62 | 1.38 |
| RVG-7 | -10.81 | 0.0239 | 3.13 | -1.01 |
| RVG-8 | -8.15 | 0.0154 | 1.12 | -1.16 |
| RVG-9 | -9.74 | 0.0208 | 1.33 | -1.12 |
| RVG-10 | -9.76 | 0.0209 | 2.65 | 1.38 |
| DMSO | | | | |
| RVG-1 | -5.82 | 0.0068 | 0.17 | 0.84 |
| RVG-2 | -5.05 | 0.0042 | 0.59 | -1.48 |
| RVG-3 | -6.15 | 0.0078 | 1.10 | -1.99 |
| RVG-4 | -5.09 | 0.0047 | 0.33 | 1.07 |
| RVG-5 | -5.76 | 0.0065 | 0.86 | -0.48 |
| RVG-6 | -5.72 | 0.0061 | 0.42 | 1.25 |
| RVG-7 | -5.65 | 0.0057 | 0.09 | 0.60 |
| RVG-8 | -5.95 | 0.0069 | 0.08 | -0.64 |
| RVG-9 | -5.89 | 0.0069 | 0.09 | 0.58 |
| RVG-10 | -5.70 | 0.0064 | 0.20 | -0.82 |

These values suggest that there is good agreement between experimental and calculated solubility values. So, the modified Apelblat equation can be used as model for the evaluation of solubility of these compounds in different solvents.

According to van't Hoff analysis, the standard enthalpy change of solution is obtained from the slope the $\ln x$ versus $1/T$ plot. However, in recent thermodynamic treatments, some modifications have been introduced in the van't Hoff equation to diminish the propagation of errors and consequently to separate the chemical effects from those due to statistical treatment used when enthalpy-entropy compensation plots are developed⁴¹. For this reason, the mean harmonic temperature (T_{hm}) is used in the van't Hoff analysis, which is calculated by the following equation.

$$T_{hm} = \frac{n}{\sum_i^n (1/T)} \quad \text{..... (3.4.5)}$$

where n is the number of temperatures studied and T is absolute temperature of the experiment. In the present case, the T_{hm} value obtained is 308 K.

So, the modified van't Hoff equation is^{42, 43}.

$$\frac{\partial \ln x}{\partial \left(\frac{1}{T} - \frac{1}{T_{hm}} \right)_P} = - \frac{\Delta H_{sol}}{R} \quad \text{..... (3.4.6)}$$

where ΔH_{sol} is the heat of solution and R is the gas constant.

Figure 3.4.2 shows the van't Hoff plots for RVG-1 in DMF and DMSO solutions. The slope of these linear plots gives the values of ΔH_{sol} whereas Gibb's free energy of dissolution (ΔG_{sol}) was evaluated from the intercept using the following equation⁴².


$$\Delta G_{sol} = -RT_{hm} \cdot \text{intercept} \quad \text{..... (3.4.7)}$$

Using these evaluated ΔH_{sol} and ΔG_{sol} values, the entropies of solutions ΔS_{sol} were obtained from equation⁴⁴ 3.4.8

$$\Delta S_{sol} = \frac{\Delta H_{sol} - \Delta G_{sol}}{T_{hm}} \quad \dots (3.4.8)$$

All these thermodynamic parameters are given in Tables 3.4.4.

Figure 3.4.1: The mole fraction solubility (x) against temperature (T/K) for RVG-1 in DMF (A) and DMSO (B).







Figure 3.4.2: van't Hoff plots for for RVG-1 in [A] DMF and [B] DMSO.






Table 3.4.4: The thermodynamic function of Dihydropyrimidinones in DMF and DMSO at 308 K (T_{hm}).

| Compound code | ΔH_{sol} $kcal.mol^{-1}$ | ΔG_{sol} $kcal.mol^{-1}$ | ΔS_{sol} $cal.mol^{-1}.K^{-1}$ | ΔH_{sol} $kcal.mol^{-1}$ | ΔG_{sol} $kcal.mol^{-1}$ | ΔS_{sol} $cal.mol^{-1}.K^{-1}$ |
|---------------|-------------------------------------|-------------------------------------|---|-------------------------------------|-------------------------------------|---|
| | DMF | | | DMSO | | |
| RVG-1 | 2.53 | 2.19 | 1.09 | 1.28 | 2.27 | -3.22 |
| RVG-2 | 3.90 | 2.26 | 5.33 | 0.78 | 2.30 | -4.93 |
| RVG-3 | 2.66 | 2.02 | 2.06 | 1.46 | 2.30 | -2.72 |
| RVG-4 | 3.57 | 1.99 | 5.15 | 0.89 | 2.22 | -4.32 |
| RVG-5 | 4.07 | 2.14 | 6.25 | 1.22 | 2.30 | -3.49 |
| RVG-6 | 3.54 | 2.10 | 4.67 | 1.15 | 2.34 | -3.84 |
| RVG-7 | 4.51 | 2.11 | 7.77 | 1.07 | 2.38 | -4.24 |
| RVG-8 | 2.90 | 2.09 | 2.61 | 1.29 | 2.34 | -3.40 |
| RVG-9 | 3.90 | 2.04 | 6.03 | 1.30 | 2.30 | -3.24 |
| RVG-10 | 3.94 | 2.02 | 6.21 | 1.20 | 2.28 | -3.50 |

It is evident from Table 3.4.4 that for all the compounds ΔH_{sol} and ΔG_{sol} values are positive whereas ΔS_{sol} values are negative for DMSO and positive values for DMF. When stronger bonds are broken and weaker bonds are formed, energy is consumed and so, ΔH_{sol} becomes positive⁴⁴. This indicates endothermic dissolution of compounds where the enthalpy term contributes to an unfavorable positive value of ΔG_{sol} ⁴⁴.

Thus, positive values of ΔG_{sol} indicates that the dissolution process is not spontaneous^{44, 45}.

The negative value of entropy indicates less randomness in solutions⁴⁴.

LITERATURE CITED

- (1) Shayanfar, A., Soltanpour, S., Jabbaribar, F., Hamidi, A. A., Acree. W. E., Jouyban, A. "Solubility of Anthracene in quaternary solvent mixtures of 2,2,4-Trimethyl pentane + 2-Propanone + Methanol + Alcohols at 298.15 K." *J. Chem. Eng. Data*, **2008**, 53, 2250–2253.
- (2) Nti-Gyabaah, J., Chmielowski, R., Chan, V., Chiew, Y. C. "Solubility of Lovastatin in a family of six Alcohols: Ethanol, 1-Propanol, 1-Butanol, 1-Pentanol, 1-Hexanol and 1-Octanol." *Int. J. Pharm.*, **2008**, 359, 111-117.
- (3) Niea, Q., Wang, J. K. "Solubility of 16, 17-Epoxyprogesterone in six different solvents. *J. Chem. Eng. Data*, **2005**, 50, 1750-1752.
- (4) Elder, J. P. "Thermal behavior of Lovastatin." *Thermochem. Acta.*, **1998**, 134, 41-45.
- (5) Ren, G. B., Wang, J. K. "Solubility of DL-p-hydroxyphenylglycine sulfate in binary acetone + water solvent mixtures." *J. Chem. Eng. Data*, **2004**, 49, 1376-1378.
- (6) Samuel, G. "Thermodynamics for Chemist", Litton Edu. Pub., Inc., New York, **1946**.

- (7) Fatima, L. M., Antonio, J. Q., Simao, P. P., Eugenia, A. M. "Aqueous Solubility of Some Natural Phenolic Compounds." *Ind. Eng. Chem. Res.* **2008**, 47, 5182–5189.
- (8) Sashina, E. S., Bochek, A. M., Novoselov, N. P., Kirichenko, D. A. "Structure and Solubility of Natural Silk Fibroin." *Russ. J. Appl. Chem.*, **2006**, 79, 869-876.
- (9) Daneshfar, A., Ghaziaskar, H. S., Homayoun, N. "Solubility of Gallic acid in Methanol, Ethanol, Water and Ethyl Acetate." *J. Chem. Eng. Data.*, **2008**, 53, 776-778.
- (10) Shalmashi, A., Eliassi, A. "Solubility of L-(+)-Ascorbic Acid in Water, Ethanol, Metanol, Propan-2-ol, Acetone, Acetonitrile, Ethyl Acetate and Tetrahydrofuran from (293 to 323) K" *J. Chem. Eng. Data.*, **2008**, 53, 1332-1334.
- (11) Voutsas, E. K., Panteli, E. K. "Solubilities of Cinnamic Acid Esters in Ionic liquids" *J. Chem. Eng. Data.*, **2009**, 54, 812-818.
- (12) Yuichi, H., Violette, D. "Effect of temperature on the solubility of aroma compounds in polyethylene film" *Poly. Test.*, **2006**, 25, 690-696.
- (13) Mansoori, G. A., Eliassi, A., Modarress, H. "Measurement of Activity of water in aqueous Poly(ethylene glycol)Solutions (Effect of Excess Volume on the Flory-Huggins X-Parameters)." *J. Chem. Eng. Data.*, **1999**, 44, 52-55.
- (14) Pradhan, A. A., Vera, J. H. "Effect of acids and bases on the solubility of amino acids". *Fluid Phase Equil.*, **1998**, 152, 121-132.
- (15) Macedo, E. A. "Solubility of amino acids, sugars, and proteins" *Pure Appl. Chem.*, **2005**, 77, 559–568.
- (16) Zhou, C. R., Kong, M. Z., Shi, X. H., Cao, Y. C. "Solubility of Imidaclopride in different solvents" *J. Chem. Eng. Data.*, **2008**, 53, 615–618.
- (17) Jouyban, A., Acree, W. E., Fakhree, M. A. A., Shayanfar, A. "Solubility of Lamotrigine, Diazepam and Clonazepam in Ethanol + Water mixtures at 298.15K." *J. Chem. Eng. Data.*, **2009**, 54, 1107-1109.
- (18) Yin, Q., Guo, K., Zhang, M., Wang, J. "Solubility of Losartan Potassium in different binary solvents from (293.15 to 343.15) K". *J. Chem. Eng. Data*, **2008**, 53, 1138-1140.
- (19) Ren, Q., Chen, W., Su, B., Xing, H., Yang, Y. "Solubility of Demosterol in five organic solvents." *J. Chem. Eng. Data.*, **2008**, 53, 2715-2717.
- (20) Song, C. Y., Shen, H. Z., Wang, L. C., Zhao, J. H., Wang, F. A. "Solubilities of Vitamin K₃ in benzene, toluene, ethylbenzene, o-Xylene, m-Xylene and

- p-Xylene between (299.44 and 344.24) K." *J. Chem Eng. Data*, **2009**, 53, 1245-1247.
- (21) James, W. H., Kristina, L. F., Mark, P. H., Markus, M. H. "Surprisingly high solubility of the ionic liquid trihexyltetradecylphosphonium chloride in dense carbon dioxide" *Green Chem.*, **2005**, 7, 475-478.
- (22) Takashi, K. "Mutual Solubility of Hydrophobic Ionic Liquids and Water in Liquid-Liquid Two-phase Systems for Analytical Chemistry". *Anal. Sci.*, **2008**, 24, 1221-1230.
- (23) Kumar, A., Shukla, J., Mohandas, V. P. "Effect of pH on the Solubility of $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ in Aqueous NaCl solutions and physicochemical solution properties at 35 °C" *J. Chem. Eng. Data*, **2008**, 53, 2797-2800.
- (24) Buryukin, F. A., Tverdokhlebov, V. P., Fedorov, V. A., Tetenkova, E. V., Fedorova, A. V. and Azanova, O. O. "The Solubility of Acetylferrocene and Diacetylferrocene in dimethylsulfoxide and its mixtures with water" *Russ. J. Phy. Chem. A.*, **2008**, 82, 1545-1548.
- (25) Yan, W., Dang, L., Wei, Z., Wei, H. "Solubility of Ammonium Thiocyanate in different solvents" *J. Chem. Eng. Data.*, **2009**, 54, 1063-1064.
- (26) Enrique, M., Jose, L. A. "Polymer Solubility Parameters of Poly (propylene oxide) Rubber from Inverse Gas Chromatography Measurements" *Poly. J.*, **1996**, 28, 127-130.
- (27) Price, G. J., Shillcock, I. M., "Inverse gas chromatographic measurement of solubility parameters in liquid crystalline systems." *J. Chromato. A*, **2002**, 964, 199-204.
- (28) Hou, G. Y., Yin, Q. X., Yang, Y., Hu, Y., Zhang, M. J., Wang, J. K. "Solubilities of adefovir dipivoxil in different binary solvents at 298.15 K." *J. Chem. Eng. Data.*, **2008**, 53, 1021-1023.
- (29) Kanji, S., Masahiro, T. "Solubility control by UV irradiation of a copolymer having both photo base generating groups and photo acid generating groups" *Chem. Commun.*, **1997**, 6, 853-854.
- (30) Bettina, H., Sabine, G., Michael, A., Christoph, S. "Comparison of Nephelometric, UV-Spectroscopic and HPLC Methods for High-Throughput Determination of Aqueous Drug Solubility in Microtiter Plates" *Anal. Chem.* **2009**, 81, 3165-3172.
- (31) Jaroslav, N. "Solid-Liquid Phase Equilibria" *Elsevier Sci. Pub.Co. New York*, **1977**.
- (32) Ma, P. S., Xia, Q. "Determination and Correlation for Solubility of Aromatic Acids in Solvents." *Chin. J. Chem. Eng.*, **2001**, 9, 39-44.

- (33) Wei, D., Pei, Y. "Measurement and correlation of solubility of diphenyl carbonate in alkanols." *Ind. Eng. Chem. Res.*, **2008**, 47, 8953–8956.
- (34) Shipra, B., Rahul, B., Mehul, B., Nayan, V., Ravi G. "Solubility of Enrofloxacin Sodium in Various Solvents at Various Temperatures" *J. Chem. Eng. Data.*, **2008**, 53, 2897–2899.
- (35) Shipra, B., Ravi, G., B., Rahul, Nayan, V. and Mehul B. "Solubility of Difloxacin in Acetone, Methanol, and Ethanol from (293.15 to 313.15) K" *J. Chem. Eng. Data.*, **2009**, 54, 1091–1093.
- (36) Shipra, B. "Physicochemical Studies of Some Schiff Bases Derived From 6-Ethylbenzene 1,3-diol" *E- J. Chem.*, **2004**, 1, 199–205.
- (37) Riddick, J. A., Bunger, W. B., Sakano, T. "Organic Solvents-Physical Properties and methods of purification." *4th Edi., Techniques of Chem.*, II, A Wiley-Interscience Publication, John Wiley, New York **1986**.
- (38) Zhu, M. "Solubility and Density of the Disodium Salt Hemiheptahydrate of Ceftriaxone in Water + Ethanol Mixtures." *J. Chem. Eng. Data.*, **2001**, 46, 175–176.
- (39) Apelblat, A., Manzurola, E. "Solubilities of o-acetylsalicylic, 4-aminosalicylic, 3,5-di nitrosalicylic, and p-toluic acid, and magnesium-DL-aspartate in water from $T = (278 \text{ to } 348) \text{ K}$." *J. Chem. Thermo.*, **1999**, 31, 85–91.
- (40) Gao, J., Wang, Z. W., Xu, D. M., Zhang, R. K. "Solubilities of Triphenylphosphine in Ethanol, 2-Propanol, Acetone, Benzene and Toluene." *J. Chem. Eng. Data*, **2007**, 52, 189–191.
- (41) Aragon, D. M., Ruidiaz, M. A., Vargas, E. F., Bregni, C., Chiappetta, D. A., Sosnik, A., Martinez, F. "Solubility of the antimicrobial agent Triclosan in organic solvents of different hydrogen bonding capabilities at several temperatures." *J. Chem. Eng. Data*, **2008**, 53, 2576–2580.
- (42) Krug, R. R., Hunter, W. G., Grieger, R. A. "Enthalpy-entropy compensation. 2. Separation of the chemical from the statistical effects." *J. Phys. Chem.*, **1976**, 80, 2341–2351.
- (43) Bustamante, P., Romero, S., Pena, A., Escalera, B., Reillo A. "Nonlinear enthalpy-entropy compensation for the solubility of drugs in solvent mixtures: paracetamol, acetanilide and nalidixic acid in dioxane-water." *J. Pharma. Sci.*, **1998**, 87, 1590–1596.
- (44) Kalsi, P. S. "Organic reactions and their mechanisms," *New age international (P) limited- New Delhi, 2nd edition*, **2004**, 119.
- (45) El-Bindary, A. A., El-Sonbati, A. Z., El-Mosalamy, E. H., Ahmed, R. M. "Potentiometric and Thermodynamic Studies of Azosulfonamide Drugs. X", *Chem. Pap.*, **2003**, 57, 255–258.



Section-V

Thermal Properties

INTRODUCTION

Studies on thermal properties of substances are of great importance from both scientific and practical of view. Scientific and technological achievements together with demands based on industrial requirement have permitted the development of various types of materials that can withstand at much higher temperatures and more corrosive environments.

Various techniques have been developed for the measurement of thermal properties. Some of the most commonly used techniques are Differential Scanning Calorimetry (DSC), Differential Thermal Analysis (DTA), Thermo Gravimetric Analysis (TGA), Evolved Gas Detection (EGD), Evolved Gas Analysis (EGA) etc. In the present study, thermal analysis is done by DSC and TGA techniques.

Thermogravimetric analysis covers a wide spectrum of thermoanalytical techniques, which monitor one or more physical properties of a substance that is undergoing a temperature programmed heating as a function of time or temperature. This technique is widely used in the fields of gas-solid interactions, fuels, catalysis, polymer and chemical synthesis. This method can not distinguish the actual nature of the material evolved in the course of the process and is also handicapped in resolving overlapped thermal events.

This technique has been used in analyzing and characterizing wide variety of materials like coals, rocks and minerals¹, petroleum coke², rubber and styrene butadiene blends³, hydrocarbon sludge from petrochemical plants⁴, polyimide resins⁵, superconducting materials⁶, hydrogen storage materials⁷, and even rocks from moon.

Apart from its traditional use in investigations of chemical reactions, temperature programmed decompositions, study of reaction kinetics and mechanism, determination of material purity etc., this technique is also used in the areas of forensic science⁸, archeology^{9, 10}, catalysis^{11, 12}, polymers¹³⁻¹⁷, nanotubes^{18, 19}, and pharmaceuticals²⁰⁻²⁵.

One of the most important applications of thermo gravimetry is to determine thermal stability of polymer and binding materials. Houriez and co-workers²⁶ have reported the thermal analysis of polymer dispersed liquid crystal by using photo pyroelectric technique. Kurt and Trilogica²⁷ have

reported the quality assurance of polymer and plastic by thermal analysis and differential scanning calorimetry. The thermal study of honey²⁸, tobacco²⁹ and edible oil³⁰ has also been reported. Bei et al.³¹ have reported the thermal decomposition kinetic of 5-flourouracil from thermogravimetric analysis.

The decomposition temperature, adsorption and desorption phenomena, ash content, flame retardancy, rates of evaporation or sublimation can also be determined by TGA³². The information from TGA can be used to compliment differential scanning calorimetry (DSC) data. Some instruments are equipped to acquire both DSC and TGA data simultaneously. In this case, the DSC data provides the thermodynamic account of different chemical and physical reactions occurring and the TGA data provides an account of any mass changes that are coupled to the reactions. TGA method is useful for plant engineers in designing safe and economical plant and process for the manufacture of various chemicals.

DSC is a versatile thermal technique used to provide quantitative and qualitative information about physical and chemical changes involving endothermic or exothermic processes or heat capacity changes. It also provides useful information about crystallinity, stability of crystallites, glass transition temperature, cross-linking and heat of polymerization etc.

Literature survey shows that several investigations have been carried out on the application of thermal methods in pharmaceutical industry. Mamede et al.³³ have reported thermal behavior of some pharmaceuticals and drugs. Bogdan and co-workers³⁴ have reported the thermal stability and kinetic decomposition of diclofenac sodium. Lapuerta and co-workers³⁵ have reported the thermogravimetric analysis of diesel particulate matter. Wu and Hung³⁶ have done the thermal analysis of thin films of 1,4-phenylene-cis-benzobisoxazole or PBO. Thermal studies of some organic and inorganic compounds have also been studied³⁷⁻⁴⁵. However, little work has been done of thermal analysis of pyrimidine moiety⁴⁶⁻⁴⁸.

In the present study, thermal properties of some new synthesized dihydropyrimidinones have been studied by DSC and TGA techniques. Using thermograms, various kinetic parameters have also been evaluated.

THEORY

From TGA curves, various kinetic parameters can be evaluated by several methods. In all these methods, it is assumed that thermal and diffusion barriers are negligible because small quantity of material is used. The shape of any TGA curve depends on the nature of apparatus and the way in which it is used. Further, Arrhenius equation is valid in all these methods.

The kinetic treatments are generally based on the relationship of the type:

$$dC/dt = K f(C) \quad \dots (3.5.1)$$

where C is the degree of conversion, t is time and K is rate constant. $f(C)$ is a temperature independent function of C .

The constant K is assumed to have the Arrhenius form:

$$K = A e^{-E/RT} \quad \dots (3.5.2)$$

C can also be defined as:

$$C = 1 - (W/W_0) \quad \dots (3.5.3)$$

where W_0 and W are the initial weight at $t=0$ and weight at any time t of the material.

Equation (3.5.3) can be written as:

$$(W/W_0) = (1-C) \quad \dots (3.5.4)$$

W/W_0 is known as residual weight fraction.

Thus, the rate of conversion is,

$$dC/dt = - (1/W_0) (dW/dt) \quad \dots (3.5.5)$$

For homogeneous kinetics, the conversion is assumed to be of the form:

$$f(C) = (1-C)^n \quad \dots (3.5.6)$$

where n is the order of the reaction.

Substituting the values from equation (3.5.2) and (3.5.6) in equation (3.5.1) gives:

$$dC/dt = A e^{-E/RT} (1-C)^n$$

$$\text{or} \quad dC/dt = (A/\beta) e^{-E/RT} (1-C)^n \quad \dots (3.5.7)$$

where A is the frequency factor, β is the rate of heating and E is the energy of activation.

Various methods for single and multiple heating rates have been reported. The methods of single heating rate are as follows:

1. Freeman-Carroll⁴⁹ and Anderson-Freeman Method⁵⁰:

At a single heating rate, Freeman and Carroll gave the following relation to analysis TGA data:

$$\ln (dC/dt) / \ln (1-C) = n - E/R [(1/T) / (\Delta \ln(1-C))] \quad \dots (3.5.8)$$

A plot of left hand side against $(1/T) / (\Delta \ln(1-C))$ gives a straight line with a slope equal to $-E/R$ and the intercept is equal to n .

Anderson and Freeman then derived the following equation by using equation (3.5.8):

$$(\Delta \ln[dC/dt]) = n (\Delta \ln(1-C)) - E/R \Delta(1/T) \quad \dots (3.5.9)$$

The plot of $(\Delta \ln[dC/dt])$ against $(\Delta \ln(1-C))$ for equal intervals of $\Delta(1/T)$ gives a straight line with slope equal to n and intercept $-E/R \Delta(1/T)$.

2. Sharp-Wentworth method⁵¹:

To analyse the TGA data for first order kinetics ($n=1$), Sharp and Wentworth gave the relation:

$$\log [(dC/dt)/(1-C)] = \log (A/\beta) - (E/2.303R) \cdot (1/T) \quad \dots (3.5.10)$$

The plot of $\log [(dC/dt)/(1-C)]$ against $1/T$ would be a straight line with slope equal to $-(E/2.303R)$ and intercept equal to $\log (A/\beta)$.

3. Chatterjee Method⁵²:

Based on the weight units, the following relation was developed by Chatterjee:

$$n = [\log(dW/dt)_1 - \log(dW/dt)_2] / (\log W_1 - \log W_2) \quad \dots (3.5.11)$$

where W_1 and W_2 are the sample weights.

4. Horowitz and Metzger method⁵³:

In this method, the value of energy of activation E can be determined from a single TG curve by the relation:

$$\ln [\ln(1-C)^{-1}] = (E/RT_s^2)\theta \quad \dots (3.5.12)$$

where $\theta = T - T_s$. T_s is the temperature at which the rate of decomposition is maximum. The frequency factor A and entropy change ΔS can be determined by the following equations:

$$\ln E - \ln (RT_s^2) = \ln A - \ln \beta - E/RT_s \quad \dots (3.5.13)$$

$$A = (k_b T / h) e^{\Delta S/R} \quad \dots (3.5.14)$$

where k_b is Boltzmann constant and h is Planck's constant.

EXPERIMENTAL

Thermo gravimetric analysis (TGA) and Differential Scanning Calorimetry (DSC) measurements were made on the instrument “Pyris-1, Perkin Elmer Thermal Analysis” at the heating rate of 10°C/min in nitrogen atmosphere for all the dihydropyrimidinones.

RESULTS AND DISCUSSION

The TGA thermo grams of RVG-9 and RVG-10 are given in Figure 3.5.1. Various thermal properties such as initial decomposition temperature (IDT), the decomposition temperature range and the maximum degradation along with the percentage weight loss and endothermic transitions are reported in Table 3.5.1.

For some compounds, degradation is single step process whereas for others, it is multi step process. For RVG-1, and RVG-3, multi step degradation takes place whereas in other compounds, degradation is single step process.

Table 3.5.1 shows that RVG-9 and RVG-10 is unstable whereas RVG-6 and RVG-8 is most stable followed by RVG-4. RVG-10 has meta chloro, para fluoro side chain whereas RVG-9 has two chloro groups at meta and para positions. In other compounds, various substituent groups are attached. Thus, thermal stability is affected by the substituent group. Further, the positions of substituent also affect stability of the studied compounds. When chloro group is present at meta position (as in RVG-8) stability is higher than RVG-4, containing chloro group at para position. However, the presence of chloro groups at both meta and para positions make the compound most unstable as in RVG-9. Similarly, the presence of fluoro group at para position (iRVG-6) increases the stability.

Various kinetic parameters, such as order of the degradation (n), energy of activation (E), frequency factor (A) and entropy change (ΔS°) have also been calculated from the thermograms for each step and are reported in Table 3.5.2.

It is evident from Tables 3.5.2 that order of reaction is quite different in different steps for different dihydropyrimidinones. For single step degradation compound, order of reaction varies from 1.75 to 10.80, whereas for multi steps it varies from 2.68 to 9.31.

Figure 3.5.1: The TGA graphs of RVG-9 and RVG-10.







Table 3.5.1: TGA/DSC data for synthesized Dihydropyrimidinone derivatives.

| Comp. Code | Amt. mg. | Initial Decomp. Temp. °C | Decomp. range °C | % Wt. loss | Residual Wt. Loss mg. | Transition | DSC °C | Open capillary method °C | ΔH J.g⁻¹ |
|-----------------------|---------------------|---|---------------------------------|-----------------------|--------------------------------------|-------------------|-------------------|---|---|
| RVG-1 | 0.933 | 150 | 150-700 | 97.96 | 0.913 | Endo. | 260.8 | 257 | 139.04 |
| RVG-2 | 2.714 | 170 | 170-550 | 90.69 | 2.461 | Endo. | 265.7 | 268 | 169.50 |
| RVG-3 | 3.247 | 165 | 165-480 | 59.86 | 1.943 | Endo. | 271 | 275 | 183.66 |
| RVG-4 | 2.527 | 175 | 175-450 | 58.81 | 1.481 | Endo. | 256 | 253 | 121.27 |
| RVG-5 | 2.616 | 145 | 145-510 | 58.85 | 1.539 | Endo. | 241 | 239 | 127.32 |
| RVG-6 | 9.477 | 210 | 210-610 | 57.33 | 5.433 | - | - | - | - |
| RVG-7 | 7.571 | 150 | 150-380 | 84.05 | 6.363 | - | - | - | - |
| RVG-8 | 4.175 | 210 | 210-485 | 72.50 | 3.026 | - | - | - | - |
| RVG-9 | 5.834 | 100 | 100-500 | 75.37 | 4.397 | - | - | - | - |
| RVG-10 | 8.599 | 100 | 100-490 | 52.01 | 4.472 | - | - | - | - |

Table 3.5.2: The kinetic parameters of Dihydropyrimidinone derivatives.

| Comp. code | <i>n</i> | <i>E</i> <i>kJ.mol⁻¹</i> | <i>A</i> <i>s⁻¹</i> | ΔS <i>J.mol⁻¹.K⁻¹</i> |
|---|-----------------|--|---|---|
| <i>RVG-1 1st step</i> | 1.66 | 339.64 | 3.46×10^{17} | 234.18 |
| <i>RVG-1 2nd step</i> | 2.68 | 20.216 | 0.1473 | -117.51 |
| <i>RVG-2</i> | 1.78 | 130.41 | 4.55×10^7 | 46.72 |
| <i>RVG-3 1st step</i> | 7.42 | 296.06 | 1.11×10^{21} | 303.86 |
| <i>RVG-3 2nd step</i> | 9.31 | 26.894 | 2.6752 | -90.88 |
| <i>RVG-4</i> | 8.55 | 171.10 | 5.48×10^{11} | 125.67 |
| <i>RVG-5</i> | 10.80 | 209.76 | 6.34×10^{14} | 184.39 |
| <i>RVG-6</i> | 9.34 | 2.11 | 0.0043 | -143.46 |
| <i>RVG-7</i> | 3.64 | 50.56 | 734.03 | -43.42 |
| <i>RVG-8</i> | 4.09 | 150.09 | 1.24×10^{11} | 113.97 |
| <i>RVG-9</i> | 1.75 | 265.37 | 2.73×10^{21} | 312.33 |
| <i>RVG-10</i> | 4.06 | 305.05 | 2.32×10^{24} | 368.29 |

Figure 3.5.2: The DSC graphs of RVG-2 and RVG-3.



For single step degradation compounds, energy of activation (E) is maximum for RVG-10 and minimum for RVG-6. The frequency factor (A) also varies in the same order. For multi step degradation compounds, in first step, RVG-1 is found to be maximum whereas in second step, RVG-3 is maximum. The frequency factor A is maximum for RVG-3 in first step and minimum for RVG-1 in second step.

Further, change in entropy (ΔS^0) for all these reactions were calculated by equation (3.5.14) and are reported in Table 3.3.2. These values are both positive and negative for different compounds. The positive values of ΔS^0 indicate that the transition state is less ordered than the original compound whereas negative value of ΔS^0 corresponds to an increase in the order of transition state than the reactants. The highest negative value of ΔS^0 is in RVG-6 which has lowest energy of activation (E) and frequency factor (A). The same but opposite order has shown in RVG-10 for single step degradation.

Further, Table 3.5.1 shows DSC data along with the melting temperature determined by open capillary method. It is observed that the melting temperatures determined by the two methods are in good agreement. However, no correlation could be established between heat of reaction, kinetic parameters, melting temperature, thermal stability and substitution group.

Thus, it is concluded that thermal stability depends not only on type of substitution but also its position in aromatic ring skeleton. Further, different compounds have different rates of decomposition.

LITERATURE CITED

- (1) Heide, K., Gerth, K., Buchel, G., Hartmann, E. "EGA [evolved gas analysis] -a fingerprint characterization of minerals and rocks." *J. Thermal Anal.* **1997**, 48, 73-81.
- (2) Zimny, T., Weber, J. V., Fixari, B. "Coal-tar pitch/sulfur and coal-tar pitch/coke/sulfur mixtures. Study of the thermal behavior and the optical texture." *Petrochemie*, **1994**, 47(6), 225-259.
- (3) Ramesan, M. T. "Thermogravimetric analysis, flammability and oil resistance properties in natural rubber and dichlorocarbene modified styrene butadiene rubber blends." *Reactive and Functional Poly.*, **2004**, 59, 267-274.
- (4) Remmler, M., Kopinke, F. D., Stottmeister, U. "Thermoanalytical methods for characterizing hydrocarbon-sludge-soil mixtures." *Thermochim. Acta.*, **1995**, 263, 101-112.
- (5) Ozawa, T., Ariei, T., Kishi, A. "Thermogravimetry and evolved gas analysis of polyimide." *Thermochim. Acta.*, **2000**, 352-353, 177-180.
- (6) Wai, S. K., Hamid, N. A., Noor, E. S., Yew, P. J., Ismail, A. B., Ahmad, B. "Electromagnetic and thermal analysis of high-T_c superconductor in application of current limiting devices." *J. Electroceramics*, **2008**, 21, 365-369.
- (7) Kamruddin, M., Ajikumar, P. K. Dash, S., Tyagi, A. K., Baldev, R. "Thermogravimetry-evolved gas analysis-mass spectrometry system for materials research." *Bull. Mater. Sci.*, **2003**, 26(4), 449-460.
- (8) Raja, S., Thomas, P. S., Stuart, B. H., Guerbois, J. P., O'Brien, C. "The estimation of pig bone age for forensic application using thermogravimetric analysis." *J. Thermal Anal. Calor.*, **2009**, 98(1), 173-176.
- (9) Frost, R. L., Wills, R. A., Weier, M. L., Musumeci, A. W., Martens, W. "Thermal decomposition of natural and synthetic plumbojarosites: Importance in 'archeochemistry'." *Thermochim. Acta*, **2005**, 432(1), 30-35.
- (10) Sanchez, R. S., Bosch, R. F., Gimeno, A. J. V., Yusa, M. D. J., Domenech, C. A. "Application of XRF, XRD, thermal analysis, and voltammetric techniques to the study of ancient ceramics." *Anal. Bioanal. Chem.*, **2002**, 373(8), 893-900.
- (11) Youssef, T. A., Mekewi, M. A. "Thermal stability eminence of PMMA prepared in presence of photosynthesized ruthenium carbonyl Schiff base as a catalyst." *J. Appl. Poly. Sci.*, **2009**, 114(3), 1503-1510.

- (12) Babenko, V. S.; Pakhomov, N. A.; Buyanov, R. A. "Study of thermal stability of aluminum-chromium catalysts for a single-stage process of butane dehydrogenation." *Kataliz v Promyshlennosti*, **2009**, 1, 13-19.
- (13) Ajloo, D., Sharifian, A., Behniafar, H. "Prediction of thermal decomposition temperature of polymers using QSPR methods." *Bull. Kor. Chem. Soc.*, **2008**, 29(10), 2009-2016.
- (14) Radhakrishnan, S., Somanathan, N., Thelakkat, M. "Thermal Degradation Studies of Polythiophenes Containing Hetero Aromatic Side Chains." *Inter. J. Thermophys.*, **2009**, 30(3), 1074-1087.
- (15) Erol, I. "Synthesis and characterization of a new methacrylate polymer with side chain benzofurane and cyclobutane ring: thermal properties and antimicrobial activity." *High Perform. Poly.*, **2009**, 21(4), 411-423.
- (16) Lazaro, M., Juan, M., Chattah, A. K., Monti, G. A., Leal, D. M. F., Buldain, G. Y., Campo Dall'Orto, Viviana. "New copper(II) complexes of polyampholyte and polyelectrolyte polymers: Solid-state NMR, FTIR, XRPD and thermal analyses." *Polymer*, **2008**, 49(25), 5482-5489.
- (17) Koptelov, I. A.; Karyazov, S. V. "Methodological errors in studying of polymer thermal degradation kinetics. I. Differential thermal analysis." *Plasticheskie Massy*, **2008**, 7, 24-28.
- (18) Feng, X., Liu, K., Xie, X., Zhou, R., Zhang, L., Li, Q., Fan, S., Jiang, K. "Thermal Analysis Study of the Growth Kinetics of Carbon Nanotubes and Epitaxial Graphene Layers on Them." *J. Phys. Chem. C*, **2009**, 113(22), 9623-9631.
- (19) Lewicki, J. P., Liggat, J. J., Hayward, D., Pethrick, R. A., Patel, M. "Degradative thermal analysis and dielectric spectroscopy studies of aging in polysiloxane nanocomposites." *ACS Symposium Series: Polymer Degradation and Performance.*, **2009**, 1004, 239-254.
- (20) Ulu, S. T., Elmali, F. T. "UV-Second Derivative Spectrophotometric and Colorimetric Methods for the Determination, Validation and Thermogravimetric Analysis of New Oral Antidiabetic Pioglitazone in Pure and Pharmaceutical Preparations." *Anal. Lett.*, **2009**, 42(14), 2254-2270.
- (21) Salvio, N. H., Novak, C., Matos, J. R. "Thermal analysis and compatibility studies of prednicarbate with excipients used in semi solid pharmaceutical form." *J. Thermal Anal. Calor.*, **2009**, 97(1), 367-374.
- (22) Matthews, K. H., Stevens, H. N. E., Auffret, A. D., Humphrey, M. J., Eccleston, G. M. "Formulation, stability and thermal analysis of lyophilized wound healing wafers containing an insoluble MMP-3 inhibitor and a non-ionic surfactant." *Int. J. Pharma.*, **2008**, 356(1-2), 110-120.

- (23) Simonis, L., Ruminski, M., Primiano, L., Sood, M., Riga, A. T., Alexander, K. S. "Quality control and identification of dental materials by thermal analysis." *Proc. NATAS Ann. Conf. Thermal Analysis and Applications*, **2007**.
- (24) Gardikis, K., Demetzos, C. "Differential scanning calorimetry (DSC): a tool for studying the thermal behavior of pharmaceutical products and designing lipidic carriers incorporating bioactive molecules." *Pharmaceutike*, **2006**, 19(3), 68-76.
- (25) Lerdkanchanaporn, S., Dollimore, D., Alexander, K. S. "A simultaneous TG-DTA study of the degradation in nitrogen of cellulose to carbon alone and in the presence of other pharmaceutical excipients." *Thermochim. Acta.*, **1998**, 324, 25-32.
- (26) Houriez, N., Delenclos, S., Longuemart, S., Dadarlat, D., Hadj, S. A. "Thermal analysis of Polymer Dispersed Liquid Crystal under electric field using photopyroelectric technique." *Laser Phys.*, **2009**, 19(6), 1345-1349.
- (27) Kurt, P., Trilogica G. "Quality assurance of polymers by thermal analysis." *CHEMagazin.*, **2008**, 18(1), 22-23.
- (28) Maria, L. F., Cristiane, B. C., Jivaldo, R. M., Lgia, B. A. M., Roy, E. B., "Optimization of Thermogravimetric Analysis of Ash Content in Honey." *J. Brazil. Chem. Soc.*, **2004**, 15(6), 797-802.
- (29) Wang, H. B., Guo, J. W., Xia, Q. L.; Fu-Wei, X., Hui-Min, L. "Thermogravimetric analysis of several domestic tobacco samples." *Yancao Keji*, **2009**, 9, 47-49.
- (30) Heda, L. C., Sharma, R., Tank, P., Sherwani, M. R. K. "Thermogravimetric analysis of copper (II) soaps derived from edible oils." *J. Lipid Sci. Tech.* **2008**, 40(1), 6-9.
- (31) Bei, Y. L., Qing-yang, L., Gui-bin, Q., Yuan-jun, D. "Thermal decomposition kinetics of 5-fluorouracil from thermogravimetric analysis." *Korean J. Chem. Eng.*, **2008**, 25(5), 980-981.
- (32) Rebecca, L. D. "Polymer Content in Bulk Molding Compound Using Thermogravimetric Analysis" *Alfred Uni., NTR-105621*, **2008**, 1-3.
- (33) Mamede, L. C., Caetano, B. L., Rocha, L. A., Ferreira, E. M., Cestari, A., Kfuri, C. R., Ciuffi, K. J., Calefi, P. S., Mello, C., Cunha, W. R., Nassar, E. J. "Thermal behavior of some pharmaceuticals and drugs." *Revista de Ciencias Farmaceuticas Basica e Aplicada.*, **2006**, 27(2), 151-155.
- (34) Bogdan, T., Fulias, A., Marian, E., Tita, D. "Thermal stability and decomposition kinetics under non-isothermal conditions of sodium diclofenac." *Rev. Chim*, **2009**, 60(5), 524-528.

- (35) Lapuerta, M., Ballesteros, R., Rodriguez-Fernandez, J. "Thermogravimetric analysis of diesel particulate matter." *Meas. Sci. Technol.*, **2007**, 18, 650–658.
- (36) Wu, G. M., Hung, C. H. "Mechanical properties and thermogravimetric analysis of PBO thin films." *J. Achiev. Mater. Manufact. Eng.*, **2006**, 17, 27-32.
- (37) Mamede, L. C., Caetano, B. L., Rocha, L. A., Ferreira, E. M., Cestari, A., Kfuri, C. R., Ciuffi, K. J., Calefi, P. S., Mello, C., Cunha, W. R., Nassar, E. J. "Thermal behavior of some pharmaceuticals and drugs." *Rev. de Ciencias Farmaceu. Basica Aplicada*, **2006**, 27(2), 151-155.
- (38) Zayed, M. A., Hawash, M. F., Fahmey, M. A., El-Habeeb, A. A. "Structure investigation of sertraline drug and its iodine product using mass spectrometry, thermal analyses and MO-calculations." *Spectrochim. Acta, A: Mol. Biomol. Spectr.*, **2007**, 68A(3), 970-978.
- (39) Albers, D., Galgoci, M., King, D., Miller, D., Newman, R., Peerey, L., Tai, E., Wolf, R., "Characterization of the Polymorphic Behavior of an Organic Compound Using a Dynamic Thermal and X-ray Powder Diffraction Technique." *Org. Proc. Res. Devel.*, **2007**, 11(5), 846-860.
- (40) Gardikis, K., Demetzos, C. "Differential scanning calorimetry (DSC): a tool for studying the thermal behavior of pharmaceutical products and designing lipidic carriers incorporating bioactive molecules." *Pharmakeutike*, **2006**, 19(3), 68-76.
- (41) Arslan, H., Dondas, H. A. "Thermal behavior of some spirobenzodiazepine derivatives." *Thermochim. Acta*, **2000**, 354(1-2), 107-115.
- (42) Randhawa, H. S., Sekhon, B. S., Sahai, H. K., Lakhani, R. "Thermal studies of metal complexes formed in Ag(I)-, Cd(II)-, UO₂(II)-, VO(II), Co(II)- and Hg(II)-sulphamethoxazole systems." *J. Thermal Anal. Calori.*, **1999**, 57(2), 551-558.
- (43) Landspersky, H., Balek, V., Urbanek, V. "Application of thermal analysis methods to the study of the preparation of sintered uranium dioxide. II. Modeling of the heat treatment of uranium(VI) xerogels during calcinations." *Jaderna Energie.*, **1979**, 25, 46-52.
- (44) Wei, L., Ma, Y., Duan, J., Wu, Z. "Determination of the inorganic composition contents in polymer blend by TGA." *Gong. Suliao Ying.*, **2001**, 29, 26-27.
- (45) Xi, Y., Martens, W., He, H., Frost, R. L. "Thermogravimetric analysis of organoclays intercalated with the surfactant octadecyltrimethylammonium bromide." *J. Therm. Anal. Calor.*, **2005**, 81, 91-97.

- (46) Moreno-Carretero, M. N., Salas-Peregrin, J. M., Mata-Arjona, A. "Thermal studies on metal complexes of 5-nitrosopyrimidine derivatives. I. Thermal behavior of zinc(II) complexes of 6-amino-5-nitrosouracil derivatives. *J. Thermal Anal.*, **1984**, 29(3), 553-559.
- (47) Sanchez, S. M. P.; Salas, P. J. M., Romero, M. M. A. Colacio, R. E., Moreno, C. M. "Thermal studies on metal complexes of 2-thiopyrimidines." *Thermochim. Acta.*, **1985**, 93, 77-80.
- (48) Sanchez-Sanchez, M. P., Salas-Peregrin, J. M., Romero-Molina, M. A., Ruiz Sanchez, J. "Thermal studies on metal complexes of 5-nitrosopyrimidine derivatives. VI. Thermal behavior of zinc(II), cadmium(II), mercury(II) and (I) complexes of 4-amino-1,6-dihydro-2-methylthio-5-nitroso-6-oxopyrimidine." *J. Thermal Anal.*, **1986**, 31(3), 573-80.
- (49) Freeman, E. S., Carroll, B. "The application of thermoanalytical techniques to reaction kinetics. The thermogravimetric evaluation of the kinetics of the decomposition of calcium oxalate monohydrate." *J. Phys. Chem.*, **1958**, 62, 394-397.
- (50) Anderson, D. A., Freeman, E. S. "Kinetics of the thermal degradation of polystyrene and polyethylene." *J. Polym. Sci.*, **1961**, 54, 253-260.
- (51) Sharp, J. H., Wentworth, S. A. "Kinetic analysis of thermogravimetric data." *Anal. Chem.*, **1969**, 41, 2060-2062.
- (52) Chatterjee, P. K. "Application of thermogravimetric techniques to reaction kinetics." *J. Polym. Sci.*, **1965**, 3, 4253-4262.
- (53) Horowitz, H. H., Metzger, G. "New analysis of thermogravimetric traces." *Ana. Chem.*, **1963**, 35, 1464-1468.



Section-VI

Dissociation Constants

INTRODUCTION

Dissociation constants are important parameters to indicate the extent of ionization of molecules in solution. The dissociation constant or acidity constants of organic compounds play a fundamental role in many analytical procedures such as acid-base titration, solvent extraction, complex formation and ion transport¹. It has been shown that the acid-base properties affect the toxicity², chromatographic retention behavior and pharmaceutical properties³ of organic acids and bases. Much of the theoretical foundation of modern organic chemistry is based on the observation of the effects on acid-base equilibria of changing molecular structure⁴.

Various workers have been reported the various method to determine the dissociation constants. Several methods for the determination of dissociation constants, such as potentiometric titration, spectrophotometric determination, conductometry and polarographic method have been reported⁵⁻⁹. Datta et al have reported the determination of dissociation constants from molar conductance¹⁰. Lahiri and co-workers have been reported the radiotracer method for determine the dissociation constants¹¹. Sager and Siewers have been measured the dissociation constant of 4-aminobenzophenone from ultraviolet spectra¹². Potentiometry is mostly used for the determination of dissociation constants of acids because it is economical of time. Further it can be used for acids of pKa range from 2 to 11 units.

For this measurement, glass and calomel electrodes have been used and carbonate free potassium hydroxide is the best alkali to use as a titrant. In contrast to potentiometry, the conductimetry provides a simple and accurate method for the determination of dissociation constants of acids without any primary and secondary buffer solutions required for calibration of the platinum electrode^{13, 14}.

The determination of the dissociation constant of an acid in mixtures of organic compounds with water provides useful data for the theoretical understanding of the ionization process in this media. The dissociation constants are also important for tautomeric equilibria and solvent-solute interaction¹⁵. The effect of solvent composition on the dissociation equilibria of

organic reagents is of great importance in chemical and biomedical analysis¹⁶. Dissociation constants can be a key parameter for understanding and quantifying chemical phenomena such as reaction rates, biological activity, biological uptake, biological transport, acid-base titration, solvent extraction, complex formation ion transport and environmental fate¹⁷⁻¹⁹.

For water insoluble compounds, the formation constant and other thermodynamic properties are measured in purely non-aqueous organic solvent or in a mixture of two solvents, one of which may be water. A solvent mixture containing water and water miscible organic solvent is known as mixed aqueous medium. A number of works has been done in non aqueous and mixed aqueous media^{20, 21}.

Some of the thermodynamic parameters have also been evaluated with the help of dissociation constant²²⁻²⁵. Fazary and Taha²⁶ have been reported the thermodynamics of second-stage dissociation of propenoic acid derivative in different solvent mixture. Seleem et al have been reported the dissociation constant and thermodynamic parameters of Schiff bases²⁷. Verma was reported the thermodynamic assessment of water from dissociation constant²⁸. Das and co-workers²⁹ have been reported the thermodynamics of the dissociation constants of trans-cinnamic acid. Topallar and Bayrak³⁰ have been reported the thermodynamics of dissociation constants of cromium soap.

A literature survey shows that various workers studied the dissociation constant of a number of substances. Many researchers have been reported the dissociation constant of organic compounds, amino acids, vitamins, polymeric materials, drug molecules, metal complexes³¹⁻⁴³.

The importance of pyrimidine derivatives arises from their biological, medicinal and agricultural applications⁴⁴. Uracil and Thiamine have biological importance in metabolism because some of their derivatives are building blocks for RNA and DNA⁴⁵. 5-Fluorouracil is a drug which is available for colon, rectum and breast tumors⁴⁶. These advantages of pyrimidines derivatives cause an interest to study the dissociation constants and thermodynamic parameters of these derivatives. Seleem et al have been reported the dissociation constants of Schiff bases of pyrimidine moiety²⁷.

In the present work, the dissociation constant of all synthesized dihydropyrimidinones derivatives (RVG series) are studied in DMF at different temperatures (298.15- 318.15 K) by Calvin Bjerrum pH titration technique.

EXPERIMENTAL

All solutions used for the titration are prepared using distilled water. Following are the concentrations of the solutions used for the titration. The chemicals used were of B.D.H Analar grade.

| Solutions | Concentration (M) |
|------------------|-------------------|
| Nitric acid | 1.0 |
| Sodium hydroxide | 0.5 |
| Sodium nitrate | 1.0 |
| Ligand (in DMF) | 0.1 |

Nitric acid and sodium hydroxide were standardized by titrating with 0.1 N NaOH and 0.05 M succinic acid solution respectively.

The buffer solutions used for the calibration of pH meter were 0.05 M potassium hydrogen phthalate and 0.01 M Borax buffer.

A Systronics *pH* meter (Model No. EQ 664) was used for the pH determination. The Systronic glass electrode and a saturated calomel electrode were used as indicator and reference electrodes respectively. Before operation, the glass electrode was immersed in 0.1 M HCl for twenty minutes. Then, it was washed thoroughly with distilled water. The *pH* meter was calibrated with buffer solution of known pH. A constant temperature was maintained to ± 0.05 K by using a digital controller Nova Inst. (Model No. NV 8550 E).

Calvin Bjerrum pH titration :

The following sets of mixtures were prepared for titration:


- (I) 2 ml HNO_3 (1.0M) + 4 ml water + 30 ml DMF + 4.0 ml NaNO_3 (1.0 M).
- (ii) 2 ml HNO_3 (1.0M) + 4 ml water + 28 ml DMF + 2.0 ml ligand solution (0.1M) + 4.0 ml NaNO_3 (1.0 M).

Thus, total volumes (V^0) in each set = 40.0 ml and Solvent : Water ratio 60:40 (v/v). The solvent used for the study was DMF.


The above mentioned solutions were allowed to attain a definite temperature and then titrated against standard NaOH solution (0.5 M). The same experiment was repeated at different temperatures, i.e., at 289.15 K, 308.15K and 318.15K.

THEORY

In the present work, all the compounds or ligands are of HL type. Thus, the equilibria are,



In general, these equations can be represented as:



The thermodynamic proton-ligand stability constant (TK_j^H) is given by:

$$TK_j^H = \frac{[LH_j]}{\{[LH_{j-1}][H]\}} \quad \dots (3.6.1)$$


TK_j^H is reciprocal of the thermodynamic dissociation constant of the acid LH_j dissociating as:



The overall thermodynamic proton-ligand stability constant β_j^H is given by:

$$T\beta_j^H = \frac{[LH_j]}{[L][H]^j} \quad \dots (3.6.2)$$

and it refers to the reaction:



The stoichiometric proton-ligand stability constant is given by:

$$K_j^H = \frac{[LH_j]}{\{[LH_{j-1}][H]\}} \quad \dots (3.6.3)$$

and

$$\beta_j^H = \frac{[LH_j]}{[L][H]^j} \quad \dots (3.6.4)$$

An inert electrolyte is used to determine the stability constant in a particular salt medium. Sodium nitrate is mostly preferred as supporting electrolyte, because of very slight complexing tendency of nitrate ion. Generally, the competition between nitrate ion and the ligand under study is minor importance. The molar concentrations are used in place of activities.

For the determination of dissociation constants, Bjerrum⁴⁷ introduced a relation for the determination of \bar{n}_H , which is defined as average number of hydrogen bound to each ligand.

$$\bar{n}_H = \frac{\{K_1^H [H] + 2K_1^H K_2^H [H]^2 + \dots JK_1^H K_2^H [H] \dots K_j^H [H]^j\}}{\{1 + K_1^H [H] + K_1^H K_2^H [H]^2 \dots K_1^H K_2^H \dots K_j^H [H]^j\}} \quad \dots (3.6.5)$$

From equation (3.6.4), we can write

$$\bar{n}_H = \frac{\sum_{j=1}^{\eta} j \beta_j^H [H]^j}{\sum_{j=1}^{\eta} \beta_j^H [H]^j} \quad : (\beta_0^H = 1) \quad \dots (3.6.6)$$

Equation (3.4.6) is called Bjerrum formation function of the system.

The determination of dissociation constants from experimental data comprises the following three steps: (i) evaluation of formation curve of the system (ii) calculation of stoichiometric K's of the system by direct solution of the formation function and (iii) conversion of stoichiometric constants into thermodynamic constants.

When the system consists of a ligand, which is a conjugated base of a weak acid, the pH-metric method introduced by Bjerrum has been widely used. This method is known as "Bjerrum-Calvin pH titration technique".

In this technique, the concentration of H^+ ions is measured. Thus, a large amount of data can be obtained in a short period of time. The Irving and Rossotti method⁴² has some advantages, such as:

- (i) Valid for both pure water and for the mixed solvents.
- (ii) Conversion of pH-meter reading in to stoichiometric hydrogen ion concentration is not necessary.
- (iii) Not necessary to know the stoichiometric concentration of neutral salt added to maintain the ionic strength constant.

Due to these advantages, this method is used in the present work. The pH-meter is standardized using an aqueous buffer. The $pH(B)$ is measured for two solutions:

- (1) A mixture containing a mineral acid, a chelating agent and a neutral electrolyte to keep ionic strength constant and

(2) A mixture same as above but without the chelating agent, when titrated against an alkali solution.

After each addition of standard alkali, the *pH* meter reading (*B*) is noted using a glass electrode-saturated calomel electrode combination. For both the titrations, same initial volume of the mixture and same standard alkali is used. The titration curves obtained in the above two titrations are designated as the reagent or ligand titration curve and the acid titration curve respectively.

The possible hydrolysis reactions are ignored because (i) fresh reagent solutions were used in *pH* titrations, (ii) titration times were of the order of one hour, (iii) there were no observable drifts with time in the meter readings and (iv) the concentrations of the mineral acid or alkali in the solutions were small.

Usually, a *pH*-meter calibrated with an aqueous buffer is used for aqueous solutions only. However, for the mixed aqueous media, especially aqueous dioxane solutions, van Uitert and Haas gave a relation between the glass electrode reading *B* in dioxane-water medium and the stoichiometric hydrogen ion concentration of the same in mixture of varied composition and ionic strength. They reported the relation:

$$-\log [H^+] = B + \log f + \log U_H^0 \quad \dots (3.6.7)$$

where *f* is the activity coefficient of the hydrogen ions in the solvent mixture under consideration at the same temperature and ionic strength, and U_H^0 is a correction factor at zero ionic strength, which depends only on the solvent composition and temperature. U_H^0 is taken as unity in aqueous media. The meter reading in any aqueous dioxane solution can, therefore, be converted into hydrogen ion concentration using equation (3.6.7), provided that correction factor for the appropriate solvent, salt medium, and temperature, has been determined.

Equation (3.6.7) can be written as:

$$1/\text{anti log } B = [H^+] f U_H^0 \quad \dots (3.6.8)$$

$$\therefore [H^+] = \frac{1}{[\text{anti log } B] [f U_H^0]} \quad \dots (3.6.9)$$

Substituting for $[H^+]$ in equation (3.4.5) we get,

$$\bar{n}_H = (K_1^H / f U_H^0) [1/\text{antilog } B] + \dots + ((JK_1^H K_2^H \dots K_J^H) / (f U_H^0)^J) [1/\text{antilog } B]^J$$

$$/(1+K_1^H/f U_H^0)[1/\text{antilog } B]+..+ ((K_1^H K_2^H \dots K_J^H)/(f U_H^0)^J)[1/\text{antilog } B] \dots (3.610)$$

$$K_j^H = fU_H^0 \cdot pK_j^H \dots (3.6.11)$$

$$\beta_j^H = fU_H^0 \cdot p\beta_j^H \dots (3.6.12)$$

The proton-ligands constant, pK_j^H can be obtained by the interpolation at half \bar{n}_H values. The pH value at $\bar{n}_H = 0.5$ gives pK_1 .

$$\log K_1 = \left(\bar{n}_H \right)_{0.5} \dots (3.6.13)$$

RESULTS AND DISCUSSION

Figure 3.6.1 shows that typical titrations curve of the acid in the absence and presence of compound RVG-1 at [A] 298.15K, [B] 308.15K and [C] 318.15K. It can be seen that for the same volume of NaOH added the compound titration curves showed a lower pH value than the titration curve of free acid.

From these titration curves, the average number of protons associated with the ligand ($\overline{n_H}$) can be calculated by the following equation given by Irving and Rossotti⁴⁸.

$$\overline{n_H} = Y - \left\{ (V'' - V') (N^0 + E^0) \right\} / \left\{ (V^0 + V') T_L^0 \right\} \quad \dots (3.6.14)$$

where Y is the number of displaceable protons per ligand molecule. For all compounds, Y is taken as one. V' and V'' are the volume of alkali required at the same pH for both acid and ligand titration curves respectively. V⁰ is the initial volume of the test solution.

N⁰, E⁰ and T_L⁰ are the initial concentration of the alkali, acid and ligand respectively.

The calculated values of $\overline{n_H}$ for all the studied compounds are given in Tables 3.6.1 to 3.6.3 for all temperature systems. It is evident from Tables that the values of $\overline{n_H}$ are found to be between 0 to 1 for all the compounds. suggesting thereby the presence of only one replaceable proton.

Figure 3.6.2 shows the plot of $\overline{n_H}$ values against pH for RVG-1. The pK_1^H value was evaluated at $\overline{n_H} = 0.5$ i.e., by half-integral method for all the compounds with Y=1.

Further, the $\log \overline{n_H} / (\overline{n_H} - 1)$ values are plotted against B as shown in Figure 3.6.3. The plots are straight lines from which $\log pK_1^H$ values were calculated at several B values, by the following equation:

$$\log pK_1^H = \text{pH} + \log \left[\overline{n_H} / (\overline{n_H} - 1) \right] \quad \dots (3.6.15)$$

The calculated pK_1^H values are reported in Tables 3.6.1 to 3.6.3 for all compounds at different pH values. From these $\log pK_1^H$ values, the average

values of pK_1^H was calculated and are reported in Table 3.6.4 along with the pK_1 value evaluated by half-integral method. Further, it is evident from Table 3.6.4 that these pK_1^H values are in agreement with that obtained by half-integral method. The acidity constant decreases in the order: $p\text{-OCH}_3 > o\text{-OCH}_3 > p\text{-CH}_3 > 2,5\text{-di Cl} > p\text{-Cl} > p\text{-F} > o\text{-OCH}_3 > m\text{-Cl} > 3,4\text{-di Cl} > 3\text{-Cl}, 4\text{-F}$.

Further, the pK_1^H value decreases with increasing temperature, i.e. the acidity of the compounds increases.⁴⁹ Further, pK_1^H is minimum in RVG-10 and maximum in RVG-1 suggesting thereby maximum dissociation in RVG-10 which contains fluoride and chloride groups. RVG-1 contains $-\text{OCH}_3$ group which causes a decrease in dissociation. Thus, high stability of compound RVG-1 can be attributed to the presence of the $-\text{OCH}_3$ group in the para position relative to the amide group. The presence of para- OCH_3 group (i. e., an electron donating effect) enhances the electron density by their high positive inductive or mesomeric effect thereby increasing the stability of the compound. RVG-1 is followed by RVG-5 which also contains $-\text{OCH}_3$ group in ortho position. RVG-2 and RVG-4 are also stable which contain methyl group at para and ortho positions respectively. Similar behavior was also observed by El-Sonbati et al.⁵⁰ while studying dissociation constants of nitrogen containing heterocyclic aldehydes. It was observed that $-\text{OCH}_3$ and $-\text{CH}_3$ groups at para position increases the stability of compounds.

The higher acidity of RVG-10 may be due to the para substitution of fluoride group which is most electronegative halogen. Further, next electronegative halogen chloride is also present at meta-position. These substitutions cause negative inductive effect which may decrease the stability of this compound. Thus, presence of different substituents influences the dissociation of the compound.

Further, some thermodynamic parameters such as enthalpy of solution, Gibb's energy change and entropy of solution have also been evaluated from dissociation constants at different temperatures for these systems.

The enthalpy changes (ΔH) for the dissociation process were evaluated⁵¹ from the slope of the plot pK_1^H vs. $1/T$.

Using the following equation, Gibbs energy changes (ΔG) have been evaluated.⁵¹

$$\Delta G = 2.303 RT \log K_1^H \quad \dots (3.6.16)$$

From these ΔG and ΔH values, entropy values (ΔS) were calculated using the following relationship⁵¹, assuming ΔH to be independent of temperature⁵²

$$\Delta S = (\Delta H - \Delta G) / T \quad \dots (3.6.17)$$

All these thermodynamic parameters are reported in Table 3.6.5 for both average and half-integral methods. It is observed that for both methods, values are in good agreement with each other.

The positive value of ΔH indicates that dissociation process is endothermic and is accompanied by absorption of heat and favourable at higher temperature.^{50, 52} Further, ΔG values are positive indicating thereby that the dissociation process is not spontaneous.^{50, 53} However, negative value of ΔS is due to the increased order as a result of the solvation processes⁵⁰ except RVG-7.

Figure 3.6.1: The plot of pH (B) against volume of NaOH for RVG-1 in DMF at [A] 298.15K [B] 308.15K and [C] 318.15K.




Table 3.6.1: The pH , n_H , pK_1^H and other terms for RVG series in DMF at 298.15K.

| pH | V' | V'' | $V''-V'$ | n_H | $\log n_H/(1-n_H)$ | pK_1^H |
|--------------|--------|--------|----------|--------|--------------------|----------|
| RVG-1 | | | | | | |
| 10.2 | 4.1981 | 4.2191 | 0.0210 | 0.9478 | -0.4398 | 11.6887 |
| 10.3 | 4.2035 | 4.2357 | 0.0321 | 0.9200 | -0.4460 | 11.4587 |
| 10.4 | 4.2105 | 4.2581 | 0.0476 | 0.8816 | -0.4527 | 11.3606 |
| 10.5 | 4.2176 | 4.2778 | 0.0601 | 0.8505 | -0.4622 | 11.2721 |
| 10.6 | 4.2246 | 4.2967 | 0.0721 | 0.8209 | -0.4704 | 11.2551 |
| 10.7 | 4.2316 | 4.3215 | 0.0899 | 0.7769 | -0.4786 | 11.6887 |
| 10.8 | 4.2387 | 4.3389 | 0.1002 | 0.7514 | -0.4914 | 11.4587 |
| 10.9 | 4.2457 | 4.3655 | 0.1198 | 0.7030 | -0.4993 | 11.2741 |
| 11 | 4.2528 | 4.3886 | 0.1358 | 0.6635 | -0.5153 | 11.2948 |
| 11.1 | 4.2598 | 4.4108 | 0.1510 | 0.6260 | -0.5295 | 11.3237 |
| 11.2 | 4.2669 | 4.4357 | 0.1688 | 0.5821 | -0.5439 | 11.3440 |
| 11.3 | 4.2739 | 4.4564 | 0.1825 | 0.5484 | -0.5623 | 11.3844 |
| 11.4 | 4.2809 | 4.4827 | 0.2018 | 0.5010 | -0.5777 | 11.4017 |
| 11.5 | 4.2880 | 4.5044 | 0.2164 | 0.4651 | -0.6015 | 11.4393 |
| 11.6 | 4.2950 | 4.5254 | 0.2304 | 0.4308 | -0.6215 | 11.4790 |
| 11.7 | 4.3032 | 4.5485 | 0.2453 | 0.3943 | -0.6425 | 11.5136 |
| 11.8 | 4.3142 | 4.5743 | 0.2601 | 0.3581 | -0.6673 | 11.5466 |
| 11.9 | 4.3252 | 4.5994 | 0.2742 | 0.3237 | -0.6948 | 11.5800 |
| RVG-2 | | | | | | |
| 10 | 4.1907 | 4.3011 | 0.1104 | 0.7252 | -0.5078 | 10.4214 |
| 10.1 | 4.1944 | 4.3139 | 0.1195 | 0.7026 | -0.5155 | 10.4733 |
| 10.2 | 4.1981 | 4.3248 | 0.1267 | 0.6847 | -0.5217 | 10.5367 |
| 10.3 | 4.2035 | 4.3364 | 0.1329 | 0.6693 | -0.5273 | 10.6061 |
| 10.4 | 4.2105 | 4.3594 | 0.1489 | 0.6295 | -0.5425 | 10.6302 |
| 10.5 | 4.2176 | 4.3819 | 0.1643 | 0.5913 | -0.5583 | 10.6604 |
| 10.6 | 4.2246 | 4.3998 | 0.1752 | 0.5642 | -0.5704 | 10.7122 |
| 10.7 | 4.2316 | 4.4181 | 0.1865 | 0.5362 | -0.5836 | 10.7630 |
| 10.8 | 4.2387 | 4.4292 | 0.1905 | 0.5263 | -0.5885 | 10.8458 |
| 10.9 | 4.2457 | 4.4471 | 0.2014 | 0.4993 | -0.6024 | 10.8988 |
| 11 | 4.2528 | 4.4697 | 0.2169 | 0.4608 | -0.6240 | 10.9318 |
| 11.1 | 4.2598 | 4.4914 | 0.2316 | 0.4244 | -0.6467 | 10.9677 |
| 11.2 | 4.2669 | 4.5167 | 0.2498 | 0.3793 | -0.6783 | 10.9860 |
| 11.3 | 4.2739 | 4.5303 | 0.2564 | 0.3630 | -0.6909 | 11.0557 |
| 11.4 | 4.2809 | 4.5455 | 0.2646 | 0.3427 | -0.7076 | 11.1171 |
| 11.5 | 4.288 | 4.5619 | 0.2739 | 0.3197 | -0.7280 | 11.1720 |
| RVG-3 | | | | | | |
| 10.2 | 4.3361 | 4.4955 | 0.1594 | 0.6045 | -0.5527 | 10.3843 |
| 10.3 | 4.3574 | 4.5226 | 0.1652 | 0.5903 | -0.5588 | 10.4587 |
| 10.4 | 4.3787 | 4.5508 | 0.1721 | 0.5734 | -0.5662 | 10.5285 |
| 10.5 | 4.3905 | 4.5691 | 0.1786 | 0.5574 | -0.5735 | 10.6002 |
| 10.6 | 4.4093 | 4.5936 | 0.1843 | 0.5435 | -0.5801 | 10.6758 |
| 10.7 | 4.4186 | 4.6091 | 0.1905 | 0.5282 | -0.5875 | 10.7491 |
| 10.8 | 4.428 | 4.6306 | 0.2026 | 0.4984 | -0.6029 | 10.7972 |
| 10.9 | 4.4373 | 4.6457 | 0.2084 | 0.4841 | -0.6107 | 10.8724 |
| 11 | 4.4467 | 4.6636 | 0.2169 | 0.4632 | -0.6226 | 10.9360 |
| 11.1 | 4.456 | 4.6805 | 0.2245 | 0.4445 | -0.6339 | 11.0032 |
| 11.2 | 4.4654 | 4.6991 | 0.2337 | 0.4219 | -0.6483 | 11.0631 |
| 11.3 | 4.4747 | 4.7206 | 0.2459 | 0.3918 | -0.6691 | 11.1090 |
| 11.4 | 4.4841 | 4.7357 | 0.2516 | 0.3778 | -0.6794 | 11.1834 |

| pH | V' | V'' | $V''-V'$ | n_H | $\log n_H/(1-n_H)$ | pK_1^H |
|--------------|--------|--------|----------|--------|--------------------|----------|
| RVG-4 | | | | | | |
| 9.9 | 4.2879 | 4.4466 | 0.1587 | 0.6058 | -0.5522 | 10.0867 |
| 10 | 4.2972 | 4.463 | 0.1658 | 0.5883 | -0.5596 | 10.1550 |
| 10.1 | 4.3148 | 4.4872 | 0.1724 | 0.5721 | -0.5668 | 10.2261 |
| 10.2 | 4.3361 | 4.5163 | 0.1802 | 0.5529 | -0.5756 | 10.2923 |
| 10.3 | 4.3574 | 4.5453 | 0.1879 | 0.5340 | -0.5847 | 10.3592 |
| 10.4 | 4.3787 | 4.5743 | 0.1956 | 0.5152 | -0.5941 | 10.4264 |
| 10.5 | 4.3905 | 4.5926 | 0.2021 | 0.4992 | -0.6025 | 10.4986 |
| 10.6 | 4.4093 | 4.6249 | 0.2156 | 0.4660 | -0.6210 | 10.5408 |
| 10.7 | 4.4186 | 4.6451 | 0.2265 | 0.4391 | -0.6373 | 10.5937 |
| 10.8 | 4.428 | 4.6629 | 0.2349 | 0.4184 | -0.6506 | 10.6570 |
| 10.9 | 4.4373 | 4.6824 | 0.2451 | 0.3933 | -0.6680 | 10.7117 |
| 11 | 4.4467 | 4.7035 | 0.2568 | 0.3645 | -0.6897 | 10.7585 |
| RVG-5 | | | | | | |
| 10.6 | 4.4093 | 4.5784 | 0.1649 | 0.5915 | -0.5582 | 10.7609 |
| 10.7 | 4.4186 | 4.5938 | 0.1702 | 0.5785 | -0.5639 | 10.8375 |
| 10.8 | 4.428 | 4.6085 | 0.1786 | 0.5578 | -0.5733 | 10.9009 |
| 10.9 | 4.4373 | 4.6249 | 0.1876 | 0.5356 | -0.5839 | 10.9620 |
| 11 | 4.4467 | 4.6409 | 0.1942 | 0.5194 | -0.5920 | 11.0337 |
| 11.1 | 4.456 | 4.6597 | 0.2037 | 0.4960 | -0.6042 | 11.0930 |
| 11.2 | 4.4654 | 4.6803 | 0.2149 | 0.4684 | -0.6196 | 11.1450 |
| 11.3 | 4.4747 | 4.6976 | 0.2229 | 0.4487 | -0.6313 | 11.2106 |
| 11.4 | 4.4841 | 4.7219 | 0.2378 | 0.4120 | -0.6550 | 11.2455 |
| 11.5 | 4.4934 | 4.7405 | 0.2471 | 0.3891 | -0.6710 | 11.3041 |
| 11.6 | 4.5044 | 4.7608 | 0.2564 | 0.3663 | -0.6883 | 11.3619 |
| RVG-6 | | | | | | |
| 10 | 4.1972 | 4.3496 | 0.1524 | 0.6207 | -0.5461 | 10.2139 |
| 10.1 | 4.2149 | 4.3759 | 0.161 | 0.5995 | -0.5549 | 10.2751 |
| 10.2 | 4.2362 | 4.4056 | 0.1694 | 0.5788 | -0.5638 | 10.3380 |
| 10.3 | 4.2574 | 4.4333 | 0.1759 | 0.5628 | -0.5710 | 10.4097 |
| 10.4 | 4.2787 | 4.4609 | 0.1822 | 0.5474 | -0.5782 | 10.4825 |
| 10.5 | 4.2891 | 4.488 | 0.1989 | 0.5060 | -0.5989 | 10.5104 |
| 10.6 | 4.3093 | 4.515 | 0.2057 | 0.4893 | -0.6078 | 10.5815 |
| 10.7 | 4.3187 | 4.5312 | 0.2125 | 0.4726 | -0.6172 | 10.6523 |
| 10.8 | 4.328 | 4.5514 | 0.2234 | 0.4456 | -0.6332 | 10.7052 |
| 10.9 | 4.3374 | 4.5686 | 0.2312 | 0.4264 | -0.6454 | 10.7712 |
| 11 | 4.3467 | 4.5866 | 0.2399 | 0.4049 | -0.6598 | 10.8328 |
| RVG-7 | | | | | | |
| 10.8 | 4.328 | 4.4784 | 0.1504 | 0.6268 | -0.5436 | 11.0252 |
| 10.9 | 4.3374 | 4.4972 | 0.1598 | 0.6035 | -0.5531 | 11.0825 |
| 11 | 4.3467 | 4.5114 | 0.1647 | 0.5915 | -0.5583 | 11.1607 |
| 11.1 | 4.3561 | 4.5329 | 0.1768 | 0.5615 | -0.5716 | 11.2075 |
| 11.2 | 4.3654 | 4.5499 | 0.1845 | 0.5425 | -0.5805 | 11.2741 |
| 11.3 | 4.3748 | 4.5670 | 0.1922 | 0.5236 | -0.5899 | 11.3410 |
| 11.4 | 4.3841 | 4.5858 | 0.2017 | 0.5001 | -0.6020 | 11.4002 |
| 11.5 | 4.3935 | 4.6069 | 0.2134 | 0.4712 | -0.6180 | 11.4500 |
| 11.6 | 4.4045 | 4.6294 | 0.2249 | 0.4429 | -0.6349 | 11.5003 |
| 11.7 | 4.4194 | 4.6552 | 0.2358 | 0.4161 | -0.6522 | 11.5528 |
| 11.8 | 4.4343 | 4.6810 | 0.2467 | 0.3893 | -0.6709 | 11.6044 |
| 11.9 | 4.4493 | 4.7036 | 0.2543 | 0.3707 | -0.6849 | 11.6701 |
| 12 | 4.4642 | 4.7329 | 0.2687 | 0.3353 | -0.7140 | 11.7027 |

| pH | V' | V'' | $V''-V'$ | n_H | $\log n_H/(1-n_H)$ | pK_1^H |
|---------------|--------|--------|----------|--------|--------------------|----------|
| RVG-8 | | | | | | |
| 10.4 | 4.2787 | 4.4408 | 0.1621 | 0.5973 | -0.5558 | 10.5712 |
| 10.5 | 4.2891 | 4.4591 | 0.17 | 0.5778 | -0.5643 | 10.6362 |
| 10.6 | 4.3093 | 4.4888 | 0.1795 | 0.5544 | -0.5749 | 10.6948 |
| 10.7 | 4.3187 | 4.5065 | 0.1878 | 0.5339 | -0.5847 | 10.7589 |
| 10.8 | 4.328 | 4.5221 | 0.1941 | 0.5183 | -0.5925 | 10.8319 |
| 10.9 | 4.3374 | 4.5384 | 0.201 | 0.5013 | -0.6014 | 10.9023 |
| 11 | 4.3467 | 4.5623 | 0.2156 | 0.4652 | -0.6215 | 10.9395 |
| 11.1 | 4.3561 | 4.5801 | 0.2239 | 0.4447 | -0.6337 | 11.0036 |
| 11.2 | 4.3654 | 4.5966 | 0.2312 | 0.4268 | -0.6451 | 11.0718 |
| 11.3 | 4.3748 | 4.6144 | 0.2396 | 0.4061 | -0.6590 | 11.1348 |
| 11.4 | 4.3841 | 4.6297 | 0.2456 | 0.3913 | -0.6694 | 11.2081 |
| RVG-9 | | | | | | |
| 9.6 | 3.516 | 3.6611 | 0.1451 | 0.6332 | -0.5410 | 9.8371 |
| 9.7 | 3.536 | 3.6881 | 0.1521 | 0.6157 | -0.5481 | 9.9047 |
| 9.8 | 3.556 | 3.7141 | 0.1581 | 0.6007 | -0.5543 | 9.9774 |
| 9.9 | 3.576 | 3.7429 | 0.1669 | 0.5787 | -0.5638 | 10.0378 |
| 10 | 3.596 | 3.7749 | 0.1789 | 0.5486 | -0.5776 | 10.0847 |
| 10.1 | 3.6276 | 3.8170 | 0.1894 | 0.5225 | -0.5904 | 10.1390 |
| 10.2 | 3.6621 | 3.8594 | 0.1973 | 0.5029 | -0.6005 | 10.2051 |
| 10.3 | 3.6966 | 3.9055 | 0.2089 | 0.4741 | -0.6163 | 10.2550 |
| 10.4 | 3.722 | 3.9376 | 0.2156 | 0.4576 | -0.6260 | 10.3261 |
| 10.5 | 3.7463 | 3.9704 | 0.2241 | 0.4365 | -0.6389 | 10.3891 |
| 10.6 | 3.7707 | 4.0056 | 0.2349 | 0.4097 | -0.6565 | 10.4413 |
| 10.7 | 3.7951 | 4.0402 | 0.2451 | 0.3844 | -0.6745 | 10.4955 |
| 10.8 | 3.8195 | 4.0762 | 0.2567 | 0.3556 | -0.6968 | 10.5418 |
| RVG-10 | | | | | | |
| 9.3 | 3.4569 | 3.6061 | 0.1492 | 0.6223 | -0.5454 | 9.5169 |
| 9.4 | 3.4765 | 3.6336 | 0.1571 | 0.6025 | -0.5536 | 9.5807 |
| 9.5 | 3.4961 | 3.6637 | 0.1676 | 0.5761 | -0.5650 | 9.6333 |
| 9.6 | 3.5160 | 3.6902 | 0.1742 | 0.5597 | -0.5725 | 9.7041 |
| 9.7 | 3.5362 | 3.7162 | 0.1802 | 0.5447 | -0.5795 | 9.7779 |
| 9.8 | 3.5561 | 3.7442 | 0.1882 | 0.5247 | -0.5893 | 9.8429 |
| 9.9 | 3.5760 | 3.7745 | 0.1985 | 0.4989 | -0.6026 | 9.8981 |
| 10 | 3.5961 | 3.8014 | 0.2054 | 0.4817 | -0.6120 | 9.9683 |
| 10.1 | 3.6276 | 3.8388 | 0.2112 | 0.4675 | -0.6201 | 10.0434 |
| 10.2 | 3.6621 | 3.8816 | 0.2195 | 0.4470 | -0.6324 | 10.1076 |
| 10.3 | 3.6966 | 3.9244 | 0.2278 | 0.4265 | -0.6453 | 10.1715 |
| 10.4 | 3.7220 | 3.9584 | 0.2364 | 0.4052 | -0.6596 | 10.2334 |
| 10.5 | 3.7463 | 3.9887 | 0.2424 | 0.3905 | -0.6700 | 10.3066 |

Table 3.6.2: The pH , n_H , pK_1^H and other terms for RVG series in DMF at 308.15 K.

| pH | V' | V'' | $V''-V'$ | n_H | $\log n_H/(1-n_H)$ | pK_1^H |
|--------------|--------|--------|----------|--------|--------------------|----------|
| RVG-1 | | | | | | |
| 10.2 | 4.3895 | 4.4758 | 0.0863 | 0.7866 | -0.4885 | 10.7665 |
| 10.3 | 4.4146 | 4.5148 | 0.1002 | 0.7524 | -0.4990 | 10.7827 |
| 10.4 | 4.439 | 4.5511 | 0.1121 | 0.7232 | -0.5085 | 10.8171 |
| 10.5 | 4.4634 | 4.5853 | 0.1219 | 0.6993 | -0.5166 | 10.8664 |
| 10.6 | 4.4878 | 4.6287 | 0.1409 | 0.6527 | -0.5335 | 10.8740 |
| 10.7 | 4.5063 | 4.6545 | 0.1482 | 0.6349 | -0.5404 | 10.9404 |
| 10.8 | 4.519 | 4.6738 | 0.1548 | 0.6188 | -0.5468 | 11.0105 |
| 10.9 | 4.5316 | 4.6969 | 0.1653 | 0.5932 | -0.5575 | 11.0638 |
| 11 | 4.5443 | 4.7188 | 0.1745 | 0.5708 | -0.5674 | 11.1238 |
| 11.1 | 4.557 | 4.7435 | 0.1865 | 0.5415 | -0.5810 | 11.1723 |
| 11.2 | 4.5696 | 4.7715 | 0.2019 | 0.5039 | -0.6000 | 11.2069 |
| 11.3 | 4.5823 | 4.7959 | 0.2136 | 0.4755 | -0.6155 | 11.2574 |
| 11.4 | 4.5949 | 4.8305 | 0.2356 | 0.4219 | -0.6483 | 11.2632 |
| 11.5 | 4.6057 | 4.8546 | 0.2489 | 0.3896 | -0.6707 | 11.3050 |
| 11.6 | 4.6152 | 4.8716 | 0.2564 | 0.3715 | -0.6843 | 11.3716 |
| 11.7 | 4.6248 | 4.8935 | 0.2687 | 0.3416 | -0.7085 | 11.4151 |
| 11.8 | 4.6343 | 4.9057 | 0.2714 | 0.3352 | -0.7141 | 11.5026 |
| 11.9 | 4.6438 | 4.9302 | 0.2864 | 0.2988 | -0.7482 | 11.5296 |
| RVG-2 | | | | | | |
| 10 | 4.3368 | 4.4642 | 0.1274 | 0.6839 | -0.5220 | 10.3352 |
| 10.1 | 4.3632 | 4.4991 | 0.1359 | 0.6630 | -0.5296 | 10.3939 |
| 10.2 | 4.3895 | 4.5386 | 0.1491 | 0.6305 | -0.5421 | 10.4321 |
| 10.3 | 4.4146 | 4.5718 | 0.1572 | 0.6107 | -0.5502 | 10.4955 |
| 10.4 | 4.439 | 4.6033 | 0.1643 | 0.5933 | -0.5575 | 10.5640 |
| 10.5 | 4.4634 | 4.6387 | 0.1753 | 0.5663 | -0.5694 | 10.6159 |
| 10.6 | 4.4878 | 4.6734 | 0.1856 | 0.5411 | -0.5812 | 10.6715 |
| 10.7 | 4.5063 | 4.7028 | 0.1965 | 0.5143 | -0.5946 | 10.7249 |
| 10.8 | 4.519 | 4.7199 | 0.2009 | 0.5036 | -0.6001 | 10.8063 |
| 10.9 | 4.5316 | 4.7425 | 0.2109 | 0.4790 | -0.6135 | 10.8636 |
| 11 | 4.5443 | 4.7656 | 0.2213 | 0.4535 | -0.6284 | 10.9190 |
| 11.1 | 4.557 | 4.7947 | 0.2377 | 0.4132 | -0.6541 | 10.9476 |
| 11.2 | 4.5696 | 4.8168 | 0.2472 | 0.3899 | -0.6705 | 11.0055 |
| 11.3 | 4.5823 | 4.8335 | 0.2512 | 0.3802 | -0.6776 | 11.0878 |
| 11.4 | 4.5949 | 4.8545 | 0.2596 | 0.3597 | -0.6936 | 11.1495 |
| 11.5 | 4.6057 | 4.8705 | 0.2648 | 0.3470 | -0.7039 | 11.2254 |
| RVG-3 | | | | | | |
| 10.1 | 4.4317 | 4.5776 | 0.1459 | 0.6388 | -0.5389 | 10.3476 |
| 10.2 | 4.4561 | 4.6078 | 0.1517 | 0.6246 | -0.5445 | 10.4212 |
| 10.3 | 4.4805 | 4.6403 | 0.1598 | 0.6048 | -0.5526 | 10.4848 |
| 10.4 | 4.5038 | 4.6722 | 0.1684 | 0.5838 | -0.5616 | 10.5469 |
| 10.5 | 4.5226 | 4.6965 | 0.1739 | 0.5704 | -0.5676 | 10.6230 |
| 10.6 | 4.5415 | 4.7271 | 0.1856 | 0.5416 | -0.5810 | 10.6725 |
| 10.7 | 4.5604 | 4.7602 | 0.1998 | 0.5068 | -0.5985 | 10.7118 |
| 10.8 | 4.5792 | 4.7906 | 0.2114 | 0.4784 | -0.6139 | 10.7624 |
| 10.9 | 4.5981 | 4.8213 | 0.2232 | 0.4495 | -0.6308 | 10.8119 |
| 11 | 4.6069 | 4.8384 | 0.2315 | 0.4291 | -0.6436 | 10.8760 |
| 11.1 | 4.6145 | 4.8543 | 0.2398 | 0.4088 | -0.6571 | 10.9397 |
| 11.2 | 4.6221 | 4.8679 | 0.2458 | 0.3941 | -0.6674 | 11.0131 |
| 11.3 | 4.6298 | 4.8827 | 0.2529 | 0.3767 | -0.6803 | 11.0812 |
| 11.4 | 4.6374 | 4.8992 | 0.2618 | 0.3548 | -0.6974 | 11.1404 |

| pH | V' | V'' | $V''-V'$ | n_H | $\log n_H/(1-n_H)$ | pK_1^H |
|--------------|--------|--------|----------|--------|--------------------|----------|
| RVG-4 | | | | | | |
| 9.8 | 4.366 | 4.5181 | 0.1521 | 0.6229 | -0.5452 | 10.0179 |
| 9.9 | 4.386 | 4.5441 | 0.1581 | 0.6082 | -0.5512 | 10.0910 |
| 10 | 4.4073 | 4.5685 | 0.1612 | 0.6007 | -0.5543 | 10.1774 |
| 10.1 | 4.4317 | 4.6012 | 0.1695 | 0.5804 | -0.5631 | 10.2408 |
| 10.2 | 4.4561 | 4.6317 | 0.1756 | 0.5655 | -0.5698 | 10.3144 |
| 10.3 | 4.4805 | 4.6684 | 0.1879 | 0.5353 | -0.5840 | 10.3615 |
| 10.4 | 4.5038 | 4.7086 | 0.2048 | 0.4938 | -0.6054 | 10.3892 |
| 10.5 | 4.5226 | 4.7404 | 0.2178 | 0.4619 | -0.6234 | 10.4337 |
| 10.6 | 4.5415 | 4.7651 | 0.2236 | 0.4478 | -0.6319 | 10.5090 |
| 10.7 | 4.5604 | 4.7918 | 0.2314 | 0.4288 | -0.6438 | 10.5754 |
| 10.8 | 4.5792 | 4.8178 | 0.2386 | 0.4113 | -0.6554 | 10.6442 |
| 10.9 | 4.5981 | 4.8416 | 0.2435 | 0.3994 | -0.6636 | 10.7228 |
| 11 | 4.6069 | 4.8558 | 0.2489 | 0.3862 | -0.6732 | 10.7988 |
| RVG-5 | | | | | | |
| 10.4 | 4.5038 | 4.7053 | 0.1529 | 0.6221 | -0.5455 | 10.6164 |
| 10.5 | 4.5226 | 4.7241 | 0.1596 | 0.6057 | -0.5522 | 10.6864 |
| 10.6 | 4.5415 | 4.743 | 0.1677 | 0.5858 | -0.5607 | 10.7506 |
| 10.7 | 4.5604 | 4.7619 | 0.1754 | 0.5670 | -0.5691 | 10.8171 |
| 10.8 | 4.5792 | 4.7807 | 0.1811 | 0.5531 | -0.5755 | 10.8927 |
| 10.9 | 4.5981 | 4.7996 | 0.1894 | 0.5329 | -0.5852 | 10.9571 |
| 11 | 4.6069 | 4.8084 | 0.2015 | 0.5031 | -0.6004 | 11.0054 |
| 11.1 | 4.6145 | 4.8267 | 0.2122 | 0.4768 | -0.6148 | 11.0597 |
| 11.2 | 4.6221 | 4.8452 | 0.2231 | 0.4500 | -0.6305 | 11.1129 |
| 11.3 | 4.6298 | 4.8596 | 0.2298 | 0.4336 | -0.6407 | 11.1840 |
| 11.4 | 4.6374 | 4.8741 | 0.2367 | 0.4167 | -0.6518 | 11.2539 |
| 11.5 | 4.645 | 4.8899 | 0.2449 | 0.3966 | -0.6656 | 11.3177 |
| 11.6 | 4.6527 | 4.9056 | 0.2529 | 0.3770 | -0.6800 | 11.3818 |
| RVG-6 | | | | | | |
| 9.8 | 3.8353 | 3.9895 | 0.1542 | 0.6131 | -0.5492 | 9.9998 |
| 9.9 | 3.8471 | 4.0092 | 0.1621 | 0.5933 | -0.5575 | 10.0641 |
| 10 | 3.8588 | 4.0289 | 0.1701 | 0.5734 | -0.5662 | 10.1284 |
| 10.1 | 3.8706 | 4.0492 | 0.1786 | 0.5522 | -0.5759 | 10.1910 |
| 10.2 | 3.8824 | 4.0688 | 0.1864 | 0.5328 | -0.5853 | 10.2570 |
| 10.3 | 3.8941 | 4.0866 | 0.1925 | 0.5176 | -0.5929 | 10.3306 |
| 10.4 | 3.9147 | 4.1143 | 0.1996 | 0.5000 | -0.6020 | 10.4001 |
| 10.5 | 3.9441 | 4.1554 | 0.2113 | 0.4711 | -0.6181 | 10.4497 |
| 10.6 | 3.9735 | 4.195 | 0.2215 | 0.4459 | -0.6330 | 10.5057 |
| 10.7 | 4.0015 | 4.2313 | 0.2298 | 0.4255 | -0.6459 | 10.5696 |
| 10.8 | 4.0162 | 4.2543 | 0.2381 | 0.4050 | -0.6598 | 10.6329 |
| 10.9 | 4.0309 | 4.2774 | 0.2465 | 0.3842 | -0.6747 | 10.6951 |
| 11 | 4.0456 | 4.3032 | 0.2576 | 0.3567 | -0.6960 | 10.7438 |
| RVG-7 | | | | | | |
| 9.7 | 3.8235 | 3.9814 | 0.1579 | 0.6037 | -0.5531 | 9.8827 |
| 9.8 | 3.8353 | 4.0035 | 0.1682 | 0.5779 | -0.5642 | 9.9365 |
| 9.9 | 3.8471 | 4.0213 | 0.1742 | 0.5630 | -0.5709 | 10.0100 |
| 10 | 3.8588 | 4.0389 | 0.1801 | 0.5483 | -0.5778 | 10.0842 |
| 10.1 | 3.8706 | 4.0601 | 0.1895 | 0.5249 | -0.5892 | 10.1432 |
| 10.2 | 3.8824 | 4.079 | 0.1966 | 0.5072 | -0.5983 | 10.2125 |
| 10.3 | 3.8941 | 4.0955 | 0.2014 | 0.4953 | -0.6046 | 10.2918 |
| 10.4 | 3.9147 | 4.1273 | 0.2126 | 0.4675 | -0.6201 | 10.3434 |
| 10.5 | 3.9441 | 4.1676 | 0.2235 | 0.4405 | -0.6364 | 10.3962 |
| 10.6 | 3.9735 | 4.2076 | 0.2341 | 0.4144 | -0.6533 | 10.4498 |
| 10.7 | 4.0015 | 4.2427 | 0.2412 | 0.3970 | -0.6653 | 10.5185 |
| 10.8 | 4.0162 | 4.2654 | 0.2492 | 0.3772 | -0.6799 | 10.5823 |

| | | | | | | |
|---------------|--------|--------|--------|--------|---------|---------|
| 10.9 | 4.0309 | 4.2873 | 0.2564 | 0.3594 | -0.6937 | 10.6491 |
| RVG-8 | | | | | | |
| 10.2 | 3.8824 | 4.0293 | 0.1469 | 0.6318 | -0.5416 | 10.4344 |
| 10.3 | 3.8941 | 4.0483 | 0.1542 | 0.6136 | -0.5490 | 10.5008 |
| 10.4 | 3.9147 | 4.0768 | 0.1621 | 0.5940 | -0.5572 | 10.5652 |
| 10.5 | 3.9441 | 4.1139 | 0.1698 | 0.5750 | -0.5655 | 10.6312 |
| 10.6 | 3.9735 | 4.1611 | 0.1876 | 0.5307 | -0.5863 | 10.6534 |
| 10.7 | 4.0015 | 4.1983 | 0.1968 | 0.5080 | -0.5978 | 10.7139 |
| 10.8 | 4.0162 | 4.2196 | 0.2034 | 0.4917 | -0.6065 | 10.7856 |
| 10.9 | 4.0309 | 4.2444 | 0.2135 | 0.4666 | -0.6206 | 10.8419 |
| 11 | 4.0456 | 4.2717 | 0.2261 | 0.4353 | -0.6396 | 10.8870 |
| 11.1 | 4.0603 | 4.2998 | 0.2395 | 0.4021 | -0.6618 | 10.9277 |
| 11.2 | 4.075 | 4.3214 | 0.2464 | 0.3850 | -0.6740 | 10.9967 |
| 11.3 | 4.0897 | 4.3485 | 0.2588 | 0.3543 | -0.6979 | 11.0394 |
| 11.4 | 4.1061 | 4.3695 | 0.2634 | 0.3431 | -0.7072 | 11.1179 |
| RVG-9 | | | | | | |
| 9.4 | 3.7796 | 3.9097 | 0.1301 | 0.6731 | -0.5259 | 9.7137 |
| 9.5 | 3.8101 | 3.9389 | 0.1389 | 0.6512 | -0.5341 | 9.7711 |
| 9.6 | 3.8122 | 3.962 | 0.1498 | 0.6239 | -0.5448 | 9.8198 |
| 9.7 | 3.8244 | 3.9902 | 0.1658 | 0.5838 | -0.5616 | 9.8470 |
| 9.8 | 3.8366 | 4.0111 | 0.1745 | 0.5621 | -0.5713 | 9.9085 |
| 9.9 | 3.8488 | 4.0344 | 0.1856 | 0.5344 | -0.5845 | 9.9599 |
| 10 | 3.8610 | 4.0599 | 0.1989 | 0.5012 | -0.6014 | 10.0020 |
| 10.1 | 3.8732 | 4.0921 | 0.2189 | 0.4512 | -0.6298 | 10.0149 |
| 10.2 | 3.8854 | 4.1112 | 0.2258 | 0.4340 | -0.6405 | 10.0847 |
| 10.3 | 3.8976 | 4.1367 | 0.2391 | 0.4009 | -0.6626 | 10.1255 |
| 10.4 | 3.9242 | 4.1754 | 0.2512 | 0.3709 | -0.6847 | 10.1706 |
| 10.5 | 3.9545 | 4.2169 | 0.2624 | 0.3433 | -0.7070 | 10.2183 |
| 10.6 | 3.9848 | 4.2593 | 0.2745 | 0.3135 | -0.7338 | 10.2596 |
| RVG-10 | | | | | | |
| 9.1 | 3.7184 | 3.8596 | 0.1412 | 0.6447 | -0.5366 | 9.3588 |
| 9.2 | 3.7388 | 3.8895 | 0.1507 | 0.6210 | -0.5459 | 9.4145 |
| 9.3 | 3.7592 | 3.9185 | 0.1593 | 0.5996 | -0.5548 | 9.4753 |
| 9.4 | 3.7796 | 3.9469 | 0.1673 | 0.5796 | -0.5634 | 9.5395 |
| 9.5 | 3.8104 | 3.9752 | 0.1752 | 0.5600 | -0.5723 | 9.6047 |
| 9.6 | 3.8122 | 3.9983 | 0.1861 | 0.5328 | -0.5853 | 9.6570 |
| 9.7 | 3.8244 | 4.0237 | 0.1993 | 0.4998 | -0.6022 | 9.6996 |
| 9.8 | 3.8366 | 4.0425 | 0.2059 | 0.4833 | -0.6111 | 9.7710 |
| 9.9 | 3.8488 | 4.0652 | 0.2164 | 0.4571 | -0.6262 | 9.8254 |
| 10 | 3.8610 | 4.0849 | 0.2239 | 0.4385 | -0.6376 | 9.8926 |
| 10.1 | 3.8732 | 4.1119 | 0.2387 | 0.4015 | -0.6622 | 9.9267 |
| 10.2 | 3.8854 | 4.1311 | 0.2457 | 0.3841 | -0.6747 | 9.9950 |
| 10.3 | 3.8976 | 4.1543 | 0.2567 | 0.3568 | -0.6959 | 10.0440 |

Table 3.6.3: The pH , n_H , pK_1^H and other terms for RVG series in DMF at 318.15 K.

| pH | V' | V'' | $V''-V'$ | n_H | $\log n_H/(1-n_H)$ | pK_1^H |
|--------------|---------|--------|----------|--------|--------------------|----------|
| RVG-1 | | | | | | |
| 10.5 | 3.9125 | 4.0314 | 0.1189 | 0.7030 | -0.5153 | 10.8741 |
| 10.6 | 3.9250 | 4.0561 | 0.1311 | 0.6727 | -0.5261 | 10.9128 |
| 10.7 | 3.9375 | 4.0870 | 0.1495 | 0.6270 | -0.5435 | 10.9255 |
| 10.8 | 3.95041 | 4.1102 | 0.1602 | 0.6005 | -0.5544 | 10.9770 |
| 10.9 | 3.9625 | 4.1389 | 0.1764 | 0.5604 | -0.5721 | 11.0054 |
| 11 | 3.9750 | 4.1644 | 0.1894 | 0.5283 | -0.5875 | 11.0491 |
| 11.1 | 3.9875 | 4.1899 | 0.2024 | 0.4962 | -0.6041 | 11.0934 |
| 11.2 | 4.0000 | 4.2198 | 0.2198 | 0.4532 | -0.6286 | 11.1185 |
| 11.3 | 4.0172 | 4.2445 | 0.2273 | 0.4349 | -0.6399 | 11.1862 |
| 11.4 | 4.0345 | 4.2743 | 0.2398 | 0.4042 | -0.6603 | 11.2315 |
| 11.5 | 4.0517 | 4.3062 | 0.2545 | 0.3681 | -0.6868 | 11.2654 |
| 11.6 | 4.0690 | 4.3384 | 0.2694 | 0.3316 | -0.7172 | 11.2957 |
| 11.7 | 4.0862 | 4.3674 | 0.2812 | 0.3028 | -0.7442 | 11.3378 |
| RVG-2 | | | | | | |
| 10 | 3.8535 | 3.9941 | 0.1405 | 0.6476 | -0.5355 | 10.2642 |
| 10.1 | 3.8651 | 4.0147 | 0.1496 | 0.6248 | -0.5444 | 10.3216 |
| 10.2 | 3.8767 | 4.0361 | 0.1594 | 0.6004 | -0.5545 | 10.3768 |
| 10.3 | 3.8884 | 4.0541 | 0.1657 | 0.5847 | -0.5612 | 10.4486 |
| 10.4 | 3.9012 | 4.0784 | 0.1784 | 0.5530 | -0.5756 | 10.4924 |
| 10.5 | 3.9125 | 4.0981 | 0.1856 | 0.5351 | -0.5841 | 10.5610 |
| 10.6 | 3.925 | 4.1236 | 0.1986 | 0.5027 | -0.6007 | 10.6046 |
| 10.7 | 3.9375 | 4.1431 | 0.2056 | 0.4853 | -0.6101 | 10.6744 |
| 10.8 | 3.9510 | 4.1687 | 0.2187 | 0.4526 | -0.6289 | 10.7175 |
| 10.9 | 3.9625 | 4.1861 | 0.2236 | 0.4405 | -0.6364 | 10.7962 |
| 11 | 3.9750 | 4.2104 | 0.2354 | 0.4112 | -0.6555 | 10.8440 |
| 11.1 | 3.9875 | 4.2351 | 0.2476 | 0.3808 | -0.6772 | 10.8889 |
| 11.2 | 4.0012 | 4.2541 | 0.2541 | 0.3647 | -0.6895 | 10.9591 |
| RVG-3 | | | | | | |
| 9.8 | 3.8462 | 3.9948 | 0.1486 | 0.6272 | -0.5434 | 10.0259 |
| 9.9 | 3.8718 | 4.0261 | 0.1543 | 0.6131 | -0.5491 | 10.1000 |
| 10 | 3.8974 | 4.0582 | 0.1608 | 0.5971 | -0.5559 | 10.1708 |
| 10.1 | 3.929 | 4.0976 | 0.1686 | 0.5778 | -0.5642 | 10.2363 |
| 10.2 | 3.9613 | 4.1405 | 0.1792 | 0.5516 | -0.5762 | 10.2900 |
| 10.3 | 3.9935 | 4.1791 | 0.1856 | 0.5359 | -0.5837 | 10.3625 |
| 10.4 | 4.0286 | 4.2213 | 0.1927 | 0.5186 | -0.5924 | 10.4323 |
| 10.5 | 4.0643 | 4.2654 | 0.2011 | 0.4980 | -0.6031 | 10.4965 |
| 10.6 | 4.1102 | 4.3154 | 0.2154 | 0.4627 | -0.6229 | 10.5351 |
| 10.7 | 4.4238 | 4.6516 | 0.2278 | 0.4359 | -0.6393 | 10.5881 |
| 10.8 | 4.1476 | 4.3833 | 0.2357 | 0.4127 | -0.6544 | 10.6468 |
| 10.9 | 4.1714 | 4.4129 | 0.2415 | 0.3986 | -0.6642 | 10.7214 |
| 11 | 4.1952 | 4.4493 | 0.2541 | 0.3676 | -0.6873 | 10.7643 |
| RVG-4 | | | | | | |
| 9.6 | 3.7967 | 3.9434 | 0.1467 | 0.6315 | -0.5417 | 9.8340 |
| 9.7 | 3.8205 | 3.9729 | 0.1524 | 0.6174 | -0.5474 | 9.9079 |
| 9.8 | 3.8462 | 4.0064 | 0.1602 | 0.5981 | -0.5554 | 9.9726 |
| 9.9 | 3.8718 | 4.0398 | 0.168 | 0.5788 | -0.5638 | 10.0380 |
| 10 | 3.8974 | 4.0756 | 0.1782 | 0.5535 | -0.5753 | 10.0932 |
| 10.1 | 3.929 | 4.1144 | 0.1854 | 0.5358 | -0.5838 | 10.1622 |
| 10.2 | 3.9613 | 4.1596 | 0.1983 | 0.5038 | -0.6000 | 10.2066 |
| 10.3 | 3.9935 | 4.2041 | 0.2106 | 0.4734 | -0.6167 | 10.2538 |

| pH | V' | V'' | $V''-V'$ | n_H | $\log n_H/(1-n_H)$ | pK_1^H |
|--------------|--------|--------|----------|--------|--------------------|----------|
| 10.4 | 4.0286 | 4.252 | 0.2234 | 0.4419 | -0.6355 | 10.2985 |
| 10.5 | 4.0643 | 4.2961 | 0.2318 | 0.4213 | -0.6487 | 10.3622 |
| 10.6 | 4.1101 | 4.3431 | 0.2431 | 0.3936 | -0.6678 | 10.4123 |
| 10.7 | 4.4238 | 4.6802 | 0.2564 | 0.3651 | -0.6892 | 10.4597 |
| 10.8 | 4.1476 | 4.4085 | 0.2609 | 0.3499 | -0.7015 | 10.5310 |
| RVG-5 | | | | | | |
| 10.2 | 3.9613 | 4.108 | 0.1467 | 0.6329 | -0.5412 | 10.4366 |
| 10.3 | 3.9935 | 4.1481 | 0.1546 | 0.6134 | -0.5490 | 10.5006 |
| 10.4 | 4.0286 | 4.1909 | 0.1623 | 0.5945 | -0.5570 | 10.5662 |
| 10.5 | 4.0643 | 4.2424 | 0.1781 | 0.5554 | -0.5744 | 10.5966 |
| 10.6 | 4.1012 | 4.2842 | 0.1842 | 0.5405 | -0.5815 | 10.6706 |
| 10.7 | 4.4238 | 4.6194 | 0.1956 | 0.5157 | -0.5939 | 10.7272 |
| 10.8 | 4.1476 | 4.1476 | 0.2018 | 0.4972 | -0.6036 | 10.7951 |
| 10.9 | 4.1714 | 4.3863 | 0.2149 | 0.4648 | -0.6217 | 10.8388 |
| 11 | 4.1952 | 4.4188 | 0.2236 | 0.4435 | -0.6345 | 10.9014 |
| 11.1 | 4.2258 | 4.4605 | 0.2347 | 0.4162 | -0.6521 | 10.9531 |
| 11.2 | 4.2581 | 4.5039 | 0.2458 | 0.3891 | -0.6711 | 11.0041 |
| 11.3 | 4.2903 | 4.5492 | 0.2589 | 0.3570 | -0.6957 | 11.0444 |
| 11.4 | 4.3146 | 4.5778 | 0.2632 | 0.3467 | -0.7042 | 11.1248 |
| RVG-6 | | | | | | |
| 9.7 | 3.8186 | 3.9698 | 0.1512 | 0.6204 | -0.5462 | 9.9134 |
| 9.8 | 3.8302 | 3.9897 | 0.1595 | 0.5997 | -0.5547 | 9.9756 |
| 9.9 | 3.8419 | 4.0103 | 0.1684 | 0.5775 | -0.5644 | 10.0357 |
| 10 | 3.8535 | 4.0283 | 0.1748 | 0.5615 | -0.5716 | 10.1075 |
| 10.1 | 3.8651 | 4.0507 | 0.1856 | 0.5346 | -0.5844 | 10.1602 |
| 10.2 | 3.8767 | 4.0679 | 0.1912 | 0.5207 | -0.5913 | 10.2359 |
| 10.3 | 3.8884 | 4.0907 | 0.2023 | 0.4930 | -0.6058 | 10.2878 |
| 10.4 | 3.9100 | 4.1135 | 0.2135 | 0.4650 | -0.6216 | 10.3392 |
| 10.5 | 3.9125 | 4.1336 | 0.2211 | 0.4461 | -0.6329 | 10.4061 |
| 10.6 | 3.9250 | 4.1546 | 0.2296 | 0.4250 | -0.6463 | 10.4688 |
| 10.7 | 3.9375 | 4.172 | 0.2345 | 0.4129 | -0.6543 | 10.5472 |
| 10.8 | 3.9510 | 4.1916 | 0.2416 | 0.3953 | -0.6666 | 10.6154 |
| 10.9 | 3.9625 | 4.2123 | 0.2498 | 0.3750 | -0.6816 | 10.6781 |
| RVG-7 | | | | | | |
| 9.5 | 3.7946 | 3.9514 | 0.1568 | 0.6062 | -0.5520 | 9.6873 |
| 9.6 | 3.8072 | 3.9717 | 0.1647 | 0.5864 | -0.5604 | 9.7517 |
| 9.7 | 3.8186 | 3.9905 | 0.1719 | 0.5685 | -0.5684 | 9.8197 |
| 9.8 | 3.8302 | 4.0104 | 0.1802 | 0.5478 | -0.5780 | 9.8832 |
| 9.9 | 3.8419 | 4.0293 | 0.1874 | 0.5298 | -0.5867 | 9.9518 |
| 10 | 3.8535 | 4.0448 | 0.1913 | 0.5202 | -0.5916 | 10.0350 |
| 10.1 | 3.8651 | 4.0616 | 0.1965 | 0.5072 | -0.5982 | 10.1126 |
| 10.2 | 3.8767 | 4.0831 | 0.2064 | 0.4825 | -0.6116 | 10.1697 |
| 10.3 | 3.8884 | 4.106 | 0.2176 | 0.4546 | -0.6277 | 10.2209 |
| 10.4 | 3.9210 | 4.1234 | 0.2234 | 0.4402 | -0.6365 | 10.2957 |
| 10.5 | 3.9125 | 4.1443 | 0.2318 | 0.4193 | -0.6500 | 10.3587 |
| 10.6 | 3.9250 | 4.1706 | 0.2456 | 0.3850 | -0.6741 | 10.3965 |
| 10.7 | 3.9375 | 4.1936 | 0.2561 | 0.3588 | -0.6942 | 10.4479 |
| RVG-8 | | | | | | |
| 10 | 3.8535 | 4.0056 | 0.1521 | 0.6185 | -0.5470 | 10.2098 |
| 10.1 | 3.8651 | 4.0234 | 0.1583 | 0.6030 | -0.5533 | 10.2816 |
| 10.2 | 3.8767 | 4.0456 | 0.1689 | 0.5766 | -0.5648 | 10.3341 |
| 10.3 | 3.8884 | 4.0671 | 0.1787 | 0.5521 | -0.5760 | 10.3909 |
| 10.4 | 3.9010 | 4.0869 | 0.1869 | 0.5317 | -0.5858 | 10.4551 |
| 10.5 | 3.9125 | 4.105 | 0.1925 | 0.5178 | -0.5928 | 10.5309 |
| 10.6 | 3.9252 | 4.1241 | 0.1991 | 0.5014 | -0.6013 | 10.6024 |
| 10.7 | 3.9375 | 4.1511 | 0.2136 | 0.4652 | -0.6214 | 10.6395 |

| pH | V' | V'' | V''-V' | n_H | log n_H/(1-n_H) | pK₁^H |
|---------------|-----------|------------|---------------|----------------------|--|-----------------------------------|
| 10.8 | 3.9530 | 4.1798 | 0.2298 | 0.4248 | -0.6464 | 10.6684 |
| 10.9 | 3.9625 | 4.1979 | 0.2354 | 0.4110 | -0.6556 | 10.7437 |
| 11 | 3.9751 | 4.2217 | 0.2467 | 0.3829 | -0.6756 | 10.7927 |
| 11.1 | 3.9875 | 4.2396 | 0.2521 | 0.3696 | -0.6857 | 10.8681 |
| RVG-9 | | | | | | |
| 9.2 | 3.7541 | 3.8962 | 0.1421 | 0.6428 | -0.5373 | 9.4551 |
| 9.3 | 3.7676 | 3.9162 | 0.1486 | 0.6265 | -0.5437 | 9.5247 |
| 9.4 | 3.7811 | 3.9359 | 0.1548 | 0.6111 | -0.5500 | 9.5962 |
| 9.5 | 3.7946 | 3.963 | 0.1684 | 0.5770 | -0.5646 | 9.6349 |
| 9.6 | 3.8074 | 3.9824 | 0.1754 | 0.5596 | -0.5725 | 9.7040 |
| 9.7 | 3.8186 | 4.0120 | 0.1814 | 0.5446 | -0.5795 | 9.7777 |
| 9.8 | 3.8302 | 4.0291 | 0.1989 | 0.5008 | -0.6016 | 9.8014 |
| 9.9 | 3.8419 | 4.0506 | 0.2087 | 0.4764 | -0.6150 | 9.8589 |
| 10 | 3.8535 | 4.0703 | 0.2168 | 0.4562 | -0.6268 | 9.9237 |
| 10.1 | 3.8651 | 4.0889 | 0.2238 | 0.4388 | -0.6375 | 9.9931 |
| 10.2 | 3.8767 | 4.1064 | 0.2297 | 0.4241 | -0.6468 | 10.0672 |
| 10.3 | 3.8884 | 4.1252 | 0.2368 | 0.4065 | -0.6587 | 10.1356 |
| 10.4 | 3.9102 | 4.1457 | 0.2457 | 0.3844 | -0.6745 | 10.1954 |
| RVG-10 | | | | | | |
| 9 | 3.7270 | 3.8615 | 0.1345 | 0.6617 | -0.5301 | 9.2913 |
| 9.1 | 3.7405 | 3.8797 | 0.1392 | 0.6499 | -0.5346 | 9.3687 |
| 9.2 | 3.7541 | 3.9028 | 0.1487 | 0.6262 | -0.5439 | 9.4240 |
| 9.3 | 3.7676 | 3.9224 | 0.1548 | 0.6109 | -0.5500 | 9.4960 |
| 9.4 | 3.7811 | 3.9449 | 0.1638 | 0.5885 | -0.5596 | 9.5553 |
| 9.5 | 3.7946 | 3.9741 | 0.1795 | 0.5491 | -0.5774 | 9.5857 |
| 9.6 | 3.8071 | 4.0051 | 0.1981 | 0.5026 | -0.6007 | 9.6045 |
| 9.7 | 3.8186 | 4.0231 | 0.2045 | 0.4866 | -0.6093 | 9.6768 |
| 9.8 | 3.8302 | 4.0470 | 0.2168 | 0.4559 | -0.6270 | 9.7232 |
| 9.9 | 3.8419 | 4.0663 | 0.2244 | 0.4370 | -0.6386 | 9.7899 |
| 10 | 3.8535 | 4.0854 | 0.2319 | 0.4183 | -0.6507 | 9.8568 |
| 10.1 | 3.8651 | 4.1044 | 0.2393 | 0.3999 | -0.6633 | 9.9237 |
| 10.2 | 3.8767 | 4.1225 | 0.2458 | 0.3838 | -0.6750 | 9.9943 |

Figure 3.6.2: The plot of n_H against pH (B) for [A] RVG-1 and [B] RVG-2 in DMF at different temperatures.





Figure 3.6.3: The plot of $\log n_H/(1-n_H)$ against B for [A] RVG-1 and [B] RVG-2 in DMF at different temperatures.





Table 3.6.4: The pK_1^H values for all the studied compounds calculated by different methods in DMF at different temperatures.

| <i>Comp. code</i> | <i>T/K</i> | <i>Half-integral method</i> | <i>Average method</i> | <i>Comp. code</i> | <i>T/K</i> | <i>Half-integral method</i> | <i>Average method</i> |
|-------------------|------------|-----------------------------|-----------------------|-------------------|------------|-----------------------------|-----------------------|
| | | pK_1^H | | | | pK_1^H | |
| RVG-1 | 298.15 | 11.44 | 11.41 | RVG-6 | 298.15 | 10.54 | 10.52 |
| | 308.15 | 11.20 | 11.19 | | 308.15 | 10.32 | 10.38 |
| | 318.15 | 11.08 | 11.09 | | 318.15 | 10.26 | 10.29 |
| RVG-2 | 298.15 | 10.82 | 10.87 | RVG-7 | 298.15 | 11.32 | 11.35 |
| | 308.15 | 10.80 | 10.79 | | 308.15 | 10.20 | 10.26 |
| | 318.15 | 10.62 | 10.61 | | 318.15 | 9.98 | 10.08 |
| RVG-3 | 298.15 | 10.76 | 10.79 | RVG-8 | 298.15 | 10.82 | 10.88 |
| | 308.15 | 10.70 | 10.71 | | 308.15 | 10.72 | 10.77 |
| | 318.15 | 10.40 | 10.41 | | 318.15 | 10.48 | 10.57 |
| RVG-4 | 298.15 | 10.44 | 10.44 | RVG-9 | 298.15 | 10.16 | 10.20 |
| | 308.15 | 10.34 | 10.40 | | 308.15 | 9.94 | 9.99 |
| | 318.15 | 10.16 | 10.19 | | 318.15 | 9.82 | 9.82 |
| RVG-5 | 298.15 | 11.0 | 11.07 | RVG-10 | 298.15 | 9.86 | 9.90 |
| | 308.15 | 10.92 | 11.0 | | 308.15 | 9.68 | 9.70 |
| | 318.15 | 10.74 | 10.78 | | 318.15 | 9.66 | 9.63 |

Table 3.6.5: Thermodynamic parameters for the dissociation of RVG series in DMF-Water mixture at different temperatures for half integral and average methods.

| compounds | T/K | ΔG_{Hal} <i>kJ mol⁻¹</i> | ΔG_{Ave} <i>kJ mol⁻¹</i> | ΔH_{Hal} <i>kJ mol⁻¹</i> | ΔH_{Ave} <i>kJ mol⁻¹</i> | ΔS_{Hal} <i>J mol⁻¹ K⁻¹</i> | ΔS_{Ave} <i>J mol⁻¹ K⁻¹</i> |
|------------------|------------|---|---|---|---|---|---|
| RVG-1 | 298.15 | 63.34 | 63.17 | 32.79 | 29.16 | -0.1056 | -0.1176 |
| | 308.15 | 66.08 | 66.02 | | | -0.1080 | -0.1196 |
| | 318.15 | 67.49 | 67.55 | | | -0.1091 | -0.1207 |
| RVG-2 | 298.15 | 59.90 | 60.18 | 17.99 | 23.50 | -0.1449 | -0.1268 |
| | 308.15 | 63.72 | 63.66 | | | -0.1484 | -0.1303 |
| | 318.15 | 64.69 | 64.63 | | | -0.1468 | -0.1293 |
| RVG-3 | 298.15 | 59.57 | 59.73 | 32.44 | 34.28 | -0.0938 | -0.0880 |
| | 308.15 | 63.13 | 63.19 | | | -0.0996 | -0.0938 |
| | 318.15 | 63.35 | 63.41 | | | -0.0972 | -0.0916 |
| RVG-4 | 298.15 | 57.80 | 57.80 | 25.33 | 25.33 | -0.1123 | -0.1220 |
| | 308.15 | 61.00 | 61.36 | | | -0.1158 | -0.1260 |
| | 318.15 | 61.89 | 62.07 | | | -0.1149 | -0.1243 |
| RVG-5 | 298.15 | 60.90 | 61.28 | 23.50 | 22.52 | -0.1293 | -0.1214 |
| | 308.15 | 64.43 | 64.90 | | | -0.1328 | -0.1257 |
| | 318.15 | 65.42 | 65.66 | | | -0.1318 | -0.1241 |
| RVG-6 | 298.15 | 58.35 | 58.24 | 25.42 | 20.92 | -0.1139 | -0.1291 |
| | 308.15 | 60.89 | 61.24 | | | -0.1151 | -0.1308 |
| | 318.15 | 62.50 | 62.68 | | | -0.1165 | -0.1312 |
| RVG-7 | 298.15 | 62.67 | 62.83 | 122.52 | 116.18 | 0.2070 | 0.1845 |
| | 308.15 | 60.18 | 60.53 | | | 0.2023 | 0.1806 |
| | 318.15 | 60.79 | 61.40 | | | 0.1940 | 0.1722 |
| RVG-8 | 298.15 | 59.90 | 60.23 | 30.72 | 28.05 | -0.1009 | -0.1113 |
| | 308.15 | 63.25 | 63.54 | | | -0.1055 | -0.1152 |
| | 318.15 | 63.84 | 64.38 | | | -0.1041 | -0.1142 |
| RVG-9 | 298.15 | 56.25 | 56.47 | 30.96 | 34.53 | -0.0875 | -0.0759 |
| | 308.15 | 58.64 | 58.94 | | | -0.0898 | -0.0792 |
| | 318.15 | 59.82 | 59.82 | | | -0.0907 | -0.0795 |
| RVG-10 | 298.15 | 54.58 | 54.81 | 18.31 | 24.63 | -0.1255 | -0.1043 |
| | 308.15 | 57.11 | 57.23 | | | -0.1259 | -0.1058 |
| | 318.15 | 58.84 | 58.66 | | | -0.1274 | -0.1069 |

*Hal: Half integral method

*Avg: Average method.

LITERATURE CITED

- (1) Ghasemi, J., Ahmadi, S., Kubista, M., Forootan, A. "Determination of Acidity Constants of 4-(2-Pyridylazo)resorcinol in Binary Acetonitrile + Water Mixtures." *J. Chem. Eng. Data*, **2003**, 48, 1178-1182.
- (2) Zhao, Y. H., Yuan, L. H., Wang, L. S. "Quantative Structure-Activity Relationships of organic acids and bases." *Bull Environ. Contam. Toxicol.*, **1996**, 57, 242-249.
- (3) Rochester, H. "Acidity Functions: Academic Press." *New York*, **1971**.
- (4) Hammett, L. P. "Physical Organic Chemistry" *McGraw-Hill: New York*, **1940**.
- (5) Beltran, J. L., Codony, R. M., Granados, A., Izquierdo, M. D. "Study on partition equilibria of metal complexes in non-ionic micellar solutions from spectrophotometric data." *Talanta*, **2000**, 52 (2), 225-232.
- (6) Afkhami, A., Khalafi, L. "Spectrophotometric determination of conditional acidity constant as a function of β -cyclodextrin concentration for some organic acids using Rank annihilation factor analysis." *Anal. Chim. Acta.*, **2006**, 569, 267-274.
- (7) Gerakis, A. M., Koupparis, M. A., Efstathiou, C. E. "Micellar acid-base potentiometric titrations of weak acidic and/or insoluble drugs." *J. Pharm. Biomed. Anal.*, **1993**, 11, 33-41.
- (8) Niazi, M. S. K. "Conductometric determination of thermodynamic dissociation constants of benzoic and nitrobenzoic acid in methanol-water mixtures at 298 K." *Bull. Chem. Soc. Jap.*, **1989**, 62, 1253-1257.
- (9) Kapoor, R. C., Agrawal, O. P., Seth, T. D. "Polarographic determination of the dissociation constant of the sulphhydryl group and diffusion coefficient measurements of thiomalic acid." *Can. J. Chem.*, **1961**, 39, 2236-2239.
- (10) Samik, N., Datta, D. "Determination of acid dissociation constants of some monobasic organic acids in acetonitrile from molar conductance." *Ind. J. Chem.*, **2007**, 46, 1263-1265.
- (11) Lahiri, S., Das, A., Ghosh, K., Banerjee, A. Nayak, D. "Determination of dynamic dissociation constants of chromium-poly(N-vinylpyrrolidone) complex by the radiotracer technique." *Ind. J. Chem.*, **2009**, 48, 672-675.
- (12) Sager, E. E. Siewers, I. J. "Dissociation Constants of 4-Aminobenzophenone Calculated From Ultraviolet Spectra at Several Temperatures." *J. Res. National Bureau of Standards.*, **1950**, 45(6), 489-496.

- (13) Strong, L. E., Kinney, T., Fischer, P. "Ionization of aqueous benzoic acid: conductance and thermodynamics." *J. Sol. Chem.*, **1979**, 8, 329-345.
- (14) Pal, A., Lahiri, S. C. "Conductometric studies on the dissociation constants of benzoic acid in 2-propanol + water mixtures." *Zeitschrift fuer Physikalische Chemie.*, **1987**, 268, 378-384.
- (15) Galic, N., Cimerman, Z., Tomisic, V. "Spectrometric study of tautomeric and protonation equilibria of o-vanillin Schiff base derivatives and their complexes with Cu(II)." *Spectrochimica Acta, A: Mole. Biomole. Spect.*, **2009**, 71A, 1274-1280.
- (16) Fan, J., Shen, X, Wang, J. "Dissociation constants of methyl orange in aqueous alcohol solvents." *Anal. Chim. Acta.*, **1998**, 364, 275-280.
- (17) Gluck, S. J., Steele, K. P., Benko, M. H. "Determination of acidity constants of monoprotic and diprotic acids by capillary electrophoresis." *J. Chromatogr. A.*, **1996**, 745, 117-125.
- (18) Zhao, Y. H., Yuan, L. H., Wang, L. S. "Quantative structure activity relationships of organic acids and bases." *Bull. Environ. Contam. Toxicol.*, **1996**, 57, 242-249.
- (19) Khalafi., L., Rohani, M., Afkhami, A. "Acidity constants of some organic acids in the presence of β -cyclodextrin in binary ethanol-water mixtures by Rank Annihilation Factor Analysis." *J. Chem. Eng. Data*, **2008**, 53, 2389-2392.
- (20) Azimi, G., Khoobi, A., Zamani, K., Zolgharnein, J. "Multiwavelength spectrophotometric determination of acidity constants of newly synthesized 1,2,4-triazole derivatives in ethanol-water mixtures." *J. Chem. Eng. Data*, **2008**, 53, 1862-1866.
- (21) Martins, C. T., Sato, B. M., El-Seoud, O. A. "First Study on the Thermo-Solvatochromism in Aqueous 1-(1-Butyl)-3-methylimidazolium Tetrafluoro borate: A Comparison between the Solvation by an Ionic Liquid and by Aqueous Alcohols." *J. Phy. Chem. B*, **2008**, 112, 8330-8339.
- (22) Sengupta, M., Pal, K., Chakravarti, A., Mahapatra, P. "Dissociation constants of toluic acids in aqueous solution at different temperatures and study of related thermodynamic parameters." *J. Chem. Eng. Data*, **1978**, 23, 103-107.
- (23) Meloun, M., Syrový, T., Vrana, A. "The thermodynamic dissociation constants of ambroxol, antazoline, naphazoline, oxymetazoline and ranitidine by the regression analysis of spectrophotometric data." *Talanta*, **2004**, 62, 511-522.

- (24) Hamborg, S. E., Versteeg, G. F. "Dissociation constants and thermodynamic properties of Amines and Alkanolamines from (293 to 353) K." *J. Chem. Eng. Data*, **2009**, 54, 1318-1328.
- (25) Bari, I., Hafiz, A., Khanzada, A. W. K. "Determination of dissociation constants and thermodynamic parameters of 8-aminoquinoline by potentiometry in aqueous and mixed organic-water solvent systems." *J. Chem. Soc. Pak.*, **2000**, 22, 1-5.
- (26) Taha, M., Fazary, A. E. "Thermodynamics of the second-stage dissociation of 2-[N-(2-hydroxyethyl)-N-methylaminomethyl]-propenoic acid in water at different ionic strength and different solvent mixtures." *J. Chem. Thermod.*, **2005**, 37, 43-48.
- (27) Seleem, H. S., El-Shetary, B. A., Khalil, S. M. E., Shebl, M. "Potentiometric and spectrophotometric studies of the complexation of Schiff-base hydrazones containing the pyrimidine moiety." *J. Serb. Chem. Soc.*, **2003**, 68(10), 729-748.
- (28) Verma, M. P. "A thermodynamic assessment of dissociation constant of water." Proceedings, *Twenty-Eighth Workshop on Geothermal Reservoir Engineering., Stanford University, Stanford, California*, **2003**, 27-29,
- (29) Das, R. C., Dash, U. N., Panda, K. N. "Thermodynamics of the dissociation of trans-cinnamic acid." *Can. J. Chem.*, **1976**, 54, 1916.
- (30) Topallar, H., Bayrak, Y. "Thermodynamics of the Dissociation of Chromium Soap Solutions in Benzene-Dimethyl Formamide." *Turk. J. Chem.*, **1999**, 23, 31-39.
- (31) Avendano, S. C., Hernandez, A. R., Romero-Romo, M. A., Pardave, M. P., Silva, M. T. R. "Study on the stability of nonadrenaline and on the acidity constants." *Spectrochimica Acta A: Molecu. Biomolecu. Spectro.*, **2005**, 61, 3139-3144.
- (32) Ocak, M., Nurhan, G., Ocak, U., Buschmann, H. J., Schollmeyer, E. "The synthesis of new triazole ligands and determination of complex stability constants with transition metal cations in aqueous media." *J. Sol. Chem.*, **2008**, 37, 1489-1497.
- (33) Blanco, S. E., Ferretti, F. H. "Calculations of acidity constants of pyridines in aqueous solution." *Chem. Phys. Lett.*, **2005**, 403, 400-404.
- (34) Oegretir, C., Demirayak, S., Tay, N. F. "Spectroscopic Determination of Acid Dissociation Constants of Some Pyridyl-Substituted 2-Aminothiazole Derivatives." *J. Chem. Eng. Data*, **2006**, 51, 946-951.

- (35) Srour, R. K., McDonald, L. M. "Determination of the acidity constants of methyl red and phenol red indicators in binary methanol- and ethanol-water mixtures." *J. Chem. Eng. Data*, **2008**, 53, 116-127.
- (36) Shivakumar, K., Shashidhar., Halli, M. B. "the potentiometric and thermodynamic properties of some divalent metal ions with biologically active 2-phenyl-3-(2'-hydroxy-5'-methylbenzylidene)-quinazoline-4-(3H)-one Schiff's base." *Russ. J. Phys. Chem.*, **2008**, 82(13), 2269-2276.
- (37) Yang, L., Yuan, Z. "Determination of dissociation constants of amino acids by capillary zone electrophoresis." *Electrophoresis.*, **1999**, 20, 2877-2883.
- (38) Ghasemi, J., Ghobadi, S., Abbasi, B., Kubista, M. "Spectrophotometric determination of acidity constants of group B vitamins in different ionic strengths at 25 °C." *J. Korean. Chem. Soc.*, **2005**, 49, 269-277.
- (39) Aleksic, M., Savic, V., Popovic, G., Buric, N., Kapetavonic, V. "Acidity substituent at C₃ position." *J. Pharma. Biomed. Ana.*, **2005**, 39, 752-756.
- (40) Rivera, A. E. S., Avendano, S. C., Angeles, G. A., Hernandez, A. R., Silva, M. T. R., Romero-Romo, M. A. "Spectrophotometric study on the stability of dopamine and the determination of its acidity constants." *Anal. Chim. Acta.*, **2003**, 499, 199-204.
- (41) Wu, Y. C., Koch, W. F., Berezansky, P. A., Holland, L. A. "The dissociation constant of amino acids by the conductimetric method. I. pK₁ of MOPSO-hydrochloride at 25 °C." *J. Solu. Chem.*, **1992**, 21, 597-605.
- (42) Ghasemi, J., Ghobadi, S., Abbasi, B., Kubista, M. "Spectrophotometric determination of acidity constants of group B vitamins in different ionic strengths at 25 ± 0.1 °C." *J. Korean Chem. Soc.*, **2005**, 49, 269-277.
- (43) Qi, Li., Lin, Y., Jing, X. "The relationship between glass transition temperature and ion dissociation of polymer solid electrolytes." *Huaxue Yanjiu Yu Yingyong.*, **2001**, 13, 137-140.
- (44) Jocelyn, H., Leslie, M. W. "Investigations into the synthesis of 6-Ethyl-5-(4-pyridinyl)-2,4-pyrimidinediamine as a potential antimalarial agent." *J. Heter. Chem.*, **1984**, 21, 741-744.
- (45) Abdel-Rahman, R. M. "Selective reactions with 3-chloro-5,6-diphenyl-1,2,4-triazine." *Commun. Fac. Sci. l'Uni. d'Ankara, Serie B: Chimie*, **1986**, 32(1-2), 87-98.
- (46) Patil, V. D., Wise, D. S., Townsend, L. B. "The synthesis of thieno[2,3-d]pyrimidine nucleosides related to the naturally occurring nucleosides cytidine and uridine." *J. Chem. Soc. Perkin Trans. 1*, **1980**, 1853-1858.
- (47) Bjerrum, J. "Metal Ammine Formation in aqueous solution." P. Hease and Sons, Copenhagen **1941**.

- (48) Irving, H., Rossoti, H. S. "The calculation of formation curves of metal complexes from pH titration curves in mixed solvents." *J. Chem. Soc.*, **1954**, 12, 2094-2010.
- (49) Issa, Y. M., Abdel-Latif, S. A., Abdel-Salam, H. A. "Spectroscopic studies of some mandelic hydrazone derivatives, Modelling." *Measurement & Control.*, **1998**, 57, 1-12.
- (50) El-Sonbati, A. Z., El-Bindary, A. A., Shoair, A. G. F., Younes, R. M. "Stereochemistry of new nitrogen containing heterocyclic aldehyde. VII. Potentiometric, Conductometric and Thermodynamic studies of novel quinoline azodyes and their metal complexes with some transition metals." *Chem. Pharm. Bull.*, **2001**, 49(10), 1308-1313.
- (51) El-Bindary, A. A., El-Sonbati, A. Z., El-Mosalamy, E. H., Ahmed, R. M. "Potentiometric and thermodynamic studies of azosulfonamide drugs." *Chem. Pap.*, **2003**, 57, 255-258.
- (52) Bustamante, S. P., Romero, A. P., Escalera, B., Reillo, A. "Nonlinear enthalpy-Entropy Compensation for the Solubility of Drugs in Solvent Mixtures: Paracetamol, Acetanilide and Nalidixic acid in dioxane-water." *J. Pharma. Sci.*, **1998**, 87, 1590-1596.
- (53) Kalsi, P. S. "Organic reactions and their mechanisms." *New age international (P) limited- New Delhi, 2nd edition*, **2004**, 119.



Chapter-4

Biological Activities

INTRODUCTION

The extensive use of antibiotics has led to the appearance of multi-drug resistant microbial pathogens¹. This highlights the incessant need for the development of new classes of antimicrobial agents and alteration of known drugs in such a way that would allow them to retain their resistance to the pathogen. The design of novel chemotherapeutic agents is particularly beneficial due to their dissimilar mode of action which can avoid cross resistance to known drugs.

To overcome the alarming problem of microbial resistance to antibiotics, the discovery of novel active compounds against new targets is a matter of urgency. Many of the crude drugs, which are sources of medicinal preparations, still originate from wild-growing material. However, plant-based drugs have shortened the life span of the source of material. There is a continuous search for more potent and cheaper raw materials to feed the industry.

The major classes of almost all antibiotics have encountered resistance in clinical applications^{2, 3}. The increasing antibiotic resistance of Gram positive bacteria is becoming a serious problem for human beings⁴. In order to overcome this rapid development of drug resistance, new agents should preferably consist of chemical characteristics that clearly differ from those of existing agents⁵.

In recent years, multiple drug resistance has developed due to indiscriminate use of existing antimicrobial drugs in the treatment of infectious diseases. In addition to this, antibiotics are sometimes associated with adverse effects on the host-like hypersensitivity. Therefore, there is a need to develop alternative antimicrobials drugs for the treatment of infectious diseases from other sources.

Similar groups/structures often exhibit similar biological activities. However, they usually exhibit different potency. The traditional structure activity relationship (SAR) is a useful tool in the search for new drugs. However, SAR is usually determined by making minor changes to the structure of the existing compound and assessing the effect on its biological activity. Similarly, structural analogy has played vital role in designing compounds with higher potency. One of such structural analogy is seen

between 4-aryl-1,4-dihydropyridines (DHPs) of the nifedipine type and dihydropyrimidines (DHPMs).

Dihydropyrimidines belong to an important group of heterocyclic compounds, which are always an attraction point for researchers because of its efficiency towards various pharmacological usages. Dihydropyrimidines have some characteristic properties like manifestations of novel structures, thermal stability, relevant biological properties, high synthesis flexibility and medicinal utility, so a wide range of dihydropyrimidines have been synthesized and extensively studied^{6, 7}. Many researchers have worked on antimicrobial activity of dihydropyrimidine moiety. Fathalla and co-workers⁸ reported the antibacterial and anticancer activity of pyrimidine derivatives. Mishra et al⁹ have reported the antimicrobial activity of 3,4dihydropyrimidinone derivatives. Shinde and Bahekar have reported the anti-inflammatory activity of dihydropyrimidine derivatives.¹⁰

In the past decades, a broad range of biological effects, including antiviral , anti tumor, antibacterial and anti-inflammatory activities has been ascribed to these partly reduced pyrimidine derivatives¹¹. More recently, DHPMs have emerged as orally active antihypertensive agents. A very recent highlight in this context been the identification of the structurally rather simple DHPM monastrol as a mitotic kinesin motor protein inhibitor and potential new lead for the development of anticancer drugs.

In the present chapter, antibacterial activities of the synthesized compounds have been screened against some Gram positive and Gram negative bacteria.

EXPERIMENTAL

The antibacterial activities of all synthesized compounds were studied in DMF and DMSO.

All the synthesized compounds were re crystallized prior to use. The solvents, DMF and DMSO were also purified before use by standard method¹.

For all the compounds, agar well diffusion method was used.

Test Microorganisms:

The synthesized compounds were tested for its antibacterial activity against Gram positive bacteria viz. *staphylococcus aureus* ATCC 25923, *bacillus cereus* ATCC 11778, *bacillus megaterium* ATCC 9885 and Gram negative bacteria viz. *proteus mirabilis* NCIM 2241, *salmonella typhimurium* ATCC 23564, and *klebsiella pneumoniae* NCIM 2719.

Microorganisms were obtained from National Chemical Laboratory (NCL), Pune, India and were maintained at 4°C on nutrient agar slants.

Preparation of test compounds:

The solutions were prepared at a concentration of 1 mg/μl for all the compounds.

Preparation of the plates and microbiological assay:

The antibacterial evaluation was done by agar well diffusion method^{2,3} using Mueller Hinton Agar No.2 as the nutrient medium. The agar well diffusion method was preferred to be used in this study because it was found to be better than the disc diffusion method as suggested by Parekh et al.³ The bacterial strains were activated by inoculating a loop full of test strain in 25 ml of N-broth and the same was incubated for 24 h in an incubator at 37° C. 0.2 ml of the activated strain was inoculated in Mueller Hinton Agar. Mueller Hinton Agar kept at 45°C was then poured in the Petri dishes and allowed to

solidify. After solidification of the media, 0.85 cm ditch was made in the plates using a sterile cork borer and these were completely filled with the test solution. The plates were incubated for 24 h at 37°C. The mean value obtained for the three wells was used to calculate the zone of growth inhibition of each sample. The controls were maintained for each bacterial strain and each solvent. The inhibition zone formed by these compounds against the particular test bacterial strain determined the antibacterial activities of these synthesized compounds.

RESULTS AND DISCUSSION

1, 4 dihydropyrimidinones

Figure 4.1 shows inhibition against Gram positive bacteria in DMSO. It is observed that against *S. aureus*, RVG-7 showed maximum inhibition and RVG-8 showed minimum inhibition. RVG-1, RVG-2, RVG-4, RVG-6, RVG-9 and RVG-10 showed no inhibition at all.

For *B. cereus*, again RVG-7 shows maximum inhibition whereas RVG-9 and RVG-10 exhibited minimum inhibition. RVG-4 and RVG-6 showed no inhibition at all.

Against *B. megaterium*, only RVG-7 shows maximum inhibition whereas RVG-10 exhibited minimum inhibition. RVG-4 showed no inhibition.

Thus, in DMSO *S. aureus* is most resistant bacteria and compound RVG-7 is the most effective compound for the studied bacteria.

Figure 4.2 shows inhibition against Gram positive bacteria in DMF. Compound RVG-7 shows maximum inhibition for all three studied Gram positive bacteria in DMF. For *S. aureus* and *B. cereus*, RVG-4 and RVG-6 shows no inhibition at all whereas in *B. megaterium*, all the compounds show inhibition.

Thus, in DMF also, *B. megaterium* is most susceptible bacteria. In both the solvents, RVG-4 and RVG-6 shows no inhibition in *S. aureus* and *B. cereus*.

The inhibition depends on the solvent, compound structure and bacterial strain. RVG-4 and RVG-6 contain o-methyl benzene and p-fluorobenzene respectively whereas RVG-7 contains 2,5-dichlorobenzene. Thus, the presence of 2,5-dichlorobenzene increases the inhibition in the studied solvents for the above mentioned bacteria.

Against Gram negative bacteria in DMSO, Figure 4.3 shows zone of inhibition for the studied compounds. Again, activity is maximum for RVG-7 all the three studied bacteria. For *K. pneumoniae*, RVG-1, RVG-4, RVG-5 and RVG-9 had no effect. For *P. mirabilis*, RVG-4 and RVG-6 showed no inhibition. Against *S. typhimurium*, only RVG-5 and RVG-7 showed inhibition and other compounds had no inhibition at all.

Figure 4.1: Antibacterial activity of 1, 4-dihydropyrimidinones against Gram positive bacteria in DMSO.




Figure 4.2: Antibacterial activity of 1, 4-dihydropyrimidinones against Gram positive bacteria in DMF.




Figure 4.3: Antibacterial activity of 1, 4-dihydropyrimidinones against Gram negative bacteria in DMSO.





Figure 4.4: Antibacterial activity of 1, 4-dihydropyrimidinones against Gram negative bacteria in DMF.



Thus, for Gram negative bacteria also, maximum is observed by RVG-7 containing 2,5–dichlorobenzene. Whereas, RVG-4 and RVG-6 containing o-methyl and p-fluoro substitution in aromatic ring, had no effect on these bacteria. Thus, in DMSO, RVG-7 is most effective compound. *S. typhimurium* is most resistant bacteria and *P. mirabilis* is the most susceptible bacteria.

Figure 4.4 shows zone of inhibition against Gram negative bacteria in DMF. For all the three bacteria, RVG-7 showed maximum activity. For *K. pneumoniae*, all the compounds show inhibition. For *P. mirabilis*, RVG-10 shows minimum inhibition. RVG-1, RVG-2, RVG-4, RVG-5 and RVG-6 had no effect at all. Against *S. typhimurium*, inhibition is quite less in comparison to *K. pneumoniae* and *P. mirabilis*. However, minimum is observed by RVG-2 whereas RVG-9 showed no inhibition.

Thus in DMF also, *K. pneumoniae* is most susceptible bacteria and compound with -Cl substitution at 3nd and 4th position is most effective.

So overall, in case of RVG series although there is slight change in inhibition in the two solvents, the presence of substituents group compound plays important role in inhibition.

Dihydropyrimidinethiones

Figure 4.5 shows zone of inhibition against Gram positive bacteria in DMSO. It is observed that against *S. aureus*, SRG-1, SRG-2 and SRG-7 are not effective but SRG-6 shows maximum inhibition. For *B. cereus*, all compounds showed inhibition and maximum is observed by SRG-5. For, *B. megaterium*, only SRG-4 had no inhibition and SRG-5 and SRG-6 exhibited maximum inhibition.

So, overall SRG-6 containing p-flouro group in side chain is most effective in DMSO for these Gram positive bacteria and *B. cereus* is most susceptible bacteria.

In DMF, all the compounds exhibited inhibition against all the three Gram positive bacteria as evident from Figure 4.6. Again SRG-6 containing p-flouro group in side chain, showed maximum inhibition against all the three bacterial strains.

Figure 4.5: Antibacterial activity of Dihydropyrimidinethiones against Gram positive bacteria in DMSO.




Figure 4.6: Antibacterial activity of Dihydropyrimidinethiones against Gram positive bacteria in DMF.




Figure 4.7: Antibacterial activity of aminopyrimidines against Gram negative bacteria in DMSO.





Figure 4.8: Antibacterial activity of aminopyrimidines against Gram negative bacteria in DMF.



Comparison of inhibition in DMSO and DMF shows that inhibition is greater in DMF than in DMSO.

Figure 4.7 and 4.8 show zone of inhibition against Gram negative bacteria in DMSO and DMF respectively. In DMSO, SRG-10 showed no activity against *K. pneumoniae* and SRG-5 exhibited maximum inhibition. For *P. mirabilis*, all the compounds show inhibition and SRG-5 and SRG-6 shows maximum inhibition. Against *S. typhimurium*, only SRG-5 showed inhibition. Other compounds had no effect at all. Thus, the presence of o-methoxy (as in SRG-5) and p-flouro (as in SRG-6) increases the inhibition against all the three bacterial strains.

In DMF, for all the three bacterial strains, SRG-5 shows maximum inhibition. SRG-1 and SRG-2 showed no inhibition against *K. pneumoniae*. Whereas, against *P. mirabilis*, inhibition is not shown by SRG-1 and SRG-4. For *S. typhimurium*, SRG-8 showed no activity. Overall, there is not much difference in activity of these compounds against studied bacteria. For all the studied bacteria SRG-6 shows maximum inhibition in both the solvents.

Thus, in this series of compounds, solvent and substitution play an important role in inhibition. Against both Gram positive bacteria, DMF is good solvent where almost all the substituents are effective. SRG-6 contains p-flouro substitution which is found to be most effective in both DMSO and DMF against Gram positive bacteria. Against Gram negative bacteria also, DMF is found to be good solvent.

LITERATURE CITED

- (1) Anderson, I., Van, Scheltinga, A. C., Vallengard, K. *Cell Mol. Life Sci.* **2001**, 58, 1897.
- (2) Rybak, M. J., Akins, R. L. "Emergence of methicillin-resistant staphylococcus aureus with intermediate glycopeptide resistance: clinical significance and treatment options." *Drugs*, **2001**, 61, 1-7.
- (3) Lipsitch, M. "The rise and fall of antimicrobial resistance." *Trends Microbio.* **2001**, 9, 438-444.
- (4) Rapp, R. P. "Antimicrobial resistance in gram-positive bacteria: the myth of the MIC. Minimum inhibitory concentration." *Pharmacotherapy*, **1999**, 19, 112-119.
- (5) Marchese, A., Schilo, G. C., Debbia, E. A. "The evolving threat of antibiotic resistance in Europe: new data from the Alexander Project." *J. Antimicro. Chemo.*, **2000**, 46, 3-9.
- (6) Rakesh, K., Shakshi, M., Ramesh, C. "Synthesis and antimicrobial activity of 4-[5-chloro-3-methyl-1-phenyl-1H Pyrazol-4-yl]-dihydropyrimidines and 4-[5-chloro-3-methyl-1-phenyl-1H Pyrazol-4-yl]-3, 4-dihydropyrimidin-2-ones." *Ind. J. Chem.*, **2009**, 48B, 718-724.
- (7) Venkatesan, J., Pandeya, S. N., Selvakumar, D. "Synthesis and biological evaluation of 4, 6-diaryl substituted-4, 5-dihydro-2-amino pyrimidines." *Ind. J. Pharma. Sci.*, **2007**, 69(4), 586-589.
- (8) Fathalla, O. A., Zeid, I. F., Haiba, M. E., Soliman, A. M., Abd-Elmoez, Sh. I., El-Serwy, W. S. "Synthesis, Antibacterial and Anticancer Evaluation of Some Pyrimidine Derivatives" *World J. Chem.*, **2009**, 4, 127-132.
- (9) Mishra, R., Mishra, B., Hari Narayan Moorthy, N. S. "Synthesis and Antimicrobial evaluation of some 3,4-dihydropyrimidine-2-one derivatives." *Trends in Appl. Sci. Res.*, **2008**, 3, 203-208.
- (10) Sinde, D. B., Bahekar, S. S. "Synthesis and anti-inflammatory activity of some [2-amino-6-(4-substituted aryl)-4-(4-substituted phenyl)-1,6-dihydropyrimidine-5-yl]-acetic acid derivatives." *Acta Pharm.*, **2003**, 53, 223-229.
- (11) Patil, P. A., Bhole, R. P., Chikhale, R. V., Bhusari, K. P. "Synthesis of 3,4-Dihydropyrimidine-2(1H)-one Derivatives using Microwave for their Biological screening." *Int. J. Chem. Tech. Res.*, **2009**, 1, 373-384.
- (12) Riddick, J. A., Bunger, W. B., Sakano, T. "Organic solvents-physical properties and methods of purification." 4th Edi., *Techniques of Chemistry*, II, A Wiley-Interscience Pub., John Wiley, New York, **1986**.

- (13) Perez, C., Paul, M., Bazerque, P. *Acta Biol. Med. Exp.*, **1990**, 15, 113.
- (14) Parekh, J., Inamdhar, P., Nair, R., Baluja, S., Chanda, S. "Synthesis and antibacterial activity of some Schiff bases derived from 4-aminobenzoic acid." *J. Serb. Chem. Soc.*, **2005**, 70, 1155.

*A Comprehensive
Summary of the Work*

A COMPREHENSIVE SUMMARY OF THE WORK

The present work was divided into following chapters:

CHAPTER - 1: This chapter describes the general introduction about the importance of heterocyclic compounds with aims and objective of the present work.

CHAPTER - 2: Part-1 deals with the synthesis of dihydropyrimidinones, dihydropyrimidinethiones, N-methyl pyrimidines and tetrahydropyrimidines bearing pyrimidine moiety whereas in **Part-2** synthesis of Schiff-bases having benzothiazole moiety are described along with their physical constant data. In both the parts I and II, the characterizations of synthesized compounds are done by IR, NMR and mass spectral data. The spectra and the characteristic peak positions of IR and NMR spectra of some compounds are reported. Further, mass spectra and possible fragmentation schemes are given in both parts.

CHAPTER - 3: The physicochemical properties of synthesized dihydropyrimidinones was also studied. The different properties are given in different sections. For the study of all physicochemical properties, DMF and DMSO are used.

Section-I: This section describes the acoustical properties of dihydropyrimidinones in solutions of various concentrations at 298.15 K. The various acoustical parameters helped to understand the different types of interactions occurring in the solutions. It is observed that for both the series, in the studied solvents, solute-solvent interactions dominate.

Section-II: In this section, the densities of dihydropyrimidinones were measured in DMF and DMSO solutions at 298.15 K. The experimental density values are found to be different than those calculated theoretically for all the studied systems, which may be due to solvation of ions in solutions. In solutions of different solvents, density is found to be different due to different

interactions. The molar refraction and refractive index of compounds are also found to be different in each solvent.

Section-III: This section deals with the conductance of studied compounds in solutions of DMF and DMSO at 298.15 K. It is observed that for all the studied compounds, conductivities are less in DMSO than in DMF. Further, all the studied compounds are found to be weak electrolytes in nature.

Section-IV: This section describes the solubility of all the studied compounds in DMF and DMSO at different temperatures (298.15 - 318.15 K). It is observed that the solubility of all the compounds increases linearly with temperature in both the solvents. Comparison of solubility of these compounds in DMF and DMSO shows that overall solubility is greater in DMF than in DMSO. The Gibb's free energy (ΔG_{sol}), enthalpy change (ΔH_{sol}) and entropy (ΔS_{sol}) of different solutions have also been evaluated. ΔH_{sol} and ΔG_{sol} values are found to be positive whereas ΔS_{sol} values are negative. Positive ΔH_{sol} indicates endothermic dissolution of compounds. Whereas positive ΔG_{sol} suggests that the dissolution process is not spontaneous. Further, the negative values of entropy indicate less random ness in solutions.

Section-V: The thermal properties of synthesized dihydropyrimidinones are described in this section. DSC and TGA thermo grams were scanned at the heating rate of 10^0 C per minute. Out of ten studied compounds, multi-step degradation takes place in RVG-3 whereas others decompose by single step. Further, RVG-9 and RVG-10 are found to be most unstable whereas RVG-6 and RVG-8 are most stable followed by RVG-4. All the studied compounds have the same central moiety but different substituents. Thus, stability is affected by different substituent groups. RVG-10 has meta-chloro and para flouro groups whereas RVG-9 has two chloro groups at meta and para positions which decreased the stability. RVG-6 has a flouro group at para positin and RVG-8 has chloro group at meta position. Further, the position of substituent affects stability of the studied compounds.

Further, the melting points determined by DSC and by open capillary methods are found to be in good agreement. The heat of reaction is found to be maximum for RVG-3 and minimum for RVG-4. However, no correlation could be established between heat of reaction, kinetic parameters, melting temperature, thermal stability and substitution groups.

Various kinetic parameters such as order of reaction, energy of activation, frequency factor and entropy change were also calculated for each step. The order of reaction is quite different in different steps for different compounds. Further, the change in entropy values is found to be both positive and negative in different steps. The positive values of entropy change indicate that the transition state is less ordered than the original compound whereas negative value of entropy change corresponds to an increase in the order of transition state than the reactants.

Section-VI: In this section, the dissociation constants of dihydropyrimidinones in water-DMF and water-DMSO mixtures are reported at 298.15 K, 308.15 K and 318.15 K. The dissociation or acidic constant depends not only on the solvent but also on the type of substituent groups present in the compound. Different groups interact differently with the solvent, which affect their dissociation.

Further, the acidity constant value decreases with increasing temperature. Further, acidic constant is minimum in RVG-10 and maximum in RVG-1 suggesting thereby maximum dissociation in RVG-10 which contains fluoride and chloride groups. RVG-1 contains $-\text{OCH}_3$ group which causes a decrease in dissociation. Thus, presence of different substituent groups influences the dissociation of the compound.

Further, some thermodynamic parameters such as enthalpy of solution (ΔH), Gibb's energy change (ΔG) and entropy of solution (ΔS) have also been evaluated from dissociation constants at different temperatures. All these thermodynamic parameters are reported in Table 3.6.5 for both average and half-integral methods. It is observed that for both methods, values are in good agreement with each other.

The positive value of ΔH indicates that dissociation process is endothermic and is accompanied by absorption of heat and favorable at higher temperature. Further, ΔG values are positive indicating thereby that the dissociation process is not spontaneous. However, negative value of ΔS is due to the increased order as a result of the solvation processes except RVG-7.

CHAPTER - 4: The antibacterial activities of the synthesized compounds in DMF and DMSO are explained in this chapter. Different Gram positive and Gram negative bacterial strains behave differently in different solvents. Further, presence of different substituents also affects inhibition.

List of Papers

List of Papers

Published Papers

- (1) "Solubility of Enrofloxacin Sodium in various solvents at various Temperature." S. Baluja, R. Bhalodia, M. Bhatt, N. Vekaria and **R. Gajera**. *J. Chem. Eng. Data*, **2008**, 53, 2897-2899.
- (2) "Solubility of Difloxacin in Acetone, Methanol and Ethanol from (293.15 to 313.15 K)." S. Baluja, **R. Gajera**, R. Bhalodia, N. Vekaria, and M. Bhatt. *J. Chem. Eng. Data*, **2009**, 54, 1091-1093.
- (3) "Synthesis and Ultrasonic Studies of Some Dihydro Pyrimidines in Different Solvents at 298.15 K." **R. Gajera**, R. Bhalodia and S. Baluja. *Int. J. App. Chem.*, **2009**, 5(1), 47-55.
- (4) "Acoustical properties of Schiff bases solutions in DMF." S. Baluja, K. P. Vaishnani, **R. Gajera** and N. Kachhadia. *Latin Amer. Appl. Res. (LAAR)*, 2009.
- (5) "Adiabatic Compressibilities of some synthesized derivatives of dihydro pyrimidines in N, N-Dimethylformamide and Dimethylsulfoxide at 298.15 K." **R. Gajera**, R. Bhalodia and S. Baluja, *Russ. J. Phys. Chem.*, **2010**, 85(5), 1-5.
- (6) "Solubility of Cholesterol in some alcohols from 293.15 to 318.15 K." S. Baluja, **R. Gajera**, N. Vekariya, M. Bhatt and R. Bhalodia. *Scholar Res. Lib., Arc. Apl. Sci. Res.*, **2009**, 1(2) 263-270.

Accepted Papers

- (1) "Solubility of Biologically Active Chalcones in 1,4-Dioxane and N,N-Dimethyl Formamide from (298.15 to 318.15) K." **R. Gajera**, A. Kulshreshtha and S. Baluja. *J. Chem. Eng. Data*, Accepted, **2009**.
- (2) "Synthesis and Acoustical studies of some chalcones of Furaldehyde in different solvents at 308.15 K." **R. Gajera**, N. Vekaria, A. Kulshreshtha and S. Baluja. *Int. J. App. Chem.*, Accepted, **2009**.
- (3) "Solubility of some synthetic Dihydropyrimidines in DMF at (298.15 to 318.15 K)." **R. Gajera**, R. Bhalodia and S. Baluja. *Int. J. Appl. Chem.*, Accepted, **2009**.
- (4) "Solubility of some synthetic dihydropyrimidines in dimethylsulfoxide at (298.15 to 318.15) K." **R. Gajera**, R. Bhalodia and S. Baluja. *Int. J. Pure App. Phys.*, Accepted, **2009**.

- (5) "Synthesis and antibacterial activity of some chalcones." **R. Gajera**, S. Baluja, N. Godvani, S. Chanda, Y. Vaghasiya, J. Parekh, and M. Bhatt. *J. Ind. Chem. Soc.*, Accepted, **2009**.
- (6) "Synthesis, characterization and antibacterial activity of some thiazolidinones." **R. Gajera**, S. Baluja, N. Godvani, S. Chanda, Y. Vaghasiya, J. Parekh, and M. Bhatt. *J. Chem. Pharma. Sci.*, Accepted, **2009**.
- (7) "Solubility of Ofloxacin in 1, 2 dichloromethane, chloroform, carbon tetrachloride and water from (293.15 to 313.15) K" S. Baluja, **R. Gajera**, R. Bhalodia, M. Bhatt and N. Vekariya. *J. Chem. Eng. Data.*, Accepted, **2009**.
- (8) "Density and Refractive index of some Dihydropyrimidines derivatives in DMF and DMSO at 298.15 K." **R. Gajera**, R. Bhalodia and S. Baluja. *Phys. Chem.: An Ind. J.*, Accepted, **2009**.
- (9) "Solubility of 5-amino salicylic acid in different solvents at various temperatures." A. Patel, A. Vaghasiya, **R. Gajera** and S. Baluja. *J. Chem. Eng. Data*. **2010**

Communicated Papers

- (1) "Solubility of a pharmacological intermediate drug Isatin in different solvents at various temperatures." **R. Gajera**, R. Bhalodia, M. Bhatt, N. Vekariya and S. Baluja. *J. Chem. Eng. Data*.
- (2) "Determination of some Thermodynamic parameters of dihydropyrimidine derivatives by dissociation constant." R. Bhalodia, **R. Gajera** and S. Baluja. *Russ. J. Phys. Chem.*
- (3) "Synthesis and antimicrobial screening of 1, 6-dihydropyrimidine derivatives." S. Baluja, **R. Gajera**, A. Kulshrestha, A. Patel, S. Chanda, Y. Vaghasiya, Y. Barvaliya.